prevalence of SJVP was determined. Review of electronic records whether this was mentioned in the CT scan reports by board-certified radiologists was done.

Result Of 1355 patients, 743 have sacroiliac joint vacuum phenomenon. There is a higher prevalence of SJVP in females, 61% and in patients belonging to ≥ 60 years old age group, 43%. Among 743 patients with SJVP, only 4 (0.5%) were reported in the official CT results.

Conclusion SJVP is a prevalent finding and has higher rates in women and in the elderly. The phenomenon is an underreported finding in CT scan examination reports signed out by different board-certified radiologists.

REFERENCES

- 1 Greenspan A, Steinbach LS. Orthopedic Imaging: A Practical Approach. Lippincott Williams Wilkins, 2010.
- 2 Lo SS, Atceken Z, Carone M, Yousen DM. Sacroiliac joint vacuum phenomenon—underreported finding. *Clinical Imaging* 2011;35:465–9.
- 3 Shibata Y, Shirai Y, Miyamoto M. The aging process in the sacroiliac joint: helical computed tomography analysis. *Journal of orthopedic science: Official journal of the Japanese Orthopaedic Association* 2002.
- 4 Stallenberg B, Madani A, Burny F, et al. The vacuum phenomenon: a CT sign of nonunited fracture. *American Journal of Roentgenology* 2001.
- 5 Sze M, Mulligan M. Reliability of vacuum phenomenon in the sacroiliac joint as a sign of traumatic injury. *Emergency Radiology* 2002.
- 6 Fafila CP, Prassopoulos PK, Daskalogiannaki ME, et al. Variation in the appearance of the normal sacroiliac joint on pelvic CT. *Clinical Radiology* 1998;742–6.
- 7 Peh W, Ooi G. Vacuum phenomena in the sacroiliac joints and in association with sacral insufficiency fractures. Incidence and significance. *Spine* 1997.
- 8 Hou Z, Riehl J, Smith W, et al. Severe postpartum disruption of the pelvic ring: report of two cases and review of the literature. *Patient Safety in Surgery* 2011;5:2.

007

SACROILIAC JOINT VACUUM PHENOMENON AMONG PATIENTS WHO UNDERWENT LUMBAR AND ABDOMINOPELVIC CT-SCANS IN A TERTIARY PRIVATE HOSPITAL. A STUDY ON PREVALENCE AND REPORTING RATES

Leiza Joy Saguid Tabora. *Radiology, Cardinal Santos Medical Center, San Juan, Philippines*

10.1136/bmjopen-2015-forum2015abstracts.7

Background Sacroiliac joint vacuum phenomenon (SJVP) is a common incidental finding but its presence is often not included in the official CT scan reports. It is uncommonly found in the literature, and there is lack of data on its prevalence and clinical significance. Determination of SJVP can possibly be used as an indicator in predicting the development or presence of certain diseases and conditions such as infection, degenerative sacroiliac joint diseases, osteonecrosis and other arthropathies.

Objectives The objectives of this study are to determine the prevalence of sacroiliac joint vacuum phenomenon among adult patients who underwent lumbar and abdominopelvic CT scan in a tertiary private hospital for a period of one year and to determine how many patients diagnosed with SJVP were correctly identified based on the official CT scan results.

Methods Multiplanar abdominopelvic and lumbar CT images from January 1, 2013 to December 31, 2013 were retrospectively reviewed for the presence of linear gas density in the sacroiliac joints, known as sacroiliac joint vacuum phenomenon. This included 1355 subjects, (709 males and 646 females) excluding patients less than 20 years old, who seldom undergo CT scan because of the known effects of ionizing radiation. The