

BMJ Open How much sugar is hidden in drinks marketed to children? A survey of fruit juices, juice drinks and smoothies

Jane Boulton,¹ Kawther M Hashem,² Katharine H Jenner,² Ffion Lloyd-Williams,³ Helen Bromley,³ Simon Capewell³

To cite: Boulton J, Hashem KM, Jenner KH, *et al*. How much sugar is hidden in drinks marketed to children? A survey of fruit juices, juice drinks and smoothies. *BMJ Open* 2016;**6**:e010330. doi:10.1136/bmjopen-2015-010330

► Prepublication history and additional material is available. To view please visit the journal (<http://dx.doi.org/10.1136/bmjopen-2015-010330>).

JB and KH are joint first authors.

Received 21 October 2015
Revised 14 December 2015
Accepted 8 January 2016



CrossMark

¹School of Dentistry, University of Liverpool, Liverpool, UK

²Wolfson Institute of Preventive Medicine, Queen Mary, University of London, London, UK

³Department of Public Health and Policy, University of Liverpool, Liverpool, UK

Correspondence to

Dr Ffion Lloyd-Williams; ffionlw@liverpool.ac.uk

ABSTRACT

Objective: To investigate the amount of sugars in fruit juices, juice drinks and smoothies (FJJDS) marketed to children.

Design: We surveyed the sugars content (per 100 ml and standardised 200 ml portion) of all FJJDS sold by seven major UK supermarkets (supermarket own and branded products). Only products specifically marketed towards children were included. We excluded sports drinks, iced teas, sugar-sweetened carbonated drinks and cordials as being not specifically marketed towards children.

Results: We identified 203 fruit juices (n=21), juice drinks (n=158) and smoothies (n=24) marketed to children. Sugars content ranged from 0 to 16 g/100 ml. The mean sugars content was 7.0 g/100 ml, but among the 100% fruit juice category, it was 10.7 g/100 ml. Smoothies (13.0 g/100 ml) contained the highest amounts of sugars and juice drinks (5.6 g/100 ml) contained the lowest amount. 117 of the 203 FJJDS surveyed would receive a Food Standards Agency 'red' colour-coded label for sugars per standardised 200 ml serving. Only 63 FJJDS would receive a 'green' colour-coded label. 85 products contained at least 19 g of sugars—a child's entire maximum daily amount of sugars. 57 products contained sugar (sucrose), 65 contained non-caloric sweeteners and five contained both. Seven products contained glucose-fructose syrup.

Conclusions: The sugars content in FJJDS marketed to children in the UK is unacceptably high. Manufacturers must stop adding unnecessary sugars and calories to their FJJDS.

INTRODUCTION

Free sugars are the most important dietary cause of dental caries.^{1 2} When consumed as a drink, fructose derived from fruit is as likely to cause dental caries as all other sugars.³ In the 2012 Children's Local Dental Health Survey, 28% of children aged 5 years in England had obvious decay in their primary teeth, rising to 35% of children aged 5 years in the North West.⁴ In 2013,

Strengths and limitations of this study

- A comprehensive assessment of the sugars content of fruit juice, juice drinks and smoothies (FJJDS) marketed to children.
- Assessment of standard portion size (200 ml) provided actual amount of sugars contained.
- Only FJJDS available at supermarket chains were recorded.
- New varieties of FJJDS appear in the market steadily and may be different.

12% of children aged 3 years had experienced obvious dental decay.⁵ Dental decay is the most common reason for children in England being admitted to hospital.⁶ Furthermore, an increased intake of fructose and sugar-sweetened drinks has been associated with childhood obesity.^{7 8}

'Free sugars' refer to monosaccharides (such as glucose, fructose) and disaccharides (such as sucrose or table sugar) added to foods and drinks by the manufacturer, cook or consumer, and sugars naturally present in honey, syrups, fruit juices and fruit juice concentrates.⁹ It is a term used to distinguish between the sugars that are naturally present in fully unrefined carbohydrates such as fruit.

Consumption of free sugars, particularly in the form of beverages, can lead to an increase in total sugars intake and a reduction in the consumption of more nutritionally valuable food, thus leading to an unhealthy diet, increased weight and risk of non-communicable diseases.^{10–12} Most recent figures show one in five children aged 4–5 years and one in three children aged 10–11 years are overweight or obese.⁶ Children who are overweight are more likely to develop illnesses such as type 2 diabetes,¹³ have an increased likelihood of weight and health problems in adolescence and are more likely to become overweight or obese adults.¹⁴

Children aged between 4 and 10 years are now getting 30% of their sugars intake from soft drinks.¹⁵ In a recent survey of people's perception of sugars in drinks, the sugars content of fruit juices and smoothies was underestimated by 48% on average, whereas the sugars content of carbonated drinks was overestimated by 12%.¹⁶ Therefore, it is not surprising that the role of sugar-sweetened drinks and fruit juices in the increase in caries and obesity has recently been at the forefront of public debate.¹⁷ Furthermore, there is increasing public awareness of the negative impact of sugar-sweetened drinks on children's health.¹⁸ As a result, parents may replace sugar-sweetened carbonated drinks (that are perceived to be unhealthy) with fruit juices, juice drinks and smoothies (FJJDS) (that are perceived to be healthier). However, this can also have a negative health impact if the sugars content of FJJDS is equal to or higher than sugar-sweetened carbonated drinks.

This study aimed to record the sugars content of FJJDS marketed to children and determine which FJJDS contain sugar (sucrose), syrups, non-caloric sweeteners and free sugars derived from fruit.

Online supplementary appendix 1 provides definitions for the various types of sugars referred to in this paper.

METHODS

This survey looked at sugars per 100 ml and per standardised 200 ml portion of 203 soft FJJDS from seven supermarkets (Tesco, Asda, Sainsbury's, Marks & Spencer, Waitrose, The Co-operative and Morrisons) in the UK. FJJDS were identified by the definitions provided by the British Soft Drinks Association (box 1).

Only products that were specifically marketed towards children were included. Sugar-sweetened drinks, sports drinks and iced teas were excluded. Cordials were also excluded. Although cordials are marketed towards children, they are not single-serve portions that were the focus of this survey. Cordials should be diluted one part cordial to nine parts water (with nutritional typical values shown with this dilution); however, people

Box 1 Fruit juice, juice drinks and smoothies as defined by the British Soft Drinks Association

- ▶ Fruit juices are defined as, '100% pure juice made from the flesh of fresh fruit or from whole fruit, depending on the type used. It is not permitted to add sugars, sweeteners, preservatives, flavourings or colourings to fruit juice'
- ▶ Juice drinks are '1% to 99% juice, nectars, still flavoured waters, sports drinks and iced teas'.^{18a} As neither sports drinks or iced teas are marketed towards children, they were not included in the survey
- ▶ Smoothies labelled as 'fruit juice' should not include any additional ingredients and are subject to the same regulations as fruit juice. Non-pure fruit smoothies may contain other ingredients such as yoghurt or milk, which must be labelled

Table 1 Identification of sugars and sweeteners translated into specific terms from the product ingredients list

Added sugar	Natural sugars	Non-caloric sweeteners
Sugar	Juice	Acesulfame K
Glucose-fructose syrup	Fruit juice from concentrate	Sucralose
		Sodium saccharin

generally dilute to taste, so the sugar content of a single serve would vary from person to person. Furthermore, they are not suitable for lunchboxes.

Only single-portion cartons were recorded. This was determined using the supermarkets' online shopping system and recording products under their children's FJJDS section—for example, 'lunchbox and kids'. For supermarkets with no online shopping, size and design of packaging determined if the products were aimed towards children. Some drinks manufacturers produce products specifically designed for children—for example, Tropicana Kids, only products in these ranges were recorded. Both supermarket own and branded products were recorded.

Product sizes ranged from 65 to 500 ml and were recalculated at 200 ml from the 100 ml data. Teaspoons of sugar per standardised 200 ml portion (calculated using 4 g of sugar per teaspoon)¹⁹ were converted into teaspoons to help the researcher visualise the amount of sugar. The ingredient list was recorded for the analysis of sugar (sucrose), syrups, free sugars derived from fruit and non-caloric sweeteners (table 1).

Data were retrieved online or in-store between July and August 2014.

RESULTS

In total, 203 FJJDS were identified. Of the 203 products, 21 were fruit juices, 158 were juice drinks and 24 were smoothies (table 2).

Of the 179 fruit juices and juice drinks, 60 contained sugar (sucrose). Sugars content ranged from 0 to 16 g/100 ml. The average sugars content was 7.0 g/100 ml, but among the 100% fruit juice category, it was 10.7 g/100 ml. In total 117 of the 203 FJJDS surveyed would receive a Food Standards Agency (FSA) 'red' colour-

Table 2 Number and average sugars content of products included

Category	Number	Mean sugars (g/100 ml)
Juice drinks	158	5.6
100% fruit juice	21	10.7
Smoothies	24	13.0
Total	203	7.0

coded label for sugars for a 200 ml serving, and only 63 FJDS would merit a 'green' colour-coded label.²⁰

Large variations existed in sugars between different types of FJDS and within the same type of FJDS. On average, juice drinks contained the lowest amounts (mean 5.6 g/100 ml) and smoothies contained the highest (mean 13.0 g/100 ml).

Of the 158 juice drinks, 85 contained at least 19 g of sugars—the child's entire maximum daily amount of free sugars.

A total of 60 products contained sugar (sucrose), 65 contained non-caloric sweeteners and 5 contained both. The majority of products were sweetened with acesulfame K, sucralose and aspartame. Of the 78/203 products, 12 contained sugars and non-caloric sweeteners. Of the 203 products, 7 contained glucose-fructose syrup. A total of 140 products contained no sugars or non-caloric sweeteners.

Online supplementary appendix 2 provides the full list of results.

DISCUSSION

The World Health Assembly (WHA) recently adopted a global target of 25% relative reduction in non-communicable diseases by 2025.²¹ This includes policies to reduce the content of sugars in non-alcoholic beverages, as well as to reduce the impact of marketing of non-alcoholic beverages high in sugars on children.²¹ In the UK, a national childhood obesity prevention strategy for England is expected in summer 2016. This is likely to include measures to reduce the consumption of sugar-sweetened carbonated soft drinks.²²

Using a standardised 200 ml serving size, results from this survey show 64% of FJDS surveyed would contain at least half of a young child's maximum sugar intake for the day (9.5 g), with 42% of products containing a full days recommended sugars allowance (19 g for children aged 4–6 years).²³ Current guidelines state a 150 ml glass of pure fruit juice at meal times counts as a maximum of one of the '5 a Day'.²⁴ However, Public Health England is aiming to refresh the '5 a Day' campaign, including a reconsideration of the advice on fruit juice and smoothies.²⁵ This appears crucial, based on our findings. Of the products included in this survey, only five produced products at a 150 ml size. All other products exceeded the recommended serving size and are likely to be consumed in a single seating as a single portion, therefore greatly increasing the child's sugars intake.

Furthermore, the FSA thresholds relate to adult intake of sugars; therefore, we can speculate that the number of FJDS with a 'red' colour code is an underestimation and the number of FJDS with a 'green' colour code is an overestimation for children's sugars consumption.²⁰

WHO recently revised guidelines recommending reducing total daily energy intake from free sugars from 10% to below 5%.¹⁹ This has now been

supported by the government's Scientific Advisory Committee on Nutrition (SACN).²³ SACN recommends that the average population intake of free sugars should not exceed 5% of total dietary energy for age groups from 2 years upwards.²³ This has been endorsed by the UK government. For young children, this 5% limit will represent a maximum of around 19 g/day (nearly five teaspoons of sugar).¹⁹ A recent critical in-depth review goes further, recommending that an intake below 2–3% is necessary to avoid dental caries. The review suggests that an increase from near-zero free sugars to 5% of energy intake doubles the prevalence of caries in children.²⁶ Figures from the latest UK National Diet and Nutrition Survey data show that in children aged 4–10 years, the average intake of free sugars is 14.7% of total energy²⁷—three-fold higher than the new recommendations.

One key difference between whole fruit and juice is fibre content. Whole fruit slows down consumption and has a satiating effect.²⁸ Research shows the body metabolises fruit juice in a different way compared to whole fruit. After whole fruit consumption, the body seems to adjust its subsequent energy intake appropriately, whereas after fruit juice consumption, the body does not compensate for the energy intake.²⁹

We did not initially set out to record the number of products containing non-caloric sweeteners. However, of the 203 FJDS included in our study, 78 contained non-caloric sweeteners, the majority of which were sweetened with acesulfame K, sucralose and aspartame. Although all these non-caloric sweeteners have been declared safe by the European Food Safety Authority,³⁰ health experts believe a reduction in the overall sweetness of products is necessary in order for children's taste to become less accustomed to sweetness.³¹ Although replacing sugars with non-caloric sweeteners may lead to improved markers of long-term metabolic health and weight loss, it will not necessarily provide a long-term solution to reducing sugar consumption and subsequent health problems.

All labels on products recorded contained a reference intake, in line with European law.³² The reference intake provides one figure, which applies to an average-sized adult female doing an average amount of physical activity.³³ This figure is not appropriate for children who have very different energy intake requirements and energy expenditure. Furthermore, pure fruit juices are labelled as Fruit Juice, while products with any other added ingredients are known as Juice Drinks. With all this in mind, it is fair to say that labelling of fruit juice and juice drinks marketed towards children can be unclear and thus potentially confusing for parents trying to make healthier choices.

Study limitations

We acknowledge a number of limitations to this study. First, only FJDS available at supermarket chains were

recorded, but there may be other products available to the public in other smaller independent stores. Second, new varieties of FJDS appear in the market steadily and may have different properties from those recorded. Results, comparisons or conclusions drawn from this survey can thus only apply to the products in this time period (July and August 2014). Third, we only focused on sugars content and did not include an assessment of the portion size of beverages marketed to children, which could be considerably larger than the recommended amount. Fourth, no lactose added to food or drink is now considered as free sugars. However, in the products where some milk was added, we could not identify from the labelling what percentage came from the lactose, therefore it was not excluded. Finally, the study was based on sugars content data provided on the available packaging labels in store on the dates of collection; hence, we relied on the accuracy of the data provided on the label. Therefore, it is assumed that the manufacturers provide accurate and up-to-date information in line with EU Regulations. However, future studies should include free sugars determined through laboratory analysis to achieve a better understanding of the true free sugars content. Nevertheless, the results of this study are relevant and serve to document the sugar content of FJDS products sold in the UK supermarkets, providing a foundation for future studies and evidence for the sugar reduction programme and the soft drink industry.

CONCLUSIONS

The sugars content in FJDS marketed to children in the UK is high. Over 40% of products surveyed contained at least 19 g of sugars—a child's entire maximum daily amount of free sugars.

We suggest that FJDS with high free sugars content should not count as one of the UK government's '5 a Day' recommendations. Ideally, fruit should be consumed in its whole form, not as juice. Parents should dilute fruit juice with water, opt for unsweetened juices and only give them during meals. Portions should be limited to 150 ml a day. In order to help combat the growing problem of childhood obesity, manufacturers need to stop adding unnecessary sugars and calories to their FJDS now; otherwise, it will be essential for the government to introduce legislation to regulate the free sugars content of these products.

Contributors JB and KH conducted the fieldwork. Analysis was done by KH. KH, SC and JB led the project. KH, KJ, HB FL-W and SC helped to design and deliver the project. JB and FL-W wrote the paper with input and approval from all authors.

Funding This research received no specific grant from any funding agency in the public, commercial or not-for-profit sectors.

Competing interests None declared.

Provenance and peer review Not commissioned; externally peer reviewed.

Data sharing statement Additional data are available on request.

Open Access This is an Open Access article distributed in accordance with the terms of the Creative Commons Attribution (CC BY 4.0) license, which permits others to distribute, remix, adapt and build upon this work, for commercial use, provided the original work is properly cited. See: <http://creativecommons.org/licenses/by/4.0/>

REFERENCES

1. Sheiham A. Dietary effects on dental diseases. *Public Health Nutr* 2001;4:569–91.
2. Moynihan PJ, Kelly SA. Effect on caries of restricting sugar intake: systematic review to inform WHO guidelines. *J Dent Res* 2014;93:8–18.
3. Tahmassebi JF, Duggal MS, Malik-Kotru G, *et al*. Soft drinks and dental health: a review of the current literature. *J Dent* 2006;34:2–11.
4. Public Health England. *National Dental Epidemiology Programme for England: oral health survey of five-year-old children 2012. A report on the prevalence and severity of dental caries*. <http://www.nwph.net/dentalhealth/Oral%20Health%205yr%20old%20children%202012%20final%20report%20gateway%20approved.pdf> (accessed 16 Jun 2014).
5. Public Health England. *Dental Public Health Epidemiology Programme Oral health survey of three-year-old children 2013. A report on the prevalence and severity of dental decay*. <http://www.nwph.net/dentalhealth/reports/DPHEP%20for%20England%20OCH%20Survey%203yr%202013%20Report.pdf> (accessed 16 Jun 2014).
6. Health and Social Care Information Centre. National Child Measurement Programme. 2013. <http://www.hscic.gov.uk/ncmp> (accessed 2 Jul 2014).
7. Ludwig DS, Peterson KE, Gortmaker SL. Relation between consumption of sugar-sweetened drinks and childhood obesity: a prospective, observational analysis. *Lancet* 2001;357:505–8.
8. Lustig R. Sugar: The Bitter Truth. 2009. <http://www.youtube.com/> (accessed 2 Jul 2014).
9. Lobstein T. *Reducing consumption of sugar-sweetened beverages to reduce the risk of childhood overweight and obesity*. World Health Organisation, 2014. http://www.who.int/elena/bbc/ssbs_childhood_obesity/en/ (accessed 6 Oct 2015).
10. WHO/FAO. *Report of the joint WHO/FAO expert consultation. Diet, nutrition and the prevention of chronic diseases*. 2013 WHO Technical Report Series 916. <http://www.who.int/mediacentre/news/notes/2014/consultation-sugar-guideline/en/> (accessed 2 Jul 2014).
11. Johnson RJ, Segal MS, Sautin Y, *et al*. Potential role of sugar (fructose) in the epidemic of hypertension, obesity and the metabolic syndrome, diabetes, kidney disease, and cardiovascular disease. *Am J Clin Nutr* 2007;86:899–906.
12. Lobstein T, Brinsden H. Symposium report: the prevention of obesity and NCDs: challenges and opportunities for governments. *Obes Rev* 2014;15:630–9.
13. Haines L, Wan KC, Lynn R, *et al*. Rising incidence of type 2 diabetes in children in the UK. *Diabetes Care* 2007;30:1097–101.
14. Singh AS, Mulder C, Twisk JW, *et al*. Tracking of childhood overweight into adulthood: a systematic review of the literature. *Obes Rev* 2008;9:474–88.
15. The National Diet and Nutrition survey. Results from years 1–4 (2008–2009 and 2011–2012). 2014. <https://www.gov.uk/government/news/new-national-diet-and-nutrition-survey-shows-uk-population-is-eating-too-much-sugar-saturated-fat-and-salt> (accessed 2 Jul 2014).
16. Gill JM, Sattar N. Fruit juice: just another sugary drink? *Lancet Diabetes Endocrinol* 2014;2:444–6.
17. Credit Suisse. Sugar Consumption at a crossroads. 2013. <https://publications.credit-suisse.com/tasks/render/file/index.cfm?fileid=780BF4A8-B3D1-13A0-D2514E21EFFB0479> (accessed 2 Jul 2014).
18. The Times. Health News: Public support grows for tax on sugar. <http://www.thetimes.co.uk/to/health/news/article4584913.ece> (accessed 30 Nov 2015).
- 18a. British Soft Drinks Association, 2014. <http://www.britishtsoftdrinks.com/Soft-Drinks> (accessed 29 Feb 16).
19. World Health Organization. Sugars intake for adults and children: Guideline. 2015. http://www.who.int/nutrition/publications/guidelines/sugars_intake/en/ (accessed 7 Dec 2015).
20. Department of Health and Food Standards Agency. Guide to creating a front of pack (FoP) nutrition label for pre-packed products sold through retail outlets. June 2013. https://www.gov.uk/government/uploads/system/uploads/attachment_data/file/300886/2902158_FoP_Nutrition_2014.pdf#sthash.tlzxHErf.dpu (accessed 7 Dec 2015).

21. World Heart. Federation Follow-up to the Political Declaration of the High-level Meeting of the General Assembly on the Prevention and Control of Non-communicable Diseases. 2014. http://apps.who.int/gb/ebwha/pdf_files/WHA66/A66_R10-en.pdf World (accessed 2 Jul 2014).
22. Action on Sugar. Childhood Obesity Plan. 2014. <http://www.actiononsalt.org.uk/actiononsugar/Press%20Release%20133979.pdf> (accessed 2 Jul 2014).
23. The Scientific Advisory Committee on Nutrition. *Carbohydrates and Health Report*. July 2015. https://www.gov.uk/government/uploads/system/uploads/attachment_data/file/445503/SACN_Carbohydrates_and_Health.pdf (accessed 7 Dec 2015).
24. NHS. 5 A DAY: what counts? 2014. <http://www.nhs.uk/Livewell/5ADAY/Pages/Whatcounts.aspx> (accessed 2 Jul 2014).
25. Public Health England. Sugar reduction responding to the challenge. 2014. http://www.gov.uk/government/uploads/system/uploads/attachment_data/file/324043/Sugar_Reduction_Responding_to_the_Challenge_26_June.pdf (accessed 2 Jul 2014).
26. Sheiham A, James WP. A new understanding of the relationship between sugars, dental caries and fluoride use: implications for limits on sugars consumption. *Public Health Nutr* 2014;17: 2176–84.
27. Bates B, Lennox A, Prentice A, *et al*. National Diet and Nutrition Survey: Headline results from Years 1 to 4 (combined) of the rolling programme from 2008 and 2009 to 2011 and 2012. 2014. <https://www.gov.uk/government/publications/national-diet-and-nutrition-survey-results-from-years-1-to-4-combined-of-the-rolling-programme-for-2008-and-2009-to-2011-and-2012> (accessed 2 Jul 2014).
28. Mattes RD, Campbell WW. Effects of food form and timing of ingestion on appetite and energy intake in lean young adults and in young adults with obesity. *J Am Diet Assoc* 2009;109:430–7.
29. Flood-Obbagy JE, Rolls BJ. The effect of fruit in different forms on energy intake and satiety at a meal. *Appetite* 2009;52:416–22.
30. European Food Safety. Authority Sweeteners. 2014. <http://www.efsa.europa.eu/en/topics/topic/sweeteners.htm> EFSA (accessed 2 Jul 2014).
31. Macgregor G. Soft Drinks Survey. 2014. <http://www.actiononsalt.org.uk/actiononsugar/Sugar%20in%20the%20news/118445.html> (accessed 2 Jul 2014).
32. The Food and Drink Federation. Food labelling: a tool to help improve the food literacy of consumers. 2014. <http://www.gdalabel.org.uk/gda/reference-intakes.aspx> (accessed 2 Jul 2014).
33. NHS. Food Labels. 2014. <http://www.nhs.uk/Livewell/Goodfood/Pages/food-labelling-terms.aspx> (accessed 2 Jul 2014).

Sugar content in fruit drinks marketed to kids “unacceptably high”

Almost half of products assessed contain entire daily recommended maximum intake

The sugar content of fruit drinks—including natural fruit juices and smoothies—is “unacceptably high,” with smoothies among the worst offenders, finds research published in the online journal **BMJ Open**.

Almost half the products assessed contained at least a child’s entire daily recommended maximum sugar intake of 19 g or almost five teaspoons, show the findings, which come ahead of the publication of the UK government’s childhood obesity strategy.

Increasing public awareness of the detrimental effect sugar sweetened drinks have on kids’ teeth and waistlines has prompted many parents to opt for seemingly healthier fruit juice and smoothie alternatives.

To assess the sugar content of fruit juice drinks, 100% natural juices, and smoothies marketed specifically to children, the researchers measured the quantity of ‘free’ sugars per 100 ml in 203 standard portion sizes (200 ml) of UK branded and supermarket own label products, using the pack labelling information provided.

‘Free’ sugars refer to sugars, such as glucose, fructose, sucrose, and table sugar, which are added by the manufacturer, and naturally occurring sugars in honey, syrups, fruit juices and fruit juice concentrates, but not the naturally occurring sugars found in whole fruits and vegetables, which the body metabolises differently and which act to curb energy intake.

The results highlighted wide variations in the amount of free sugars between different types of drink and within the same type of product.

The sugar content ranged from 0-16 g/100 ml, with the average 7 g/100 ml. But sugar content was significantly higher in pure fruit juices and smoothies.

Among the 158 fruit juice drinks analysed, the average sugar content was 5.6 g/100 ml, but rose to 10.7/100 ml among the 21 pure fruit juices tested, and to 13 g/100 ml among the 24 smoothies assessed.

Nevertheless, 85 juice drinks, which add up to more than 40% of the total sample of products, contained at least 19 g of free sugars—a child’s entire daily maximum recommended amount.

Almost 60% of all the products would get a red traffic light label—a coding system designed by the UK’s Food Standards Agency to help consumers identify high levels of fat, salt, and sugar in processed food and drink—the researchers point out.

Some 78 products contained non-calorific sweeteners, such as aspartame. Although safe, health experts believe that a reduction in the overall sweetness of products is required for children’s taste buds to become accustomed to less sugar in their diets, say the researchers.

Current dietary guidelines recommend that a serving of fruit juice/drink/smoothie should be no more than 150 ml, but only six of the products assessed matched this portion size.

Furthermore, the labels on all the products contained a reference intake, which is in line with European law. But this figure applies to an average sized adult woman who does an average amount of physical activity, so wholly inappropriate for children, the researchers emphasise.

They acknowledge that only products available in supermarkets were assessed, that new products constantly appear that may have lower sugar content, and that squashes and cordials were excluded because these are diluted to taste, making portion size comparison difficult.

But based on their findings, they make several recommendations:

- Fruit juices/juice drinks/smoothies with a high free sugar content should not count as one of the UK government's '5 a day' recommendations, as is currently the case
- Fruit should preferably be eaten whole, not as juice
- Parents should dilute fruit juice with water or opt for unsweetened juices, and only allow these drinks during meals
- Portion sizes should be limited to 150 ml/day
- Manufacturers should stop adding unnecessary amounts of sugars, and therefore calories, to their fruit drink/juice/smoothie products—and if they can't do this voluntarily, the government should step in with statutory regulations