BMJ Open Establishing the Aus-ROC Australian and New Zealand out-of-hospital cardiac arrest Epistry

Ben Beck,¹ Janet Bray,^{1,2} Karen Smith,^{1,3,4} Tony Walker,³ Hugh Grantham,^{5,6} Cindy Hein,^{5,6} Melanie Thorrowgood,⁶ Anthony Smith,⁷ Tony Smith,⁸ Bridget Dicker,^{8,9} Andy Swain,¹⁰ Mark Bailey,¹⁰ Emma Bosley,^{11,12} Katherine Pemberton,¹¹ Peter Cameron,^{1,13} Graham Nichol,¹⁴ Judith Finn,^{1,2,4,7} on behalf of the Aus-ROC Steering Committee

To cite: Beck B, Bray J, Smith K, *et al.* Establishing the Aus-ROC Australian and New Zealand out-of-hospital cardiac arrest Epistry. *BMJ Open* 2016;**6**:e011027. doi:10.1136/bmjopen-2016-011027

► Prepublication history and additional material is available. To view please visit the journal (<http://dx.doi.org/10.1136/bmjopen-2016-011027>).

Received 1 January 2016
Accepted 12 February 2016

CrossMark

For numbered affiliations see end of article.

Correspondence to
Professor Judith Finn;
judith.finn@curtin.edu.au

ABSTRACT

Introduction: Out-of-hospital cardiac arrest (OHCA) is a global health problem with low survival. Regional variation in survival has heightened interest in combining cardiac arrest registries to understand and improve OHCA outcomes. While individual OHCA registries exist in Australian and New Zealand ambulance services, until recently these registries have not been combined. The aim of this protocol paper is to describe the rationale and methods of the Australian Resuscitation Outcomes Consortium (Aus-ROC) OHCA epidemiological registry (Epistry).

Methods and analysis: The Aus-ROC Epistry is designed as a population-based cohort study. Data collection started in 2014. Six ambulance services in Australia (Ambulance Victoria, SA Ambulance Service, St John Ambulance Western Australia and Queensland Ambulance Service) and New Zealand (St John New Zealand and Wellington Free Ambulance) currently contribute data. All OHCA attended by ambulance, regardless of aetiology or patient age, are included in the Epistry. The catchment population is approximately 19.3 million persons, representing 63% of the Australian population and 100% of the New Zealand population. Data are collected using Utstein-style definitions. Information incorporated into the Epistry includes demographics, arrest features, ambulance response times, treatment and patient outcomes. The primary outcome is 'survival to hospital discharge', with 'return of spontaneous circulation' as a key secondary outcome.

Ethics and dissemination: Ethics approval was independently sought by each of the contributing registries. Overarching ethics for the Epistry was provided by Monash University HREC (Approval No. CF12/3938—2012001888). A population-based OHCA registry capturing the majority of Australia and New Zealand will allow risk-adjusted outcomes to be determined, to enable benchmarking across ambulance providers, facilitate the identification of system-wide strategies associated with survival from OHCA, and allow monitoring of temporal trends in process and outcomes to improve patient care. Findings will be shared with participating ambulance services and the academic community.

Strengths and limitations of this study

- The Australian Resuscitation Outcomes Consortium (Aus-ROC) epidemiological registry (Epistry) is a population-based out-of-hospital cardiac arrest (OHCA) registry with a capture region encompassing the majority of Australia and all of New Zealand.
- The Epistry provides an ability to define consistent primary and secondary outcome measures to compare the effects of regional and ambulance service variations and understand factors associated with improved survival from OHCA.
- Combining data from multiple regions has inherent challenges, including variation in the variables collected by each of the participating ambulance services.

INTRODUCTION

Out-of-hospital cardiac arrest (OHCA) is a global health problem.¹ Despite over 50 years since the advent of cardiopulmonary resuscitation,² survival remains low with reported rates between 0.6% and 25%.³ Recognition of the variation in OHCA survival rates between regions has heightened interest in the development of multicentre cardiac arrest registries to understand and improve OHCA outcomes.

Individual OHCA registries have been established by ambulance services in most Australian states and across the whole of New Zealand. However, until recently, combining these registries has been hindered by a lack of infrastructure to host the data and to ensure a consensus of definitions and standardisation. As illustrated by the success of international groups, such as the Resuscitation Outcomes Consortium (ROC),⁴ the combining of registries across a larger region facilitates: the understanding of regional differences in the OHCA incidence and outcomes; the

Figure 1 Map of Aus-ROC Epistry highlighting the captured regions in Australia and New Zealand (shown in colour).

influence of different ambulance service structures; and the patient, ambulance service and period factors associated with survival.^{5–7}

AIMS AND OBJECTIVES OF THE PROJECT

The Australian ROC (Aus-ROC)⁸ was established as a National Health and Medical Research Council (NHRMC) Centre of Research Excellence in 2011 to increase research capacity aimed at improving OHCA survival and outcomes. One of the major objectives of this consortium was to develop an Australian and New Zealand cardiac arrest epidemiological registry (Epistry) to enhance and inform coordinated OHCA outcomes, research and evaluation. This paper describes the rationale and methods of the Aus-ROC Epistry.

METHODS

Study design

The Aus-ROC Epistry is designed as a population-based cohort study. Data collection started in 2014. Six established cardiac arrest registries, four in Australia and two in New Zealand, contribute data to the Epistry (figure 1). All six cardiac arrest registries are maintained by

ambulance services: Ambulance Victoria, SA Ambulance Service, St John Ambulance Western Australia, Queensland Ambulance Service, St John New Zealand and Wellington Free Ambulance. Approximately 19.3 million persons are served by these ambulance services (table 1); thus, the Epistry represents approximately 63% of the Australian population (total population of 23.5 million⁹) and 100% of the New Zealand population (total population of 4.5 million).¹⁰

Demographic and socioeconomic characteristics of each of the regions in Australia and New Zealand are shown in online supplementary appendices 2 and 3, respectively.^{17–19} Australian regions participating in the Epistry demonstrated similar characteristics to those of the broader Australian population. While some regions not currently participating in the Epistry demonstrated variations in the proportion of the population who were Aboriginal and Torres Strait Island peoples and those living in ‘major cities’, these regions (Australian Capital Territory, Northern Territory and Tasmania) had small total populations that reflected a combined 4.9% of the Australian population.

Differences in the classification of geographical regions did not facilitate comparisons between Australia and New Zealand. Alternate metrics may be required when making comparisons of rurality between the two countries.

Inclusion criteria

All OHCA attended by ambulance, regardless of aetiology, are included in the Epistry. All deaths attended by ambulance are classified as OHCA and thus included in the Epistry.

Data capture

Capture of OHCA cases varies between ambulance services. A combination of electronic queries of patient care records (PCRs), manual sorting through PCRs, staff standardised reporting and dispatch codes are used to identify OHCA cases. For those ambulance services that use electronic queries, subsequent manual searches are conducted by all services to maximise case capture. Hospital

Table 1 Description of the six participating Australian Resuscitation Outcomes Consortium (Aus-ROC) regions

Region	Ambulance service	Service area population	Geographic area (ha)
South Australia, Australia	SA Ambulance Service	1 685 714 ⁹	98 417 966 ¹¹
Victoria, Australia	Ambulance Victoria	5 841 667 ⁹	22 749 552 ¹²
Western Australia, Australia	St John Ambulance Western Australia	2 573 389 ⁹	252 641 786 ¹³
Queensland, Australia	Queensland Ambulance Service	4 722 447 ⁹	172 582 593 ¹⁴
<i>Australian Registries</i>		14 823 217	121 809 304
New Zealand (excluding Wellington)	St John New Zealand	4 018 370 ¹⁰	26 152 195 ¹⁵
Wellington, New Zealand	Wellington Free Ambulance	491 380 ¹⁰	813 005 ¹⁶
<i>New Zealand Registries</i>		4 509 750	26 965 200
Total		19 332 967	573 357 097

Australia population=23 490 736, Australia geographic area=758 848 540.5 ha,¹² New Zealand population=4 509 750,¹⁰ New Zealand geographic area=26 965 200 ha.¹⁵

outcomes data are commonly obtained through linkage with hospital records and state-based death registries. In one region, hospital outcomes are obtained through linkage with the state Department of Health data.

Variables

Data across all participating sites are collected in accordance with Utstein-style definitions,²⁰ although some variation in the coding of these items exists. Recoding of each variable is conducted to ensure uniformity in the variables collected. Data include information on the OHCA event, demographics, arrest features, response times, treatment and outcome. Variables were identified by consensus among representatives from participating ambulance services (table 2). All variables are collected by each of the six participating ambulance services.

The primary outcome measure is 'survival to hospital discharge', with 'return of spontaneous circulation' as a key secondary outcome. These measures are considered 'core' outcomes according to the Utstein template.²⁰ While neurological outcome and long-term quality of life measurements are important patient-centred outcomes, they are not presently collected across all sites, and thus these measures are not currently included in the Epistry.

Ethical considerations

Ethics approval was independently sought by each of the contributing registries. Overarching ethics for the Epistry was provided by Monash University HREC (Approval No. CF12/3938—2012001888). Data from each of the contributing registries will be provided in reidentifiable form with a unique identifier should the need arise to cross-check individual cases.

Data management and governance

The Epistry is coordinated and located at the Aus-ROC administrative base in the School of Public Health and Preventive Medicine at Monash University, Australia. Each participating ambulance service is responsible for regulating case ascertainment, data capture and quality control. Data upload is web-based and is stored on a secure server at Monash University. Data are then recoded by Epistry staff at Monash University for consistency with the Epistry data dictionary and then combined to form the Epistry.

An Epistry Management Committee (EMC) was established to develop processes and protocols, and oversee all ongoing aspects of the Epistry governance. The committee comprises representatives from each of the contributing ambulance services and Aus-ROC investigators. The Epistry consensus governance document guides the committee on the agreed use of the Epistry data; while maintaining data security and ethical principles.

Statistical analyses

Based on previously reported incidence of OHCA in our region,²¹ we estimate that the Epistry will capture approximately 14 000 patients with OHCA annually. We anticipate

Table 2 Mandatory variables included in the Aus-ROC Epistry

Group	Variable
Case overview	Unique Aus-ROC case identifier Event date Longitude coordinate of event Latitude coordinate of event
Demographics	Age Age qualifier Sex
Arrest features	Location of arrest Witnessed arrest Bystander CPR First arresting rhythm Aetiology of arrest
Dates and times	Date and time call received Date and time arrive at scene (first) Date and time depart scene (transporting ambulance) Date and time arrive hospital (transporting ambulance)
Treatment	Ambulance service resuscitation attempt Ambulance service defibrillation Defibrillation before ambulance service Total number of shocks Airway (final) Mechanical CPR
Outcomes	Any ROSC Scene outcome Survived event (ROSC on hospital handover) Died Discharged alive Date of discharge/death Discharge disposition Receiving hospital (PCI capability)

Aus-ROC, Australian Resuscitation Outcomes Consortium; Epistry, epidemiological registry; PCI, percutaneous coronary intervention; ROSC, return of spontaneous circulation.

using descriptive statistics to summarise baseline characteristics and survival outcomes. Annual crude and age-sex-adjusted incidence rates will be calculated using population figures from the Australian Bureau of Statistics and Statistics New Zealand. Logistic regression will be used to investigate factors associated with key outcome measures with ambulance services as random effects.

Reporting

Publically available reports will be generated on an annual basis, in which ambulance services will be de-identified. As a component of quality improvement, benchmarking reports will be confidentially provided to each of the participating ambulance services comparing their risk-adjusted outcomes with the combined outcomes of the other ambulance services participating in the Aus-ROC Epistry. Findings will also be reported in peer-reviewed publications.

DISCUSSION

The Aus-ROC Epistry was established with the aim of understanding regional, ambulance service and treatment factors associated with improved OHCA survival and outcomes. The scope of the Epistry will enable a range of important research questions to be answered, including understanding regional variations in outcome, the impact of variations in the provision of treatment for OHCA between ambulance services, temporal changes in incidence and outcomes, and the impact of changes in clinical guidelines and clinical trials, as well as increased clinical trial efficiency. The Epistry will allow risk-adjusted outcomes to be determined, to enable benchmarking across providers and identification of system-wide strategies associated with survival for patients with OHCA in Australia and New Zealand.

Variations have been identified in the incidence and outcomes of OHCA across geographical regions in Australia^{21–22} and internationally.^{23–25} To improve outcomes for these patients, there is a need to identify the underlying causes of these variations, which may include differences in the underlying risk of OHCA, and ambulance service structure, function and resuscitation practices; or simply case definitions. For example, other multicentre cardiac arrest registries have investigated regional variation in: the incidence and outcomes of OHCA,^{23–26–27} temporal trends in survival,⁶ rates of bystander cardiopulmonary resuscitation (CPR),²⁸ treatment and transport practices,⁷ and the association between receiving hospital characteristics and survival.²⁹ The Epistry will serve as a framework to address similar questions and identify factors associated with regional variation in survival. The longitudinal nature of the Epistry enables pre-post studies to assess the effect of changes in practice, such as examining whether survival from OHCA improves following the implementation of guideline updates.³⁰ The Epistry will also facilitate the evaluation of clinical trials by providing infrastructure to capture and combine prehospital and in-hospital data across multiple regions in a standardised manner. It is intended to eventually broaden the telephone assessment of quality-of-life (using validated instruments) of OHCA survivors that is conducted by the Victorian Ambulance Cardiac Arrest Registry³¹ and in Aus-ROC clinical trials,³² and use this information to study patient-centred outcomes, as well as economic evaluation of resuscitation interventions.

The Aus-ROC Epistry has been modelled on the successful ROC Epistry.⁴ In comparison to ROC who rely on data from 264 emergency medical service agencies,⁴ one of the strengths of Aus-ROC is the contribution of data from only six ambulance services. Individually, these ambulance services cover large geographical areas and populations, and represent both metropolitan and rural regions, facilitating understanding of variability in the ambulance service systems and the impact that such variability has on patient outcomes. When compared with other multicentre registries that capture smaller proportions of their relevant population, the capture population

of the Epistry represents the majority of Australia and New Zealand and is therefore more likely to be representative of the broader population. With significant heterogeneity in the reporting of OHCA outcomes,³³ another strength of the Aus-ROC Epistry is the ability to define consistent primary and secondary outcome measures to compare the effect of regional and ambulance service variations. Similarly, having a consistent and uniform denominator when defining incidence and survival in OHCA enables valid comparisons across regions.

The sensitive nature of OHCA data, particularly for ambulance services, presents challenges when establishing multicentre cardiac arrest registries.³⁴ These challenges include data ownership, access security and use. In establishing the Epistry, memorandums of understanding were developed between the lead academic institution and each of the participating ambulance services. These agreements, combined with the Epistry governance document, provide clear guidelines on the ownership, use and dissemination of Epistry data.

Combining data from multiple regions has inherent challenges. Variation existed in the variables collected by each of the participating ambulance services, which reduced the number of core variables that were collected for the Epistry. However, it is hoped that over time, all participating sites will be able to collect additional variables, such as 'neurological outcome' and other resuscitation practices and interventions. A survey of the participating ambulance services is currently underway to evaluate variation in the treatment and provision of service between Epistry sites. This survey will also collect data on case ascertainment and quality control measures within each of the participating sites. Aus-ROC also has a vision to extend the current network of participating ambulance services to include all Australian ambulance services. While four Australian ambulance services not currently participating in the Epistry have been invited to contribute data (NSW Ambulance, St John Ambulance Northern Territory, ACT Ambulance, Ambulance Tasmania), these service-based OHCA registries are in various stages of development. It is hoped that these registries will be able to contribute data in the future, and thus the Epistry will capture the prevalence of OHCA in the entire Australian population—as is currently possible for New Zealand.

SUMMARY

The Aus-ROC Epistry is a population-based OHCA registry with a capture region encompassing the majority of Australia and all of New Zealand. Combining OHCA data from six sites will facilitate an understanding of the factors that are associated with improved survival from OHCA and will allow monitoring of temporal trends in process and outcome. Furthermore, the Epistry provides infrastructure to enhance the coordination of clinical trials and improve collaboration between ambulance services, thus increasing capacity to improve outcomes of patients with OHCA in Australia and New Zealand.

Author affiliations

¹Department of Epidemiology and Preventive Medicine, Monash University, Melbourne, Victoria, Australia

²Prehospital, Resuscitation and Emergency Care Research Unit (PRECUR), Curtin University, Perth, Western Australia, Australia

³Ambulance Victoria, Doncaster, Victoria, Australia

⁴Discipline of Emergency Medicine, University of Western Australia, Perth, Western Australia, Australia

⁵Flinders University, Adelaide, South Australia, Australia

⁶SA Ambulance Service, Eastwood, South Australia, Australia

⁷St John Ambulance Western Australia, Perth, Western Australia, Australia

⁸St John, Auckland, New Zealand

⁹Auckland University of Technology, Auckland, New Zealand

¹⁰Wellington Free Ambulance, Wellington, New Zealand

¹¹Queensland Ambulance Service, Brisbane, Queensland, Australia

¹²School of Clinical Sciences, Queensland University of Technology, Brisbane, Queensland, Australia

¹³Emergency and Trauma Centre, The Alfred, Melbourne, Victoria, Australia

¹⁴University of Washington-Harborview Center for Prehospital Emergency Care, University of Washington, Seattle, Washington, USA

Twitter Follow Aus-ROC at @Aus_ROC

Acknowledgements The authors would like to acknowledge the late Professor Ian Jacobs whose contribution to the science of resuscitation is immeasurable. It was Professor Jacob's vision to establish the Aus-ROC Epistry and it is through his hard work and dedication that the Epistry is now operational. The Aus-ROC Epistry will act as one of his many legacies. The authors would also like to thank the registry staff in each of the participating ambulance services for their contribution.

Collaborators Please see online supplementary appendix 1 for a list of Aus-ROC Steering Committee members.

Contributors BB, JB, KS, TW, HG, CH, MT, AS, TS, BD, AS, MB, EB, KP, PC, GN and JF were all involved in the study design. BB drafted the manuscript and all other authors reviewed and approved the manuscript.

Funding The Australian Resuscitation Outcomes Consortium (Aus-ROC) is a National Health and Medical Research Council (NHRC) Centre of Research Excellence (#1029983; <https://www.ausroc.org.au/>). JB is supported by a co-funded NHMRC/National Heart Foundation (NHF) Fellowship (#1069985).

Competing interests None declared.

Ethics approval Monash University Human Research Ethics Committee.

Provenance and peer review Not commissioned; externally peer reviewed.

Open Access This is an Open Access article distributed in accordance with the Creative Commons Attribution Non Commercial (CC BY-NC 4.0) license, which permits others to distribute, remix, adapt, build upon this work non-commercially, and license their derivative works on different terms, provided the original work is properly cited and the use is non-commercial. See: <http://creativecommons.org/licenses/by-nc/4.0/>

REFERENCES

- Ornato JP, Becker LB, Weisfeldt ML, *et al.* Cardiac arrest and resuscitation: an opportunity to align research prioritization and public health need. *Circulation* 2010;122:1876–9.
- Kouwenhoven WB, Jude JR, Knickerbocker G. Closed-chest cardiac massage. *JAMA* 1960;173:1064–7.
- Berdowski J, Berg RA, Tijssen JGP, *et al.* Global incidences of out-of-hospital cardiac arrest and survival rates: Systematic review of 67 prospective studies. *Resuscitation* 2010; 81:1479–87.
- Morrison LJ, Nichol G, Rea TD, *et al.* Rationale, development and implementation of the Resuscitation Outcomes Consortium Epistry-Cardiac Arrest. *Resuscitation* 2008;78:161–9.
- Tijssen JA, Prince DK, Morrison LJ, *et al.* Time on the scene and interventions are associated with improved survival in pediatric out-of-hospital cardiac arrest. *Resuscitation* 2015;94:1–7.
- Daya MR, Schmicker RH, Zive DM, *et al.* Out-of-hospital cardiac arrest survival improving over time: results from the Resuscitation Outcomes Consortium (ROC). *Resuscitation* 2015;91:108–15.
- Zive D, Koprowicz K, Schmidt T, *et al.* Variation in out-of-hospital cardiac arrest resuscitation and transport practices in the Resuscitation Outcomes Consortium: ROC Epistry-Cardiac Arrest. *Resuscitation* 2011;82:277–84.
- The Australian Resuscitation Outcomes Consortium (Aus-ROC). <https://www.ausroc.org.au/> (accessed 18 May 2015).
- Australian Bureau of Statistics. 3235.0—Population by Age and Sex, Regions of Australia, 2014. 2014. <http://www.abs.gov.au/AUSSTATS/abs@.nsf/DetailsPage/3235.02014?OpenDocument> (accessed 6 Jan 2015).
- Statistics New Zealand. New Zealand Subnational Population Estimates. 2015. <http://nzdotstat.stats.govt.nz/wbos/Index.aspx?DataSetCode=TABLECODE7501#> (accessed 18 Aug 2015).
- Australian Bureau of Statistics. South Australia (S/T). 2012. http://stat.abs.gov.au/itt/r.jsp?RegionSummary®ion=4&dataset=ABS_NRP9_ASGS&geoconcept=REGION&measure=MEASURE&datasetASGS=ABS_NRP9_ASGS&datasetLGA=ABS_NRP9_LGA®ionLGA=REGION®ionASGS=REGION (accessed 5 May 2015).
- Australian Bureau of Statistics. Victoria (S/T). 2012. http://stat.abs.gov.au/itt/r.jsp?RegionSummary®ion=2&dataset=ABS_NRP9_ASGS&geoconcept=REGION&measure=MEASURE&datasetASGS=ABS_NRP9_ASGS&datasetLGA=ABS_NRP9_LGA®ionLGA=REGION®ionASGS=REGION (accessed 5 May 2015).
- Australian Bureau of Statistics. Western Australia (S/T). 2012. http://stat.abs.gov.au/itt/r.jsp?RegionSummary®ion=5&dataset=ABS_NRP9_ASGS&geoconcept=REGION&measure=MEASURE&datasetASGS=ABS_NRP9_ASGS&datasetLGA=ABS_NRP9_LGA®ionLGA=REGION®ionASGS=REGION (accessed 5 May 2015).
- Australian Bureau of Statistics. Queensland (S/T). 2012. http://stat.abs.gov.au/itt/r.jsp?RegionSummary®ion=3&dataset=ABS_REGIONAL_ASGS&geoconcept=REGION&measure=MEASURE&datasetASGS=ABS_REGIONAL_ASGS&datasetLGA=ABS_REGIONAL_LGA®ionLGA=REGION®ionASGS=REGION (accessed 9 Jan 2015).
- Statistics New Zealand. New Zealand in Profile: 2014. 2014. <http://www.stats.govt.nz/~media/Statistics/browse-categories/snapshots-of-nz/nz-in-profile/2014/nzip-2014-web.pdf> (accessed 18 Aug 2015).
- Greater Wellington Regional Council. Greater Wellington—About the Region. <http://www.gw.govt.nz/about-the-region/> (accessed 18 Aug 2015).
- Australian Bureau of Statistics. 3235.0—Population by Age and Sex, Regions of Australia, 2011. 2011. <http://www.abs.gov.au/ausstats/abs@.nsf/Products/3235.0-2011~Main+Features~Western+Australia?OpenDocument> (accessed 3 Apr 2015).
- Australian Bureau of Statistics. 3218.0—Regional Population Growth, Australia, 2013–14. 2013. <http://www.abs.gov.au/ausstats/abs@.nsf/mf/3218.0> (accessed 2 Feb 2016).
- Statistics New Zealand. New Zealand: An Urban/Rural Profile Update. 2006. http://www.stats.govt.nz/browse_for_stats/Maps_and_geography/Geographic-areas/urban-rural-profile-update.aspx (accessed 2 Feb 2016).
- Perkins GD, Jacobs IG, Nadkarni VM, *et al.* Cardiac arrest and cardiopulmonary resuscitation outcome reports: update of the Utstein resuscitation registry templates for out-of-hospital cardiac arrest: a statement for healthcare professionals from a task force of the International Liaison Committee on Resuscitation (American Heart Association, European Resuscitation Council, Australian and New Zealand Council on Resuscitation, Heart and Stroke Foundation of Canada, InterAmerican Heart Foundation, Resuscitation Council of Southern Africa, Resuscitation Council of Asia); and the American Heart Association Emergency Cardiovascular Care Committee and the Council on Cardiopulmonary, Critical Care, Perioperative and Resuscitation. *Resuscitation* 2015;96:328–40.
- Nehme Z, Bernard S, Cameron P, *et al.* Using a cardiac arrest registry to measure the quality of emergency medical service care: decade of findings from the Victorian Ambulance Cardiac Arrest Registry. *Circ Cardiovasc Qual Outcomes* 2015;8:56–66.
- Jennings PA, Cameron P, Walker T, *et al.* Out-of-hospital cardiac arrest in Victoria: rural and urban outcomes. *Med J Aust* 2006;185:135–9.
- Nichol G, Thomas E, Callaway CW, *et al.* Regional variation in out-of-hospital cardiac arrest incidence and outcome. *JAMA* 2008;300:1423–31.

24. Ro YS, Shin SD, Song KJ, *et al.* A trend in epidemiology and outcomes of out-of-hospital cardiac arrest by urbanization level: a nationwide observational study from 2006 to 2010 in South Korea. *Resuscitation* 2013;84:547–57.
25. Fosbøl EL, Dupre ME, Strauss B, *et al.* Association of neighborhood characteristics with incidence of out-of-hospital cardiac arrest and rates of bystander-initiated CPR: implications for community-based education intervention. *Resuscitation* 2014;85:1512–17.
26. Hasegawa K, Tsugawa Y, Camargo CA Jr, *et al.* Regional variability in survival outcomes of out-of-hospital cardiac arrest: the All-Japan Utstein Registry. *Resuscitation* 2013;84:1099–107.
27. Ong MEH, Shin SD, De Souza NNA, *et al.* Outcomes for out-of-hospital cardiac arrests across 7 countries in Asia: the Pan Asian Resuscitation Outcomes Study (PAROS). *Resuscitation* 2015;96:100–8.
28. Sasson C, Keirns CC, Smith D, *et al.* Small area variations in out-of-hospital cardiac arrest: does the neighborhood matter? *Ann Intern Med* 2010;153:19–22.
29. Callaway CW, Schmicker R, Kampmeyer M, *et al.* Receiving hospital characteristics associated with survival after out-of-hospital cardiac arrest. *Resuscitation* 2010;81:524–9.
30. Deasy C, Bray JE, Smith K, *et al.* Cardiac arrest outcomes before and after the 2005 resuscitation guidelines implementation: evidence of improvement? *Resuscitation* 2011;82:984–8.
31. Nehme Z, Andrew E, Bernard S, *et al.* Comparison of out-of-hospital cardiac arrest occurring before and after paramedic arrival: epidemiology, survival to hospital discharge and 12-month functional recovery. *Resuscitation* 2015;89:50–7.
32. Deasy C, Bernard S, Cameron P, *et al.* Design of the RINSE Trial: the Rapid Infusion of cold Normal Saline by paramedics during CPR. *BMC Emerg Med* 2011;11:17.
33. Whitehead L, Perkins GD, Clarey A, *et al.* A systematic review of the outcomes reported in cardiac arrest clinical trials: the need for a core outcome set. *Resuscitation* 2015;88:150–7.
34. Gräsner JT, Herlitz J, Koster RW, *et al.* Quality management in resuscitation—towards a European Cardiac Arrest Registry (EuReCa). *Resuscitation* 2011;82:989–94.