

Supplementary File 6: Crude figures for jurisdictions (as of February 2016) #

Jurisdiction (month of launch)		Breast			Colorectal			Lung			Ovarian		
		Responses #	Stage 1 Rate (%)	Stage 2 Rate~ (%)	Responses #	Stage 1 Rate (%)	Stage 2 Rate~ (%)	Responses #	Stage 1 Rate (%)	Stage 2 Rate~ (%)	Responses #	Stage 1 Rate (%)	Stage 2 Rate~ (%)
Manitoba (May 2013)	Patient	384	50.6		274	36.0		219	28.9		61	58.1	
	PCP	215	61.8		139	58.2		100	54.1		40	76.9	
Victoria (Jun 2013)	Patient	489	50.7		472	45.0		241	42.1		136	57.4	
	PCP	333	75.9		289	73.9		107	60.8		77	72.6	
England* (Jul 2013)	Patient	416	32.7	47.2	298	24.6	38.2	273	15.5	28.7	273	26.1	42.9
	PCP	326	79.9		232	79.5		206	76.6		210	70.9	
Northern Ireland* (Aug 2013)	Patient	358	48.4		277	47.6		215	32.4		77	63.1	
	PCP	357	100.0		297	100.0		248	100.0		62	80.5	
Wales* (Oct 2013)	Patient	281	34.2	51.4	314	24.6	43.0	228	12.6	30.9	98	22.3	29.6
	PCP	212	80.3		240	81.4		139	70.6		62	75.6	
Denmark (Oct 2013)	Patient	380	77.9		333	61.8		340	69.4		272	70.1	
	PCP	264	77.6		246	75.7		220	74.6		199	73.2	
Scotland* (Dec 2013)	Patient	418	36.4	48.7	337	31.5	44.5	235	20.6	33.5	140	31.6	45.3
	PCP	412	99.5		333	99.4		164	69.5		91	80.5	
Sweden (Feb 2014)	Patient	298	67.7		320	59.6		214	43.3		N/A		
Ontario (Aug 2014)	Patient	1217	27.2		899	17.6		572	15.5		109	28.3	
	PCP	476	54.1		235	37.5		148	46.0		44	65.7	
Norway[‡] (Sept 2014)	Patient	451	N/A	N/A	358	N/A	N/A	146	N/A	N/A	54	N/A	N/A
	PCP	51	24.8		38	24.5		15	18.8		4	12.9	

~ England, Scotland, Wales and Norway used a two stage recruitment process: patient surveys were sent via primary care practitioners (PCPs) who were asked to forward the pack if they felt it would be appropriate for the patient to participate in this study. 'Stage 1 rate' reflects overall response, 'stage 2 rate' is responses including only patients where the PCP responded to indicate they forwarded the questionnaire.

* Responses in the UK nations include all age groups (18 years and older). All other jurisdictions include responses from patients aged 40 years or more.

Crude figures include all respondents, including ineligible patients (e.g. more than six months after diagnosis)

‡ Norway sent patient surveys to PCPs but did not capture how many packs were sent from PCPs to patients, so it is not possible to work out a response rate.

A summary of patient and primary care recruitment across all jurisdictions (as of February 2016).

These preliminary data illustrate the degree to which the recruitment strategy has been successful.