BMJ Open Investigating the epidemiology of medication errors and error-related adverse drug events (ADEs) in primary care, ambulatory care and home settings: a systematic review protocol

Ghadah Asaad Assiri,^{1,2} Liz Grant,¹ Hisham Aljadhey,² Aziz Sheikh³

To cite: Assiri GA, Grant L, Aljadhey H, *et al.* Investigating the epidemiology of medication errors and error-related adverse drug events (ADEs) in primary care, ambulatory care and home settings: a systematic review protocol. *BMJ Open* 2016;**6**:e010675. doi:10.1136/bmjopen-2015-010675

► Prepublication history and additional material is available. To view please visit the journal (<http://dx.doi.org/10.1136/bmjopen-2015-010675>).

Received 25 November 2015

Revised 19 May 2016

Accepted 14 July 2016

CrossMark

For numbered affiliations see end of article.

Correspondence to

Ghadah Asaad Assiri;
S1373565@ed.ac.uk

ABSTRACT

Introduction: There is a need to better understand the epidemiology of medication errors and error-related adverse events in community care contexts.

Methods and analysis: We will systematically search the following databases: Cumulative Index to Nursing and Allied Health Literature (CINAHL), EMBASE, Eastern Mediterranean Regional Office of the WHO (EMRO), MEDLINE, PsycINFO and Web of Science. In addition, we will search Google Scholar and contact an international panel of experts to search for unpublished and in progress work. The searches will cover the time period January 1990–December 2015 and will yield data on the incidence or prevalence of and risk factors for medication errors and error-related adverse drug events in adults living in community settings (ie, primary care, ambulatory and home). Study quality will be assessed using the Critical Appraisal Skills Program quality assessment tool for cohort and case-control studies, and cross-sectional studies will be assessed using the Joanna Briggs Institute Critical Appraisal Checklist for Descriptive Studies. Meta-analyses will be undertaken using random-effects modelling using STATA (V.14) statistical software.

Ethics and dissemination: This protocol will be registered with PROSPERO, an international prospective register of systematic reviews, and the systematic review will be reported in the peer-reviewed literature using Preferred Reporting Items for Systematic Reviews and Meta-Analyses.

INTRODUCTION

Patient safety is a public concern in health-care systems across the world.¹ The National Coordinating Council for Medication Error Reporting and Prevention defines a medication error as ‘any preventable event that may cause or lead to inappropriate medication use or patient harm, while the medication is in the control of the healthcare professional, patient or consumer’.² Medication errors are therefore any mistakes at any stage of

medication management. Adverse drug event (ADE), on the other hand, is ‘an injury resulting from medical intervention related to a drug’, regardless of whether an error has occurred.³ While almost all medication errors can be prevented, ADEs can be categorised as preventable and non-preventable.¹ Box 1 provides definitions of the key terms employed in this systematic review protocol.

Medication errors and error-related ADEs are common and are responsible for considerable patient harm.¹ More specifically, ADEs can lead to morbidity, hospitalisation, increased healthcare costs and, in some cases, death.⁴ It has been estimated that 5–6% of all hospitalisations are drug-related.^{5 6} With estimates suggesting that ADEs causing hospital admission occur in around 10% of inpatients; approximately half of these ADEs are believed to be preventable.⁷

The cost of drug-related morbidity and mortality was estimated to be \$177.4 billion annually in 2001 in the USA alone.⁸

Medication errors and ADEs are a major problem in all care settings, including home, ambulatory and community settings.¹ Children and adults who suffer from multiple long-term conditions with associated complex drug regimens are particularly at risk.^{9–11}

Systematic reviews focusing on the safety of primary care contexts only have identified studies with vastly different prevalence estimates of the rates of medication errors, these reflect differences in definitions, sampling strategy and populations studied; none of these have investigated the risk factors for medication errors.^{12 13}

Since the release of *To Err is Human: Building a Safer Health System* by the Institute of Medicine,¹⁴ which focused on acute care settings, most patient safety research has

Box 1 Key definitions

- **Adverse drug event (ADE):** Bates *et al*³ define ADE as ‘an injury resulting from medical intervention related to a drug’. Some ADEs are caused by underlying medication errors.
- **Medication error:** The National Coordinating Council for Medication Error Reporting and Prevention defines a medication error as: ‘any preventable event that may cause or lead to inappropriate medication use or patient harm, while the medication is in the control of the healthcare professional, patient, or consumer. Such events may be related to professional practice, healthcare products, procedures and systems, including prescribing; order communication; product labelling, packaging, and nomenclature; compounding; dispensing; distribution; administration; education; monitoring; and use’.² Medication errors can result from any step of the medication-use process: selection and procurement, storage, ordering and transcribing, preparing and dispensing, administration or monitoring.¹
- **Non-prescription drugs:** Medicines that can be sold legally without a drug prescription.
- **Over-the-counter (OTC) drug:** The Food and Drug Administration defines OTC drugs as ‘drugs that have been found to be safe and appropriate for use without the supervision of a healthcare professional such as a physician, and they can be purchased by consumers without a prescription’.²⁴
- **Prescription drug:** Drugs that cannot be sold legally without a prescription.

been carried out on hospitalised patients.^{15 16} Given that patients are increasingly managed in primary, ambulatory and home settings, there is a need to also focus attention on community care contexts.

Prior to undertaking further primary work in this area, it is important to take stock of the current evidence base, reflect on the quality of the evidence, distil key findings that have the potential to provide both estimates on the frequency of medication errors and error-related ADEs, and understand the factors underpinning this important source of preventable harm. We will therefore undertake a systematic review investigating the incidence and prevalence of and risk factors for medication errors and error-related ADEs in community (ie, primary care, ambulatory and home) settings.

Research question

What are the incidence and prevalence of and risk factors for medication errors and error-related ADEs in primary care, ambulatory care and home settings?

METHODS**Design**

We will undertake a systematic review and, if possible, a meta-analysis.

Inclusion criteria**Eligibility criteria****Type of studies**

Population-based cross-sectional and cohort studies will be eligible to estimate the incidence and prevalence of

medication errors and ADEs; these study designs and case-control studies will be eligible to study risk factors for the development of error-related ADEs.

Population

The population of interest will be adults (≥ 18 years) who are dwelling in the community and living in their own homes without home healthcare or nursing at home. These patients may be self-managing, receiving care in primary care and ambulatory settings or any combination of the above.

Exposures

The exposure of interest is prescribed and/or over-the-counter medications.

Outcomes

The outcomes of interest are the incidence and prevalence of medication errors and ADEs, and risk factors for the development of medication errors and error-related ADEs. These errors may have occurred anywhere in the medicines’ management process.¹ We will work with the definitions of medication errors and error-related ADEs employed in individual studies.

Exclusion criteria

1. Studies on illegal substance abuse, herbal products, home healthcare (ie, continuous medical and/or nursing care provided to patients in their own homes), nursing home, hospitalised in-patients or those managed in emergency department settings.
2. Paediatrics (< 18 years).
3. Randomised controlled trials since these cannot be used to reliably assess the incidence and/or prevalence of the outcomes of interest.
4. Existing reviews since the focus is on the primary literature.
5. Studies focusing on specific medication errors or subgroups of populations.
6. Incompletely reported studies, for example, in the form of abstracts.

Search strategy

We will search the following biomedical databases for published research studies: Cumulative Index to Nursing and Allied Health Literature (CINAHL), EMBASE, Eastern Mediterranean Regional Office of the WHO (EMRO), MEDLINE, PsycINFO and Web of Science. These databases will be searched from January 1990 to December 2015; the start date has been chosen to reflect the time when patient safety came into the consciousness of policymakers, professionals and the public.¹⁷ In addition, we will search Google Scholar and contact an international panel of experts to search for unpublished and in progress work. The corresponding author of the eligible articles may be contacted if additional information is needed. The reference list of previous studies will be scrutinised for additional possible

eligible studies. No restriction on the language of publication will be employed.

Detailed search strategies are presented in online supplementary appendix 1.

Study selection

GA will search the databases. GA and a second reviewer will then independently screen the titles and abstracts for eligible studies according to the above detailed selection criteria. Full-text articles will be retrieved from selected studies and will be reviewed according to the selection criteria. Disagreements will be resolved by discussion between the reviewers or arbitration by a third reviewer if a decision cannot be reached. Each copy of the selected studies will be retrieved and the reason for excluding other studies will be clearly noted.

Quality assessment

The risk of bias assessments will be independently carried out on each study by two reviewers using the Critical Appraisal Skills Program quality assessment tool for cohort and case-control studies,¹⁸ and cross-sectional studies will be assessed using the Joanna Briggs Institute Critical Appraisal Checklist for Descriptive Studies.¹⁹ Any disagreements will be resolved by consensus or arbitration by a third reviewer if a decision cannot be reached. Each study will be graded as being at high, medium or low risk of bias.

Data extraction

Data will be extracted by two reviewers and recorded onto a customised data extraction sheet. Discrepancies will be resolved by discussion. The following information will be extracted:

1. Author, year;
2. Study design, study type (retrospective, prospective);
3. Population of interest;
4. Exposure of interest;
5. Outcomes of interest;
6. Main findings;
7. Conclusions;
8. Additional notes.

Data analysis

Data will be summarised in detailed data tables, which will include information on the incidence, prevalence, and relative risk and ORs, together with 95% CIs, for each study (where available). STATA (V.14) statistical software will be used to pool study data if this is considered both clinically and statistically appropriate. Meta-analyses will be undertaken using random-effects modelling.²⁰

Sensitivity analyses will be undertaken by excluding studies judged to be at the highest risk of bias.

Subgroup analyses will be undertaken comparing: adults (18–64 years) versus elderly (≥ 65 years) patients; and those who have recently been an inpatient or had a

hospital visit (<30 days) versus those who have not had a recent hospital attendance (≥ 30 days).

If possible, funnel plots will be used to assess the presence of publication bias.²¹

Registration and reporting

This systematic review will be registered with PROSPERO, an international prospective register of systematic reviews, and will be reported using Preferred Reporting Items for Systematic Reviews and Meta-Analyses²² and Meta-analysis Of Observational Studies in Epidemiology guidelines.²³

DISCUSSION

This systematic review will provide a comprehensive assessment of the epidemiology of medication errors and error-related ADEs in community settings. We anticipate reporting the findings from this study in the autumn of 2016.

Author affiliations

¹Centre for Population Health Sciences, Usher Institute of Population Health Sciences and Informatics, The University of Edinburgh, Edinburgh, UK

²King Saud University, College of Pharmacy, Riyadh, Saudi Arabia

³Centre of Medical Informatics, Usher Institute of Population Health Sciences and Informatics, The University of Edinburgh, Edinburgh, UK

Acknowledgements The authors are grateful to Marshall Dozier for her help with formulating the search strategy, Rachel Faulkner-Jones for proofreading and the Farr Institute.

Contributors GAA conceived the idea for this review. The protocol methods were developed in conjunction with AS and EG. GAA led the drafting of the manuscript and this was commented on critically by AS and EG.

Funding The systematic review protocol is a part of GAA's PhD study with The University of Edinburgh. King Saud University, College of Pharmacy funded the Scholarship. AS is supported by the Farr Institute.

Competing interests None declared.

Provenance and peer review Not commissioned; externally peer reviewed.

Open Access This is an Open Access article distributed in accordance with the Creative Commons Attribution Non Commercial (CC BY-NC 4.0) license, which permits others to distribute, remix, adapt, build upon this work non-commercially, and license their derivative works on different terms, provided the original work is properly cited and the use is non-commercial. See: <http://creativecommons.org/licenses/by-nc/4.0/>

REFERENCES

1. Mark SM LJ, Geller S, Weber RJ. Chapter 5. Principles and practices of medication safety. In: DiPiro JT, Talbert RL, Yee GC, *et al*, eds. *Pharmacotherapy: a pathophysiologic approach*. New York, NY: McGraw-Hill, 2011 (Accessed Oct 10, 2015).
2. What is a Medication Error? National Coordinating Council for Medication Error Reporting and Prevention (cited 16 October 2015). <http://www.nccmerp.org/about-medication-errors>
3. Bates DW, Cullen DJ, Laird N, *et al*. Incidence of adverse drug events and potential adverse drug events. Implications for prevention. ADE Prevention Study Group. *JAMA* 1995;274:29–34.
4. Sorensen L, Stokes JA, Purdie DM, *et al*. Medication management at home: medication risk factor prevalence and inter-relationships. *J Clin Pharm Ther* 2006;31:485–91.
5. Einarson TR. Drug-related hospital admissions. *Ann Pharmacother* 1993;27:832–40.
6. Krähenbühl-Melcher A, Schlienger R, Lampert M, *et al*. Drug-related problems in hospitals: a review of the recent literature. *Drug Saf* 2007;30:379–407.

7. Kongkaew C, Hann M, Mandal J, *et al*. Risk factors for hospital admissions associated with adverse drug events. *Pharmacotherapy* 2013;33:827–37.
8. Ernst FR, Grizzle AJ. Drug-related morbidity and mortality: updating the cost-of-illness model. *J Am Pharm Assoc (Wash)* 2001;41:192–9.
9. Kozer E. Medication errors in children. *Paediatr Drugs* 2009;11:52–4.
10. Zakharov S, Tomas N, Pelclova D. Medication errors—an enduring problem for children and elderly patients. *Ups J Med Sci* 2012;117:309–17.
11. Machado JE, Moncada JC, Mesa G. [Prescription patterns for antilipidemic drugs in a group of Colombian patients]. *Rev Panam Salud Publica* 2008;23:179–87.
12. Olaniyan JO, Ghaleb M, Dhillon S, *et al*. Safety of medication use in primary care. *Int J Pharm Pract* 2015;23:3–20.
13. Panesar SS, deSilva D, Carson-Stevens A, *et al*. How safe is primary care? A systematic review. *BMJ Qual Saf* 2016;25:544–53.
14. Kohn LT, Corrigan JM, Donaldson MS, eds. *To err is human: building a safer health system*. Washington DC: National Academies Press (US) Copyright 2000 by the National Academy of Sciences. All rights reserved, 2000.
15. Sheikh A, Panesar SS, Larizgoitia I, *et al*. Safer primary care for all: a global imperative. *Lancet Glob Health* 2013;1:e182–3.
16. Cresswell KM, Panesar SS, Salvilla SA, *et al*. Global research priorities to better understand the burden of iatrogenic harm in primary care: an international Delphi exercise. *PLoS Med* 2013;10:e1001554.
17. Emanuel L, Berwick D, Conway J, *et al*. What exactly is patient safety? In: Henriksen K, Battles JB, Keyes MA, *et al*, eds. *Advances in patient safety*. New Directions and Alternative Approaches, 2008.
18. Critical Appraisal Skills Programme checklist for cohort studies (cited 10 October 2015). http://www.casp-uk.net/wp-content/uploads/2011/11/CASP_Cohort_Appraisal_Checklist_14oct10.pdfwebcite
19. Joanna Briggs Institute. Checklist for critical appraisal of descriptive studies (cited 16 October 2015). http://joannabriggs.org/assets/docs/jbc/operations/criticalAppraisalForms/JBC_Form_CritAp_DescCase.pdf
20. Borenstein M, Hedges LV, Higgins JPT, *et al*. A basic introduction to fixed-effect and random-effects models for meta-analysis. *Res Synth Methods* 2010;1:97–111.
21. Egger M, Davey Smith G, Schneider M, *et al*. Bias in meta-analysis detected by a simple, graphical test. *BMJ* 1997;315:629–34.
22. Beller EM, Glasziou PP, Altman DG, *et al*. PRISMA for Abstracts Group. PRISMA for Abstracts: Reporting Systematic Reviews in Journal and Conference Abstracts. *PLoS Med* 2013;10:e1001419.
23. Stroup DF, Berlin JA, Morton SC, *et al*. Meta-analysis of observational studies in epidemiology: a proposal for reporting. Meta-analysis Of Observational Studies in Epidemiology (MOOSE) group. *JAMA* 2000;283:2008–12.
24. What are over-the-counter (OTC) drugs and how are they approved? U.S. Food and Drug Administration (cited 19 October 2015). <http://www.fda.gov/AboutFDA/Transparency/Basics/ucm194951.htm>

Correction: *Investigating the epidemiology of medication errors and error-related adverse drug events (ADEs) in primary care, ambulatory care and home settings: a systematic review protocol*

Assiri GA, Grant L, Aljadhey H, *et al.* Investigating the epidemiology of medication errors and error-related adverse drug events (ADEs) in primary care, ambulatory care and home settings: a systematic review protocol *BMJ Open* 2016;6:e010675. doi: 10.1136/bmjopen-2015-010675

The author, Ghadah Asaad Assiri would like to change the affiliation order in this article.

At present, the order of affiliation for Ghadah Asaad Assiri is:

1. Centre for Population Health Sciences, Usher Institute of Population Health Sciences and Informatics, The University of Edinburgh, Edinburgh, UK.
2. King Saud University, College of Pharmacy, Riyadh, Saudi Arabia.

The revised order of affiliation is:

1. King Saud University, College of Pharmacy, Riyadh, Saudi Arabia.
2. Centre for Population Health Sciences, Usher Institute of Population Health Sciences and Informatics, The University of Edinburgh, Edinburgh, UK.

As per the revised order, the affiliation for Liz Grant and Hisham Aljadhey is

2. Centre for Population Health Sciences, Usher Institute of Population Health Sciences and Informatics, The University of Edinburgh, Edinburgh, UK.
1. King Saud University, College of Pharmacy, Riyadh, Saudi Arabia.

Open access This is an open access article distributed in accordance with the Creative Commons Attribution Non Commercial (CC BY-NC 4.0) license, which permits others to distribute, remix, adapt, build upon this work non-commercially, and license their derivative works on different terms, provided the original work is properly cited, appropriate credit is given, any changes made indicated, and the use is non-commercial. See: <http://creativecommons.org/licenses/by-nc/4.0/>.

© Author(s) (or their employer(s)) 2019. Re-use permitted under CC BY-NC. No commercial re-use. See rights and permissions. Published by BMJ.

BMJ Open 2019;9:e010675corr1. doi:10.1136/bmjopen-2015-010675corr1