

Supplementary Materials.

Table A. Mental health and wellbeing 9 months after the birth by current pregnancy status

	Held (%)	Did not hold (%)	OR (95%CI)	OR _{adj} (95%CI)
Currently Pregnant (188)				
Depression	19.5 (32)	4.2 (1)	5.16 (0.67-39.78)	5.00 (0.65-38.74)
Anxiety	29.9 (49)	20.8 (5)	1.62 (0.57-4.61)	1.58 (0.55-4.52)
Poor physical health	3.0 (5)	8.3 (2)	0.57 (0.06-5.35)	0.27 (0.02-3.19)
PTSD symptoms	36.0 (59)	12.5 (3)	3.76* (1.07-13.21)	3.75* (1.07-13.20)
Relationship difficulties with partner	3.7 (6)	0.0 (0)	‡	‡
Relationship difficulties with family	7.9 (13)	0.0 (0)	‡	‡
Not Pregnant (270)				
Depression	30.2 (67)	22.9 (11)	1.41 (0.68-2.95)	1.25 (0.58-2.69)
Anxiety	35.1 (78)	16.7 (8)	2.65* (1.18-5.96)	2.26 (0.98-5.21)
Poor physical health	5.4 (12)	6.2 (3)	0.84 (0.23-3.09)	0.93 (0.23-3.74)
PTSD symptoms	40.1 (89)	29.2 (14)	1.58 (0.79-3.12)	1.19 (0.58-2.47)
Relationship difficulties with partner	11.7 (26)	12.5 (6)	0.90 (0.35-2.34)	0.80 (0.30-2.17)
Relationship difficulties with family	17.6 (39)	4.2 (2)	4.79*(1.12-20.61)	3.68 (0.84-16.12)

OR_{adj}: with adjustment for multiple pregnancy, ethnicity and fertility treatment

‡ Group sizes insufficient for analysis, *p<.05, **p<.01, ***p<.001

Table B. Mental health and wellbeing 9 months after the birth by condition of the baby (time between antepartum death and delivery, gestation of stillbirth, and presence of a congenital anomaly)

	Held % (n)	Did not hold % (n)	OR (95%CI)	OR _{adj} (95%CI)	Held % (n)	Did not hold % (n)	OR (95%CI)	OR _{adj} (95%CI)
Time between antepartum death and delivery					A few hours before or during delivery (n = 97)			
Up to the day before (or unsure) (n = 357)								
Depression	24.3 (71)	16.9 (11)	1.52 (0.75-3.07)	1.49 (0.73-3.04)	30.4 (28)	0 (0)	‡	‡
Anxiety	30.8 (90)	16.9 (11)	2.19* (1.09-4.40)	2.04* (1.01-4.13)	40.2 (55)	40.0 (2)	1.03 (0.16-6.46)	1.08 (1.52-7.69)
PTSD symptoms	36.3 (106)	21.5 (14)	2.02* (1.07-3.84)	1.76 (0.91-3.39)	45.7 (42)	20.0 (1)	‡	‡
Poor physical health	4.1 (12)	6.2 (4)	0.81 (0.22-2.98)	0.93 (0.24-3.64)	5.4 (5)	0 (0)	‡	‡
Relationship difficulties with partner	8.6 (25)	7.7 (5)	1.12 (0.41-3.05)	1.07 (0.39-2.94)	10.9 (10)	20.0 (1)	‡	‡
Relationship difficulties with family	14.4 (42)	1.5 (1)	‡	‡	14.1 (13)	20.0 (1)	‡	‡
Gestation of stillbirth					33 weeks' + (n = 289)			
< 33 weeks gestation (n = 165)								
Depression	22.6 (31)	7.1 (2)	3.82 (0.86-17.05)	5.21* (1.07-25.28)	27.9 (69)	23.8 (10)	1.19 (0.55-2.56)	1.08 (0.49-2.38)
Anxiety	29.9 (41)	7.1 (2)	5.86* (1.33-25.85)	7.52* (1.61-35.17)	35.2 (87)	26.2 (11)	1.50 (0.72-3.14)	1.31 (0.61-2.80)
PTSD symptoms	31.4 (43)	21.4 (6)	1.71 (0.65-4.53)	1.75 (0.64-4.78)	42.9 (106)	26.2 (11)	2.05 (0.98-4.27)	1.69 (0.79-3.62)
Poor physical health	5.8 (8)	7.1 (2)	0.84 (0.17-4.18)	1.46 (0.25-8.65)	3.6 (3)	7.1 (9)	0.66 (0.14-3.22)	0.69 (0.13-3.60)
Relationship difficulties with partner	5.8 (8)	3.6 (1)	1.74 (0.21-14.51)	1.50 (0.17-13.04)	10.9 (27)	11.9 (5)	0.89 (0.32-2.47)	0.75 (0.27-2.12)
Relationship difficulties with family	13.9 (19)	0 (0)	‡	‡	14.6 (36)	4.8 (2)	3.36 (0.78-14.53)	3.00 (0.69-13.16)
Congenital Anomaly					No congenital anomaly (n = 392)			
Congenital Anomaly (n = 59)								
Depression	29.2 (14)	18.2 (2)	1.63 (0.30-8.70)	1.60 (0.26-10.02)	24.8 (82)	16.4 (10)	1.65 (0.80-3.41)	1.51 (0.72-3.15)
Anxiety	27.1 (13)	18.2 (2)	1.63 (0.30-8.70)	1.32 (0.23-7.72)	32.9 (109)	18.0 (11)	2.20* (1.10-4.41)	2.03* (1.01-4.12)
PTSD symptoms	31.3 (15)	18.2 (2)	1.81 (0.34-9.63)	1.94 (0.34-11.10)	38.7 (128)	24.6 (15)	1.91* (1.02-3.57)	1.67 (0.88-3.17)
Poor physical health	2.1 (1)	0(0)	‡	‡	3.9 (13)	8.2 (5)	0.57 (0.18-1.82)	0.62 (0.19-2.07)
Relationship difficulties with partner	4.2 (2)	18.2 (2)	0.19 (0.02-1.52)	0.11 (0.01-1.07)	10.0 (33)	6.6 (4)	1.55 (0.53-4.56)	1.48 (0.50-4.44)
Relationship difficulties with family	10.4 (5)	0 (0)	‡	‡	14.5 (48)	3.3 (2)	4.94* (1.17-20.91)	4.64* (1.09-19.84)

OR_{adj}: with adjustment for multiple pregnancy, ethnicity and fertility treatment ‡ Group sizes insufficient for analysis, *p<.05, **p<.01, ***p<.001