BMJ Open Vitamin B₁₂ status in women of childbearing age in the UK and its relationship with national nutrient intake guidelines: results from two National Diet and Nutrition Surveys

Nithya Sukumar,^{1,2} Antonysunil Adaikalakoteswari,¹ Hema Venkataraman,^{1,2} Hendramoorthy Maheswaran,¹ Ponnusamy Saravanan^{1,2}

To cite: Sukumar N, Adaikalakoteswari A, Venkataraman H, *et al.* Vitamin B₁₂ status in women of childbearing age in the UK and its relationship with national nutrient intake guidelines: results from two National Diet and Nutrition Surveys. *BMJ Open* 2016;**6**:e011247. doi:10.1136/bmjopen-2016-011247

► Prepublication history and additional material is available. To view please visit the journal (<http://dx.doi.org/10.1136/bmjopen-2016-011247>).

NS and AA contributed equally.

Received 22 January 2016
Revised 22 May 2016
Accepted 14 June 2016

CrossMark

¹Warwick Medical School, University of Warwick, Coventry, UK

²Academic Department of Diabetes and Metabolism, George Eliot Hospital, Nuneaton, UK

Correspondence to

Dr Ponnusamy Saravanan;
P.Saravanan@warwick.ac.uk

ABSTRACT

Objective: To assess serum B₁₂, folate and the associated homocysteine (Hcy) levels among women of childbearing age in the UK and examine their association with dietary intake in relation to the UK Recommended Nutrient Intakes (RNIs) for B₁₂ and folate.

Design: Cross-sectional study.

Setting: Data from two publicly available National Diet and Nutrition Surveys (NDNS 2000/2001 and 2008/2012) were used. These were population-based surveys of randomly selected samples of adults which were carried out in their households.

Participants: Women of childbearing age (aged 19–39 years), representative of the UK population. Those who were pregnant or breastfeeding were excluded.

Outcome measures: The associations between micronutrient intakes and blood levels of B₁₂, folate and Hcy were assessed by correlation and stepwise linear regression. B₁₂ intake was divided into quintiles and plotted against blood B₁₂ and Hcy concentrations to determine the threshold of any associations.

Results: 299 women from the first NDNS cohort had complete intake and biomarker data. The prevalence of serum vitamin B₁₂ (≤ 150 pmol/L) and serum folate (≤ 10 nmol/L) deficiency and hyperhomocysteinemia (≥ 12 μ mol/L) was 12.4%, 6.4% and 21.2%, respectively, despite seemingly adequate B₁₂ intakes (median 3.8 μ g/day, 96% consumed more than the UK RNI of 1.5 μ g/day). B₁₂ concentrations increased across all quintiles of intake with serum levels in quintiles 4 and 5 (median intake 4.9 and 7.1 μ g/day, respectively) significantly higher than quintile 1. However, Hcy concentrations levelled off between quintiles 4 and 5. Comparison of micronutrient intake between the two surveys found that folate intake has reduced in the more recent cohort.

Conclusions: The UK RNI for B₁₂ intake should be increased for women of childbearing age with intakes of around 5–7 μ g/day likely to be associated with stable biomarker levels. B₁₂ levels should also be measured in women preconceptionally or in early pregnancy given the high rates of deficiency.

Strengths and limitations of this study

- Two publicly available data sets from the British National Diet and Nutrition Surveys (NDNS) were used to investigate the association between dietary intake of B₁₂ and folate and their blood concentrations in women of childbearing age.
- The women sampled in the surveys were representative of UK adults; therefore, the findings can be generalised to the wider UK population.
- Information about the participants, including demographics, dietary intakes, blood results and anthropometry, was used for detailed analysis of the outcomes of interest.
- The availability of two separate NDNS data sets allowed comparison of micronutrient intakes over the last 15 years in young women.
- Blood micronutrient concentrations were not analysed in the more recent survey. Therefore, the associations observed in the first survey could not be replicated.

INTRODUCTION

Vitamin B₁₂ (B₁₂), also known as cobalamin, is a micronutrient essential for cellular growth, differentiation and development.¹ Along with folate, B₁₂ is necessary for the synthesis of DNA, RNA, lipids and protein, and an essential step in this process is the conversion of homocysteine (Hcy) to methionine.² Therefore, a deficiency of either B₁₂ or folate leads to increased Hcy which can have significant clinical implications such as cardiovascular disease and atherosclerosis in adults.^{3,4}

During pregnancy, low circulating levels of B₁₂ or folate have been associated with complications such as neural tube defects (NTDs),⁵ spontaneous abortion,⁶ premature birth⁷ and possibly low birth weight.⁸

Mandatory folic acid fortification in North America has resulted in reduction of NTD by over 40% in the last 10 years,⁹ but there has been a tripling of the condition attributable to B₁₂ deficiency during the same time frame in this population.¹⁰ Maternal hyperhomocysteinaemia has been linked to early pregnancy losses,¹¹ pre-eclampsia¹² and small for gestational age babies.¹³ Although the exact mechanisms are not known, some of these effects may be mediated by vascular compromise to the fetus and insufficient placental development which occurs in the early embryonic period.¹⁴ As around half of the pregnancies are unplanned, even in developed countries,¹⁵ there is increasing attention on optimising the nutrient status of women in the periconceptional period.

B₁₂ deficiency is prevalent during pregnancy as shown by a large systematic review recently conducted by our group.⁸ We showed that the rates were in the order of 20–30% in all three trimesters across the world, with particular high rates in studies from the Indian subcontinent, Eastern Mediterranean and South American regions.^{16–18} A longitudinal study has shown that cobalamin levels can decrease by around 10–20% from preconception to the first and second trimesters, respectively.¹⁹ However, there are no studies in the UK that assessed the B₁₂ intake or circulating levels in women of reproductive age prior to conception.

The UK Department of Health (DOH) has stipulated that the Recommended Nutrient Intake (RNI) for B₁₂ and folate in adults is 1.5 and 200 µg per day, respectively, with no different recommendations for women (pregnant or otherwise) and the elderly.²⁰ These recommendations were published in 1991 and state that the RNI for B₁₂ 'represent the level of intake considered likely to be sufficient to meet the requirements of 97.5% of the group'.²¹ The evidence for this RNI is from reports in the 1970s, which estimated the average requirement to prevent diet-related B₁₂ deficiency and megaloblastic anaemia.^{22–24} B₁₂ deficiency in these studies was arbitrarily defined as <150 pmol/L without use of any functional markers of B₁₂ deficiency.^{25–26} Thus, there are little contemporary data to support how accurately the recommended intakes correlate with serum values of B₁₂ and functional indicators such as Hcy. The primary aim of this study is to determine the serum B₁₂, folate and Hcy status of women of childbearing age in the UK and to assess the correlation between estimated B₁₂ intake and blood concentrations of B₁₂ and Hcy using the data from two British National Diet and Nutrition Survey (NDNS) in 2000/2001 and 2008/2012. The secondary aim is to compare the nutritional intake between the two NDNS cohort to provide more recent data on B₁₂ intake.

METHODS

Subjects

We used data collected from the two British NDNS between July 2000 and June 2001 and 2008–2012. The

surveys provide detailed quantitative assessment of nutritional status and laboratory results of the participants. The methods used in the survey have previously been described in detail.^{27–29} The samples were made up of randomly selected adults aged 19–64 years living in private households, who were representative of the UK adult population. Their selection was done by a multi-stage, stratified, probability sampling with postal sectors as first stage units. If there were more than one adult in the same household, one was selected randomly. Women who were pregnant or breast feeding were excluded.

The selection of the participants included in our analysis is provided in online supplementary figures S1 and S2. For the first survey (2000/2001), the fieldwork was carried out over a 12-month period, with respondents being surveyed over four 3-month periods to account for seasonal variations in nutritional behaviour and content. Out of 3704 potentially eligible adults identified for the study, 299 women between 19 and 39 years of age (known as women of 'childbearing age' for the purpose of our analysis) with complete dietary and biomarker information were included in the final analysis. In an independent study,³⁰ no evidence of was found of non-response bias in this NDNS data.

The second survey (2008/2012) consisted of 2424 adults of whom 395 were women of childbearing age. Unfortunately, blood samples were only obtained from 157 of these women (ie, 39.7% of all females aged 19–39 years). The main reasons for this low return of blood samples were 'no nurse visit', 'participant refused' and 'blood sample inapplicable'. Owing to the potential bias from a sample of women not representative of the larger NDNS cohort, we only included and compared the micronutrient intakes from the two surveys to assess the secular trend.

Assessment of dietary intake

This has been described in detail elsewhere.²⁷ Briefly, dietary assessment for each participant was done by a two-stage process: (1) two face-to-face interviews using computer-assisted personal interviewing methods, and (2) a 7-day dietary record using weighed food diaries. Each participant was provided with a set of Soehnle Quanta digital food scales and two recording diaries (for use at home and outside). Additional information was obtained about the use of medicines, vitamin and mineral supplements. Information provided in the food diaries was later used to determine nutrient intakes by linking to the Food Standards Agency nutrient database, which holds details for 56 nutrients for each of 6000 food codes.²⁷

Laboratory methods

Venous blood samples were taken at the non-fasting state by trained phlebotomists in participants' homes. Serum folate and B₁₂ measurements were performed on the Abbott IMx semiautomated analyser, which uses microparticle enzyme immunoassay (MEIA) technology

(z). Quality control consisted of an internal pooling of serum samples with each run for use as a drift control, and an external quality assessment by the UK National External Quality Assessment Service (NEQAS). Outlying results were defined as serum B₁₂ level >1000 pmol/L and serum folate >60 nmol/L and were excluded. Plasma Hcy measurements were performed by the Abbott IMx assay on the IMx analyser. Quality control consisted of participation in an international external quality assessment scheme based in Denmark³¹ and by the manufacturer's QV samples.

Statistical analysis

Statistical analysis was performed with SPSS V.22.0 (IBM SPSS Statistics for Windows, Version 22.0 (program). Armonk, NY: IBM Corp, Released 2013). For continuous variables (eg, mean B₁₂/folate intakes and levels between RNI threshold groups), the Student's t-test and for categorical variables (eg, B₁₂ and folate deficiency rates), the χ^2 test for independence or the Fisher's exact test was used. Stepwise multiple linear regression analysis with each of serum B₁₂, folate and plasma Hcy as the dependent variable was done, with predictors entered or removed following the criteria: probability of F to enter ≤ 0.050 , probability of F to remove ≥ 0.100 . The regression models included the following covariates: age, body mass index (BMI), total cholesterol:high-density lipoprotein (TC:HDL) ratio, alcohol intake, smoking status, oral contraceptive use, vegan/vegetarian, serum folate and where appropriate daily folate intake (with supplements), daily B₁₂ intake (with supplements) and serum B₁₂ levels. Race was not included in the analysis as the majority of the participants (95%) were white. To determine the trend of B₁₂ and Hcy across the spectrum of B₁₂ intakes, we divided the cohort into quintiles of B₁₂ intake. One-way analysis of variance (ANOVA) with Tukey's post hoc test was applied to compare the serum levels against the lowest quintile. Logistic regression was then done for incremental B₁₂ intake thresholds of 0.25 μg to determine predictors of B₁₂ deficiency and hyperhomocysteinemia. In order to estimate the prevalence of inadequate B₁₂ intake in our population and observe how this was related to abnormal biomarker levels, we used the Estimated Average Requirement (EAR) cut-point method.³² The EAR was calculated as 1.25 $\mu\text{g/day}$ based on a coefficient of variation of 10% below the UK RNI.²⁵

The definitions for micronutrient deficiencies were as follows: serum B₁₂ ≤ 150 pmol/L,³³ serum folate ≤ 10 nmol/L³⁴ and red cell folate < 350 nmol/L.³⁴ The upper limit of normal for plasma Hcy in non-pregnant adults aged 15–65 has been variably defined as 12 $\mu\text{mol/L}$ in non-pregnant adults with folic acid fortification or supplementation³⁵ and 15 $\mu\text{mol/L}$ in those without fortification or supplementation, which applies to our population. However, Hcy levels above 10.7 $\mu\text{mol/L}$ in women during preconception has been associated with pre-eclampsia, prematurity and very low

birthweight infants if they become pregnant.¹² In addition, other experts recommend < 12 $\mu\text{mol/L}$ for all adults.³⁶ Therefore, we have presented results for 12 and 15 $\mu\text{mol/L}$ and stated these clearly throughout.

RESULTS

From the NDNS 2000/01 cohort, 299 women of childbearing age had complete dietary and serum micronutrient level results. The demographics and clinical characteristics of these women are shown in table 1. Their mean age was 31.6 years and BMI 25.3 kg/m². To determine how the B₁₂/folate intakes women of childbearing age compared with older women from the same NDNS survey, we analysed these parameters in women aged 40–64 years. The younger women consumed significantly less B₁₂ and folate (median 3.83 vs 5.16 and 237 vs 279 $\mu\text{g/day}$, respectively, $p < 0.001$ for both) and consisted of more vegetarians/vegans (8.7% vs 4.2%, $p < 0.05$) and less B₁₂ supplement users (10.4% vs 16.2%, $p < 0.05$). Folic acid supplement use was the same (data not shown).

The blood levels of B₁₂, folate and Hcy in women of childbearing age and according to categories of UK RNI for B₁₂ intake (adequate/inadequate) are given in table 2. The median serum B₁₂ concentration was 241 pmol/L. Overall, 12% of women were B₁₂ deficient (< 150 pmol/L), despite the median B₁₂ intake of the deficient women being nearly two times the UK RNI (2.96 $\mu\text{g/day}$; data not shown). In total, 3.7% of the surveyed population had B₁₂ < 150 pmol/L and Hcy > 12 $\mu\text{mol/L}$, with a significantly higher proportion having the combination of

Table 1 Demographics and B₁₂ and folate intakes of women of childbearing age (NDNS 2000/2001 cohort)

Female 19–39 years	All subjects (n=299)
Age (years)	31.6 $\pm$ 5.7
BMI (kg/m ²)	25.4 $\pm$ 5.2
Obesity, n (%)	46 (15.7)
Current smokers, n (%)	117 (39.1)
Regular alcohol drinkers, n (%)	267 (89.3)
Oral contraceptive use, n (%)	103 (34.4)
Ethnicity, n (%)	
White	283 (94.6)
Afro-Caribbean	3 (1.0)
Asian	8 (2.7)
Other	5 (1.7)
Vegetarians, n (%)	26 (8.7)
B ₁₂ supplement users, n (%)	31 (10.4)
B ₁₂ intake, diet only ($\mu\text{g/day}$)	3.82 (2.75, 5.02)
B ₁₂ intake, diet+supplements ($\mu\text{g/day}$)	3.83 (2.82, 5.20)
Folic acid supplement users, n (%)	32 (10.7)
Folate intake with supplements ($\mu\text{g/day}$)	237 (177, 315)

Continuous variables are mean $\pm$ SD or median (IQR).

Categorical variables are n (%).

BMI, body mass index; NDNS, National Diet and Nutrition Surveys.

Table 2 Comparison of B₁₂, folate and plasma homocysteine concentrations in women according to the UK RNI for vitamin B₁₂ intake

Female 19–39 years	All subjects	UK RNI (µg/day)		p value*
		<1.5	≥1.5	
Number (%)	299 (100)	11 (3.7)	288 (96.3)	
B ₁₂ intake, diet only (µg/day)	3.82 (2.75, 5.02)†	1.29 (0.64, 1.46)	3.86 (2.86, 5.09)	<0.001
B ₁₂ intake, diet+supplements (µg/day)	3.83 (2.82, 5.20)	1.29 (0.98, 1.46)	3.92 (2.88, 5.32)	<0.001
Serum B ₁₂ (pmol/L)	241 (188, 324)	169 (153, 256)	244 (189, 325)	0.05
B ₁₂ deficiency (<150 pmol/L), n (%)	36 (12.0)	2 (18.2)	34 (11.8)	NS
Serum folate (nmol/L)	19.5 (14.1, 26.7)	14.3 (13.6, 21.3)	19.7 (14.2, 27.0)	NS
Serum folate deficiency (<10 nmol/L), n (%)	18 (6.1)	0 (0)	18 (6.3)	NS
Red cell folate (nmol/L)	584 (473.9, 750.6)	460 (372, 739)	585 (478, 751)	NS
Red cell folate deficiency (<350 nmol/L), n (%)	13 (4.4)	2 (18.2)	11 (3.8)	NS
Hcy (µmol/L)	9.4 (9.1, 9.8)	11.9 (9.6, 14.3)	9.2 (7.8, 11.4)	<0.05
High Hcy (>12 µmol/L), n (%)	62 (21.2)	5 (50)	57 (20.1)	<0.05
High Hcy (>15 µmol/L), n (%)	24 (8.2)	2 (20)	22 (7.8)	NS

*Comparison between lower and higher B₁₂ intake groups. For categorical variables, Student's t-test was used (after log transformation); for continuous variables, Fisher's exact test was used.

†Median, 25–75th centile in parentheses (all such values).

Hcy, homocysteine; NS, not significant; UK RNI, UK Recommended Nutrient Intake.

abnormalities when their estimated was lower than the UK RNI (9.1% vs 3.5%, $p=0.001$). There is evidence that B₁₂ levels <258 pmol/L may be indicative of B₁₂ deficiency in certain individuals with concomitant elevation of Hcy and methylmalonic acid (MMA).³⁷ In total, 44.0% of women had B₁₂ levels in this borderline range of 150–258 pmol/L. In this subgroup, mean Hcy levels were significantly higher than the group with B₁₂>258 pmol/L (10.4 vs 9.2 µmol/L, $p=0.02$) despite similar folate levels (21.4 vs 22.0 nmol/L, $p=NS$).

The plasma Hcy concentrations were higher in the lower intake group (11.9 vs 9.2 µmol/L, $p<0.05$) although hyperhomocysteinemia (Hcy>12 µmol/L) was present in 20% of women with apparently adequate B₁₂ intake (table 2). Serum and red cell folate deficiency rates were 6.1% and 4.4%, respectively, in the whole population. There were no differences in the folate deficiency rates between above and below the UK RNI B₁₂ intake groups. In total, 34.4% of the women were taking the oral contraceptives and their B₁₂ values were lower than those who were not (median 211.5 vs 267.5 pmol/L, $p<0.001$).

In total, 8.7% of women in the whole cohort were vegetarian or vegan and their median dietary intake of B₁₂ was non-significantly lower than non-vegetarians (2.95 vs 3.87 µg/day), while folate consumption in the former group was higher (see online supplementary table S1). Vegetarians had lower serum B₁₂ concentrations (median 192 vs 248 pmol/L, $p<0.01$) but their folate or Hcy concentrations did not vary significantly.

Predictors of B₁₂, folate and Hcy

There was a positive correlation between serum B₁₂ values and daily B₁₂ intake (Pearson's $r=0.27$, $p<0.001$) (figure 1). Simple linear regression analyses of the predictors of B₁₂, folate and Hcy are shown in table 3. After

adjusting for the likely confounders, daily B₁₂ and folic acid intakes were positive predictors of serum B₁₂ ($\beta=0.28$, $p<0.001$) and serum folate ($\beta=0.33$, $p<0.001$), respectively. Along with age, serum B₁₂, serum folate and B₁₂ intake were independent predictors of Hcy, though it would appear that serum folate is strongest based on the value of the β coefficient (table 3).

Relationship between B₁₂ intake and associated biomarkers

In total, 5 out of 299 (1.7%) women consumed below the EAR of 1.25 µg/day, but none of them had B₁₂ levels <150 pmol/L. Conversely, in the 98.3% of women with 'adequate' EAR category for B₁₂ consumption, 12.2% of them had serum levels below 150 pmol/L.

In order to determine the trend of blood B₁₂ and Hcy concentrations with increasing intakes of B₁₂, we divided the cohort in quintiles of B₁₂ intake. The median levels

Figure 1 Correlation between daily B₁₂ intake and serum B₁₂ values. §Log transformed for statistical comparisons.

Table 3 Multiple linear regression analysis of predictors of serum B₁₂, folate and homocysteine

Variables	Serum B ₁₂ *		Serum folate		Homocysteine*	
	β coefficient	p Value	β coefficient	p Value	β coefficient	p Value
Age	–	NS	–	NS	0.18	0.001
BMI	–	NS	–	NS	–	NS
Smoking	–0.12	<0.05	–	NS	–	NS
Alcohol	–	NS	–	NS	–	NS
Oral contraceptive use	–0.29	<0.001	0.11	<0.05	–	NS
TC:HDL ratio*	–	NS	–	NS	–	NS
Vegetarian or vegan diet	–0.18	<0.01	–	NS	–	NS
B ₁₂ supplement use	–	NS	Not included	NA	–	NS
Daily B ₁₂ intake*	0.28	<0.001	Not included	NA	–0.16	0.001
Folic acid supplement use	Not included	NA	0.18	<0.01	–	NS
Daily folate intake*	Not included	NA	0.33	<0.001	–	NS
Serum B ₁₂ *	–	NS	–	NS	–0.20	<0.001
Serum folate*	–	NS	–	NS	–0.35	<0.001

*Log transformed for statistical comparison.

–, Tested but not significant in the model;

BMI, body mass index; NA, not applicable; NS, not significant; TC:HDL, total cholesterol:high-density lipoprotein ratio.

in each quintile and biomarker values are represented in figures 2A, B. Women in quintiles 4 and 5 (median intake 4.9 and 7.1 µg/day, respectively) had significantly higher mean B₁₂ and lower mean Hcy concentrations than quintile 1 (291 and 322 vs 229 pmol/L; 9.0 and 9.0 vs 11.4 µmol/L, respectively). Additionally, the Hcy levels appear to level off between quintiles 4 and 5, suggesting that increasing intakes above 5 µg/day is unlikely to provide higher B₁₂ at the tissue level (figure 2B). To confirm this, we performed logistic regression to determine the intake threshold at which the odds of hyperhomocysteinemia would reduce significantly after correcting for confounders. Using 0.25 µg increments, a threshold of 4.75 µg/day significantly reduced the odds of Hcy>12 µmol/L (adjusted OR (AOR) 0.35, 95% CI 0.14 to 0.88). When serum B₁₂ was added to the model, the significance was lost, suggesting that the influence of dietary B₁₂ intake on circulating Hcy is mediated through serum B₁₂ values in a folate-replete population (data not shown).

Micronutrient intake: comparisons between NDNS 2000/2001 and 2008/2012 data

Comparison of the mean nutrient intake among young women between the two surveys is shown in table 4. The proportion of women consuming below the UK RNI for folate increased in the more recent survey (44% vs 33%, p<0.01) and median intake levels fell by 14% (206 vs 235 pmol/L, p<0.001). There were no significant differences in the B₁₂ intake values.

DISCUSSION

Our study shows contemporary data of B₁₂ and folate intakes and serum levels in population-based nutritional surveys involving women of childbearing age, who are representative of the UK population. The key findings are that, despite an apparent adequate daily intake of

B₁₂, a high proportion of women have B₁₂ deficiency and hyperhomocysteinemia.

Optimum B₁₂ intake and biomarker levels

Our data showed positive correlation between B₁₂ intake and blood levels with a trend of increasing B₁₂ concentration even in intakes up to 7 µg/day, although the corresponding reductions in Hcy levels level off around the intake of 5–7 µg/day. Additionally, the consumption of around 4.75 µg/day was independently associated with a decrease in the odds of hyperhomocysteinemia. Our findings are supportive of the dose–response relationship between intake and blood levels of B₁₂ found previously, which showed doubling B₁₂ intakes would increase B₁₂ concentrations by around 10%.³⁸

A daily intake of 4–10 µg has been suggested by other studies to stabilise levels of B₁₂, Hcy and the associated MMA in adults,^{39 40} but more evidence is needed before extrapolating these figures to women of childbearing age due to the different requirements and implications should they become pregnant. In a randomised controlled trial where all subjects consumed 8.6 µg/day of B₁₂, pregnant women had higher holotranscobalamin (holoTC):total B₁₂ ratios than non-pregnant, non-lactating controls, suggesting that such high intakes were required to provide sufficient supply to the developing fetus.⁴¹ More research is required to decide on the upper limit of B₁₂ intake in women before and during pregnancy, as higher B₁₂ intake in the third trimester was positively associated with offspring birth weight in high-BMI women (who are already at high risk of having macrosomic babies), although circulating B₁₂ levels were not reported here.⁴² There is also no consistent evidence that increased B₁₂ intake or supplementation is associated with reduction in the prevalence of subclinical B₁₂ deficiency in adults (particularly neurocognitive decline in the elderly).^{43 44} Therefore, until the

availability of further studies, our data call for revision of the UK RNI (and EAR) for B₁₂ to be at least in line with the current European recommendations (4.5 µg/day).²⁴

Figure 2 Relationship between B₁₂ intake in quintiles and (A) mean serum B₁₂ and (B) mean plasma homocysteine concentrations. Mean±SEM values are plotted against the median B₁₂ intake in each quintile. One-way analysis of variance (ANOVA) test used to compare the means between the quintiles (after log transformation) and Tukey's post hoc analysis done. Mean biomarker levels differed as compared with quintile 1 as follows: *p<0.05; **p<0.01; ***p<0.001.

The estimated B₁₂ intake in preconceptional or women in early pregnancy and specifically its relationship with serum levels has not been widely studied. In developing countries, over 50% of women of reproductive age do not meet the RNIs for B₁₂^{45 46} and separate surveys from three developing countries (Turkey, Iran and China) found B₁₂ deficiency rates of 21–23%.^{47–49} An Australian study on young females found a similar rate to ours (11.4%), albeit using a lower threshold of 120 pmol/L.⁵⁰ As the B₁₂ levels fall by around 10% from preconception to early pregnancy,¹⁹ if we extrapolate our findings, the proportion of pregnant women with B₁₂ levels below 150 pmol/L is likely to be much higher. More than a third of vegetarian/vegan women in our study had B₁₂ deficiency, which was in keeping with studies from other UK adult population⁵¹ and elsewhere.⁵² Thus our findings highlight the need for specific advice for the vegetarian/vegan population about potential sources of B₁₂, as well as recommending them to have their B₁₂ levels checked (and corrected) if they are planning pregnancy.

Folate intake

Our study shows that consumption of dietary folate has fallen by around 14% in the 10 years between the two surveys and nearly 50% of women are now consuming below the UK RNI. The Scientific Advisory Committee on Nutrition within the UK Food Standards Agency has recommended mandatory folic acid fortification of food products to the UK DOH,⁵³ which, if implemented, would increase folate intake and consequently serum levels in the population as it did in the USA.⁵⁴ However, if mandatory fortification does occur, it is possible that improving folate levels can reduce B₁₂ levels due to usage of the available B₁₂.⁵⁵ In addition, B₁₂ deficiency is the strongest driver of Hcy in a folate-replete population.⁵⁶ Therefore fortification of food products with B₁₂ together with folate should be considered if there is to be a policy change in the UK or, as a minimum, B₁₂ supplements recommended for women in the periconceptional period, especially if they are at high risk of deficiency.

The strengths of our study are that this is the first study of its kind from the UK to evaluate B₁₂ and folate

Table 4 Comparison between folic acid and B₁₂ intakes between the NDNS 2000/2001 and 2008/2011 cohorts

Female 19–39 years	NDNS 2000/2001 cohort	NDNS 2008/2011 cohort
Number	299	395
Folic acid supplement users, n (%)	32 (10.7)	12 (3.0)***,†
Folate intake, diet only (µg/day)	234.9 (175.7, 302.7)‡	205.8 (162.3, 261.2)***
n (%) consuming below UK RNI of folate	99 (33.1)	173 (43.8)**
B ₁₂ supplement users, n (%)	31 (10.4)	33 (8.4)
B ₁₂ intake, diet only (µg/day)	3.82 (2.75, 5.01)	3.78 (2.64, 4.84)
n (%) consuming below UK RNI of B ₁₂	11 (3.7)	24 (6.1)

Significance: **p<0.01; ***p<0.001

†p Values from independent-samples t-test for continuous variables (after log transformation) or χ^2 test for categorical variables.

‡Median, 25–75th centile in parentheses (all such values).

UK RNI, UK Recommended Nutrient Intake; NDNS, National Diet and Nutrition Survey.

among women in the preconceptional stage. Extensive data on these women including anthropometry, biochemical markers and dietary information allowed comparison of intake and serum levels adjusted for possible confounders. There were three important limitations of this data set: (1) lack of sufficient biochemical data in the second survey, (2) lack of data on other biomarkers such as holoTC or MMA and (3) lack of haematological and clinical information relevant to the effects of B₁₂. Since we were able to compare the nutrient intakes between the two cohorts, which was predictive of serum levels in the earlier survey, we believe that the availability of serum B₁₂ in the latter cohort may not necessarily change the findings. Hcy is a readily available marker in clinical practice as opposed to holoTC and MMA. Therefore, our findings are applicable to wider clinical practice. Any future prospective studies involving preconceptional or pregnant women should include more clinical information as well as these biomarkers for a better evaluation of B₁₂ status. In conclusion, our study supports revision of the UK RNI to at least match the European recommendations and also calls for assessing maternal B₁₂ status in preconception or early pregnancy.

Acknowledgements The authors acknowledge the British National Diet and Nutrition Survey for kindly providing the public database for detailed quantitative assessment of the samples, nutritional, dietary and laboratory characteristics.

Contributors PS conceived the research question and study design. NS and AA contributed to data collection, statistical analysis and data interpretation and NS drafted the initial manuscript. HM and HV contributed to data collection. All authors contributed, revised and edited the manuscript. PS is the guarantor of this work and had full access to all the data presented in the study and takes full responsibility for the integrity and the accuracy of the data analysis.

Funding This research received no specific grant from any funding agency in the public, commercial or not-for-profit sectors. PS is partly supported by a project grant from MRC which is designed to assess the impact of B₁₂ levels in early pregnancy on gestational diabetes.

Competing interests None declared.

Ethics approval Ethical approval was obtained from a multicentre Research Ethics Committee and the Oxfordshire A Research Ethics Committee for the first and second surveys, respectively, and all local research ethics committees covering areas where the fieldwork was conducted.

Provenance and peer review Not commissioned; externally peer reviewed.

Data sharing statement No additional data are available.

Open Access This is an Open Access article distributed in accordance with the terms of the Creative Commons Attribution (CC BY 4.0) license, which permits others to distribute, remix, adapt and build upon this work, for commercial use, provided the original work is properly cited. See: <http://creativecommons.org/licenses/by/4.0/>

REFERENCES

- Yajnik CS, Deshmukh US. Fetal programming: maternal nutrition and role of one-carbon metabolism. *Rev Endocr Metab Disord* 2012;13:121–7.
- Saravanan P, Yajnik CS. Role of maternal vitamin B12 on the metabolic health of the offspring: a contributor to the diabetes epidemic? *Br J Diabetes Vasc Dis* 2010;10:109–14.
- Wald DS, Law M, Morris JK. Homocysteine and cardiovascular disease: evidence on causality from a meta-analysis. *BMJ* 2002;325:1202.
- Zhou J, Austin RC. Contributions of hyperhomocysteinemia to atherosclerosis: Causal relationship and potential mechanisms. *Biofactors* 2009;35:120–9.
- Ray JG, Blom HJ. Vitamin B12 insufficiency and the risk of fetal neural tube defects. *QJM* 2003;96:289–95.
- George L, Mills JL, Johansson AL, et al. Plasma folate levels and risk of spontaneous abortion. *JAMA* 2002;288:1867–73.
- Ronnenberg AG, Goldman MB, Chen D, et al. Preconception homocysteine and B vitamin status and birth outcomes in Chinese women. *Am J Clin Nutr* 2002;76:1385–91.
- Sukumar N, Rafnsson SB, Kandala NB, et al. Prevalence of vitamin B-12 insufficiency during pregnancy and its effect on offspring birth weight: a systematic review and meta-analysis. *Am J Clin Nutr* 2016;103:1232–51.
- Centers for Disease Control and Prevention (CDC). Spina bifida and anencephaly before and after folic acid mandate—United States, 1995–1996 and 1999–2000. *MMWR Morb Mortal Wkly Rep* 2004;53:362–5.
- Ray JG, Wyatt PR, Thompson MD, et al. Vitamin B12 and the risk of neural tube defects in a folic-acid-fortified population. *Epidemiology* 2007;18:362–6.
- Nelen WL, Blom HJ, Steegers EA, et al. Hyperhomocysteinemia and recurrent early pregnancy loss: a meta-analysis. *Fertil Steril* 2000;74:1196–9.
- Vollset SE, Refsum H, Irgens LM, et al. Plasma total homocysteine, pregnancy complications, and adverse pregnancy outcomes: the Hordaland Homocysteine study. *Am J Clin Nutr* 2000;71:962–8.
- Hogeveen M, Blom HJ, den Heijer M. Maternal homocysteine and small-for-gestational-age offspring: systematic review and meta-analysis. *Am J Clin Nutr* 2012;95:130–6.
- Koukoura O, Sifakis S, Spandidos DA. DNA methylation in the human placenta and fetal growth (review). *Mol Med Rep* 2012;5:883–9.
- Finer LB, Zolna MR. Unintended pregnancy in the United States: incidence and disparities, 2006. *Contraception* 2011;84:478–85.
- Balcı YI, Ergin A, Karabulut A, et al. Serum vitamin B12 and folate concentrations and the effect of the Mediterranean diet on vulnerable populations. *Pediatr Hematol Oncol* 2014;31:62–7.
- Dwarkanath P, Barzilay JR, Thomas T, et al. High folate and low vitamin B-12 intakes during pregnancy are associated with small-for-gestational age infants in South Indian women: a prospective observational cohort study. *Am J Clin Nutr* 2013;98:1450–8.
- García-Casal MN, Osorio C, Landaeta M, et al. High prevalence of folic acid and vitamin B12 deficiencies in infants, children, adolescents and pregnant women in Venezuela. *Eur J Clin Nutr* 2005;59:1064–70.
- Murphy MM, Molloy AM, Ueland PM, et al. Longitudinal study of the effect of pregnancy on maternal and fetal cobalamin status in healthy women and their offspring. *J Nutr* 2007;137:1863–7.
- [No authors listed]. Dietary reference values for food energy and nutrients for the United Kingdom. Report of the Panel on Dietary Reference Values of the Committee on Medical Aspects of Food Policy. *Rep Health Soc Subj (Lond)* 1991;41:1–210.
- Whitehead RG. Dietary reference values. *Proc Nutr Soc* 1992;51:29–34.
- Baker SJ, Mathan VI. Evidence regarding the minimal daily requirement of dietary vitamin B12. *Am J Clin Nutr* 1981;34:2423–33.
- Cooper BA, Lowenstein L. Vitamin B-12-folate interrelationships in megaloblastic anaemia. *Br J Haematol* 1966;12:283–96.
- EFSA NDA Panel (EFSA Panel on Dietetic Products NAA. Scientific Opinion on Dietary Reference Values for cobalamin (vitamin B12). *EFSA J* 2015;13:4150.
- Doets EL, Cavelaars AE, Dhonukshe-Rutten RA, et al. Explaining the variability in recommended intakes of folate, vitamin B12, iron and zinc for adults and elderly people. *Public Health Nutr* 2012;15:906–15.
- Institute of Medicine. *Dietary reference intakes for thiamin, riboflavin, niacin, vitamin B6, folate, vitamin B12, pantothenic acid, biotin, and choline. Dietary Reference Intakes for Thiamin, Riboflavin, Niacin, Vitamin B6, Folate, Vitamin B12, Pantothenic Acid, Biotin, and Choline*. Washington DC: National Academies Press, 1998:306–56.
- Henderson L, Gregory J, Swan G. *The national diet & nutrition survey: adults aged 19 to 64 years. Volume 1. Types and quantities of foods consumed*. London: The Stationery Office, 2002.
- Henderson L, Irving K, Gregory J, et al. *The national diet & nutrition survey: adults aged 19 to 64 years. Volume 3. Vitamin and mineral intake and urinary analytes*. London: The Stationery Office, 2003.

29. Bates B, Lennox A, Prentice A, *et al.* National Diet and Nutrition Survey: Results from Years 1, 2, 3 and 4 (combined) of the Rolling Programme (2008/2009–2011/2012). 2014. https://www.gov.uk/government/uploads/system/uploads/attachment_data/file/310995/NDNS_Y1_to_4_UK_report.pdf
30. Skinner C, Holmes D. The 2000-01 National Diet and Nutrition Survey of Adults aged 19-64 years: The impact of non-response. In National Diet and Nutrition survey Adults Aged 19–64 years. Appendix E. 2002. <http://www.foodgovuk/science/101717/ndnsdocuments/ndnsappendices>
31. Møller J, Christensen L, Rasmussen K. An external quality assessment study on the analysis of methylmalonic acid and total homocysteine in plasma. *Scand J Clin Lab Invest* 1997;57:613–19.
32. Beaton GH. Approaches to analysis of dietary data: relationship between planned analyses and choice of methodology. *Am J Clin Nutr* 1994;59(1 Suppl):253S–61S.
33. Carmel R. Biomarkers of cobalamin (vitamin B-12) status in the epidemiologic setting: a critical overview of context, applications, and performance characteristics of cobalamin, methylmalonic acid, and holotranscobalamin II. *Am J Clin Nutr* 2011;94:348S–58S.
34. [No authors listed]. Folate and vitamin B12 deficiencies. Proceedings of a WHO technical consultation held 18–21 October, 2005, in Geneva, Switzerland. *Food Nutr Bull* 2008;29(2 Suppl):S1–246.
35. Refsum H, Smith AD, Ueland PM, *et al.* Facts and recommendations about total homocysteine determinations: an expert opinion. *Clin Chem* 2004;50:3–32.
36. Fokkema MR, Weijer JM, Dijck-Brouwer DA, *et al.* Influence of vitamin-optimized plasma homocysteine cutoff values on the prevalence of hyperhomocysteinemia in healthy adults. *Clin Chem* 2001;47:1001–7.
37. Lindenbaum J, Rosenberg IH, Wilson PW, *et al.* Prevalence of cobalamin deficiency in the Framingham elderly population. *Am J Clin Nutr* 1994;60:2–11.
38. Dullemeijer C, Souverein OW, Doets EL, *et al.* Systematic review with dose-response meta-analyses between vitamin B-12 intake and European Micronutrient Recommendations Aligned's prioritized biomarkers of vitamin B-12 including randomized controlled trials and observational studies in adults and elderly persons. *Am J Clin Nutr* 2013;97:390–402.
39. Bor MV, von Castel-Roberts KM, Kauwell GP, *et al.* Daily intake of 4 to 7 microg dietary vitamin B-12 is associated with steady concentrations of vitamin B-12-related biomarkers in a healthy young population. *Am J Clin Nutr* 2010;91:571–7.
40. Vogiatzoglou A, Smith AD, Nurk E, *et al.* Dietary sources of vitamin B-12 and their association with plasma vitamin B-12 concentrations in the general population: the Hordaland Homocysteine Study. *Am J Clin Nutr* 2009;89:1078–87.
41. Bae S, West AA, Yan J, *et al.* Vitamin B-12 Status Differs among Pregnant, Lactating, and Control Women with Equivalent Nutrient Intakes. *J Nutr* 2015;145:1507–14.
42. Horan MK, McGowan CA, Gibney ER, *et al.* The association between maternal dietary micronutrient intake and neonatal anthropometry—secondary analysis from the ROLo study. *Nutr J* 2015;14:105.
43. Smith AD, Refsum H. Do we need to reconsider the desirable blood level of vitamin B12? *J Intern Med* 2012;271:179–82.
44. Smith AD, Smith SM, de Jager CA, *et al.* Homocysteine-lowering by B vitamins slows the rate of accelerated brain atrophy in mild cognitive impairment: a randomized controlled trial. *PLoS ONE* 2010;5:e12244.
45. Arsenault JE, Yakes EA, Islam MM, *et al.* Very low adequacy of micronutrient intakes by young children and women in rural Bangladesh is primarily explained by low food intake and limited diversity. *J Nutr* 2013;143:197–203.
46. Nguyen PH, Nguyen H, Gonzalez-Casanova I, *et al.* Micronutrient intakes among women of reproductive age in Vietnam. *PLoS One* 2014;9:e89504.
47. Abdollahi Z, Elmadfa I, Djazayeri A, *et al.* Folate, vitamin B12 and homocysteine status in women of childbearing age: baseline data of folic acid wheat flour fortification in Iran. *Ann Nutr Metab* 2008;53:143–50.
48. Karabulut A, Guler ÖT, Karahan HT, *et al.* Premarital screening of 466 Mediterranean women for serum ferritin, vitamin B12, and folate concentrations. *Turk J Med Sci* 2015;45:358–63.
49. Zhu JH, Hu DJ, Hao L, *et al.* Iron, folate, and B(12) deficiencies and their associations with anemia among women of childbearing age in a rural area in Northern China. *Int J Vitam Nutr Res* 2010;80:144–54.
50. Fayet-Moore F, Petocz P, Samman S. Micronutrient status in female university students: iron, zinc, copper, selenium, vitamin B12 and folate. *Nutrients* 2014;6:5103–16.
51. Gilsing AM, Crowe FL, Lloyd-Wright Z, *et al.* Serum concentrations of vitamin B12 and folate in British Male omnivores, vegetarians and vegans: results from a cross-sectional analysis of the EPIC-Oxford cohort study. *Eur J Clin Nutr* 2010;64:933–9.
52. Pawlak R, Parrott SJ, Raj S, *et al.* How prevalent is vitamin B (12) deficiency among vegetarians? *Nutr Rev* 2013;71:110–17.
53. Scientific Advisory Committee on Nutrition. *Folate and disease prevention*. Food Standards Agency and the Department of Health, 2006.
54. Pfeiffer CM, Johnson CL, Jain RB, *et al.* Trends in blood folate and vitamin B-12 concentrations in the United States, 1988–2004. *Am J Clin Nutr* 2007;86:718–27.
55. Selhub J, Morris MS, Jacques PF, *et al.* Folate–vitamin B-12 interaction in relation to cognitive impairment, anemia, and biochemical indicators of vitamin B-12 deficiency. *Am J Clin Nutr* 2009;89:702S–6S.
56. Selhub J, Morris MS, Jacques PF. In vitamin B12 deficiency, higher serum folate is associated with increased total homocysteine and methylmalonic acid concentrations. *Proc Natl Acad Sci U S A* 2007;104:19995–20000.