BMJ Open Epidemiological survey of the feasibility of broadband ultrasound attenuation measured using calcaneal quantitative ultrasound to predict the incidence of falls in the middle aged and elderly

Ling-Chun Ou,^{1,2,3} Yin-Fan Chang,⁴ Chin-Sung Chang,⁴ Ching-Ju Chiu,⁵ Ting-Hsing Chao,⁶ Zih-Jie Sun,⁴ Ruey-Mo Lin,⁷ Chih-Hsing Wu^{4,5,8}

To cite: Ou L-C, Chang Y-F, Chang C-S, *et al.* Epidemiological survey of the feasibility of broadband ultrasound attenuation measured using calcaneal quantitative ultrasound to predict the incidence of falls in the middle aged and elderly. *BMJ Open* 2017;7:e013420. doi:10.1136/bmjopen-2016-013420

► Prepublication history for this paper is available online. To view these files please visit the journal online (<http://dx.doi.org/10.1136/bmjopen-2016-013420>).

Received 13 July 2016
Revised 8 December 2016
Accepted 9 December 2016

CrossMark

For numbered affiliations see end of article.

Correspondence to

Dr Chih-Hsing Wu;
paulo@mail.ncku.edu.tw

ABSTRACT

Objectives: We investigated whether calcaneal quantitative ultrasound (QUS-C) is a feasible tool for predicting the incidence of falls.

Design: Prospective epidemiological cohort study.

Setting: Community-dwelling people sampled in central western Taiwan.

Participants: A cohort of community-dwelling people who were ≥ 40 years old (men: 524; women: 676) in 2009–2010. Follow-up questionnaires were completed by 186 men and 257 women in 2012.

Methods: Structured questionnaires and broadband ultrasound attenuation (BUA) data were obtained in 2009–2010 using QUS-C, and follow-up surveys were done in a telephone interview in 2012. Using a binary logistic regression model, the risk factors associated with a new fall during follow-up were analysed with all significant variables from the bivariate comparisons and theoretically important variables.

Primary outcome measures: The incidence of falls was determined when the first new fall occurred during the follow-up period. The mean follow-up time was 2.83 years.

Results: The total incidence of falls was 28.0 per 1000 person-years for the ≥ 40 year old group (all participants), 23.3 per 1000 person-years for the 40–70 year old group, and 45.6 per 1000 person-years for the ≥ 70 year old group. Using multiple logistic regression models, the independent factors were current smoking, living alone, psychiatric drug usage and lower BUA (OR 0.93; 95% CI 0.88 to 0.99, $p < 0.05$) in the ≥ 70 year old group.

Conclusions: The incidence of falls was highest in the ≥ 70 year old group. Using QUS-C-derived BUA is feasible for predicting the incidence of falls in community-dwelling elderly people aged ≥ 70 years.

Strengths and limitations of this study

- The different risk factors for falling incidence were analysed in the 40–70 year old and the ≥ 70 year old groups.
- The use of calcaneal quantitative ultrasound (QUS-C)-derived broadband ultrasound attenuation for predicting the incidence of falls in community dwelling elderly people aged ≥ 70 years is examined.
- Whether QUS-C measured using other brands or at different sites in different populations would yield the same findings is unknown and warrants additional study.

INTRODUCTION

It is estimated that around one-fifth¹ to one-third² of elderly people fall at least once every year. Fall-related injuries, including soft-tissue injury, bone fractures, intracranial haemorrhage, functional decline, disability and death,² are hazardous and life threatening. Therefore, developing a method to determine the clinical risk factors related to falls is important.

Many risk factors for falls have been reported: older age,³ sex,^{1 3} lower body weight,¹ living alone,³ arthritis,⁴ previous falls,² using antipsychotics, antidepressants and other psychiatric medications,² stroke,⁵ hyperglycaemia,¹ sarcopenia,⁵ nutritional deficiency (hypoalbuminemia or anaemia),^{2 6} and impaired balance or gait.² Habitual smoking and alcohol consumption might result in frailty⁷ that leads to falls.⁸ Over the

past decades, quantitative ultrasound (QUS) has emerged as a reliable method of assessing skeletal status in osteoporosis and the risk of fractures in elderly women.^{9–12} The interrelationships between QUS and quadriceps muscle strength and gait speed are believed to be associated with falls.^{4 10} However, existing evidence is mostly hospital based^{6 12} or from cross-sectional studies.^{1 2 5 9–11} Furthermore, there are few community surveys on the incidence of falls.^{8 13–16} Most studies on the incidence of falls provide data from intervention programmes,¹⁶ focused only on the elderly and not on middle-aged people. Although several risk factors related to the incidence of falls have been reported,^{8 13–16} whether the risk factors related to falling might be different between Asian and Western populations^{13 14} or between middle aged people and the elderly are inconclusive.

QUS is a convenient tool for screening community-dwelling elderly people for osteoporosis¹⁷ and related osteoporotic fractures.^{18–22} Osteoporosis is defined as low bone mass or fragile bone quality;²³ thus, QUS, and especially calcaneal QUS (QUS-C),²⁴ might help clinical decision-making when dual-energy X-ray absorptiometry and vertebral fracture assessment disagree about a diagnosis of osteoporosis.²⁵ Ou *et al*¹¹ recently reported that QUS-C is a convenient screening tool in association with the prevalence of falls in community-dwelling middle-aged and elderly people. Based on the epidemiological cohort,¹¹ in the present study, we investigated the incidence of falls and the feasibility of using of QUS-C for predicting falls in middle-aged and elderly people. We hypothesised that QUS-C-derived broadband ultrasound attenuation (BUA) would be a convenient parameter for predicting the incidence of falls.

MATERIALS AND METHODS

Study design and population

Two thousand, two hundred and eight subjects were selected using a step by step, stratified systemic cluster sample of households from 2009 to 2010, with a final cohort population of 1200 (including 686 who agreed to receive a blood test) from Kukeng and Dou-Liou townships in Yun-Lin County in central western Taiwan.¹¹ The raw response rate was 54.3%. Of the 686 enrolees who underwent a blood test in 2009–2010 and provided written informed consent, 443 (men: 186; women: 257) answered the follow-up questionnaires in a telephone interview in 2012 (figure 1). There were no significant differences in the basic characteristics between the 514 enrolees who did not take a blood test and the 686 who did; or, except for a minor difference in age, between the 443 enrolees who were followed up and the 243 who were not. The 243 enrolees who were not followed up were slightly but significantly younger (57.71 ± 11.46 years old) than the original 1200 participants (59.30 ± 11.40 years old) ($p < 0.01$). The follow-up time, defined by the date of a new fall or the date of the telephone interview if there was no new fall, ranged from 0.91 to 3.48 years (mean: 2.83 years). This study was approved by the Institutional Review Board of National Cheng Kung University Hospital (A-ER-100-347). Each participant signed a written informed consent form.

Measurements

From 2009 to 2010, participants were asked to complete a structured questionnaire¹¹ that asked about lifestyle habits (exercise, smoking and alcohol drinking), living and socioeconomic status, past medical conditions, drug history and fracture history. Dichotomous variables were, for example, exercise (participants were defined as

Figure 1 Flow chart of baseline and follow-up study populations.

moderate exercisers if they exercised more than three times per week), smoking (participants were defined as a smoker if they smoked ≥ 20 cigarettes per month for ≥ 6 months), drinking (participants were defined as a drinker if they consumed alcohol at least once per week for > 6 months), and living alone (participants were defined as living alone if they lived with no other person).¹¹ All participants underwent anthropometric and body composition measurements: skeletal muscle mass (SMM) measured using an 8-polar multi-frequency bioelectrical impedance analysis (Inbody720; Biospace, Seoul, Korea).²⁶ The SMM (kg) was estimated using Janssen's equation²⁷ and was validated in Taiwan:²⁶

$$\text{SMM} = \left[\frac{\text{Ht}^2}{\text{R} \times 0.401} + (\text{sex} \times 3.825) + (\text{age} \times -0.071) \right] + 5.102$$

where height is in cm; resistance is in ohms; for sex, men=1 and women=0; and age is in years. QUS-C-derived BUA (CUBA Clinical scanner MK2.6; McCue Ultrasonics, Winchester, UK) and coefficients of variation (1.3%) were obtained at the same time by a single QUS technician during 2009–2010.

Baseline haemoglobin and serum albumin were measured in 2009–2010 and used as a reference index of nutritional status.^{28 29} Participants were considered anaemic if haemoglobin (Hb) was < 13 g/dL in men and < 12 g/dL in women.²⁸ Participants were considered hypoalbuminaemic if their serum albumin was < 3.5 mg/dL.²⁹ Participants who reported a fall during the baseline survey were defined as having a history of falls if they also met any of the two following conditions: (1) they unpredictably tilted downward when standing up, sitting, or walking; or (2) they unintentionally contacted the floor with their body when changing position.¹¹ The same researcher who conducted the baseline interviews used, in the 2012 telephone interviews, the follow-up structured questionnaire that asked about new falls. Participants were defined as new fallers if they fell during the follow-up period, and as non-fallers if they had no recorded falls. We calculated the incidence of falls in person-years, based on Merom *et al.*,³⁰ using the following formula:

$$\left[\frac{\text{number of new fallers}}{(\text{average duration of telephone follow-up} \times \text{number of non-fallers} + \text{average duration before new fall} \times \text{number of new fallers})} \right] \times 1000(\%)$$

The time interval (years) before the new fall was defined as from the month during which participants underwent a free community osteoporosis evaluation until the new fall occurred. The time interval (year) of the telephone follow-up was defined as from the middle of the month during which participants underwent the free community osteoporosis evaluation until telephone contact.

Statistical analysis

Data were collected and analysed using SPSS V.16 for Windows (SPSSWIN, V.16.0, Chicago, Illinois, USA). The continuous variables were expressed using mean $\pm$ SD, and the categorical variables were expressed as percentages. To differentiate potentially different risk factors, participants were assigned to one of two age groups: middle aged (40–70 years old) or elderly (≥ 70 years old) because the elderly are more susceptible to osteoporosis and advised to have a vertebral fracture assessment to reduce the diagnostic discrepancy of osteoporosis.²⁴ Comparisons of categorical variables between groups were analysed using χ^2 and Fisher's exact tests, and continuous variables were analysed using an independent t test. The associated risk factors for a new faller were analysed using a binary logistic regression model with all significant variables from the bivariate comparisons and with theoretically important variables. Significance was set at $p < 0.05$ (two tailed).

RESULTS

Of the 443 participants, the incidence of falls was 28.0 per 1000 person-years in the ≥ 40 year old group, 23.3 in the 40–70 year old group, and 45.6 in the ≥ 70 year old group. For participants with a history of falls, the incidence was 56.9 per 1000 person-years, and for those without a history of falls, it was 20.9. The univariate analysis showed a significantly higher fall history for new fallers in the 40–70 year old group (table 1). There is also a significant difference between new fallers and non-fallers ≥ 70 years old and living alone, and 40–70 years old with anaemia.

A binary logistic regression analysis to determine the independent factors for the incidence of falls—not used if the number of cases was less than one: cerebrovascular accident, osteoarthritis and arrhythmia—shows that the independent predictors of falls in the overall population were being male, current smoking, current alcohol consumption, living alone, a history of falls and anaemia (table 2). In the 40–70 year old group, the independent factors were being male, current alcohol consumption, fall history and fracture history. In the ≥ 70 year old group, the independent factors were current smoking, living alone, psychiatric drug user and QUS-C-derived BUA (OR 0.93; 95% CI 0.88 to 0.99).

DISCUSSION

We found that the incidence of falls was 28.0 per 1000 person-years for the ≥ 40 year old group (all participants), 23.3 for the 40–70 year old group, and 45.6 for the ≥ 70 year old group. The incidence of falls in our study was lower than that found in Western studies,³¹ perhaps because Asians live in crowded conditions¹⁴ and fewer elderly people live alone in Asia (5.6% of our participants) than in the West (Caucasian: 53%).³ Furthermore, incidence-of-fall studies typically focus on interventional effects on the elderly with a history of

Table 1 Characteristics between fallers and non-fallers in 443 men and women during follow-up

Variables	≥70 years old (n=94)		40–70 years old (n=349)	
	New faller (%)	Non-faller (%)	New faller (%)	Non-faller (%)
Number of cases	12 (12.8)	82 (87.2)	23 (6.6)	326 (93.4)
Age (years)	77.1 (4.9)	75.0 (4.4)	53.5 (7.4)	53.0 (7.5)
Male	6 (50.0)	45 (54.9)	5 (21.7)	130 (39.9)
Body height (cm)	154.85 (7.73)	156.66 (8.71)	157.60 (6.85)	159.56 (7.70)
Skeletal muscle mass (kg)	21.93 (3.60)	22.28 (4.70)	22.7 (4.9)	23.3 (4.9)
Living alone	5 (41.7)**	7 (8.5)	1 (4.3)	12 (3.7)
Current smoking	2 (16.7)	5 (6.1)	4 (17.4)	40 (12.3)
Current alcohol consumption	1 (8.3)	7 (8.5)	6 (26.1)	46 (14.1)
Moderate exercise habit	1 (8.3)	20 (24.4)	7 (30.4)	116 (35.6)
Fall history	6 (50.0)	24 (29.3)	8 (34.8)*	51 (15.6)
Fracture history	3 (25.0)	15 (18.3)	9 (39.1)	67 (20.6)
Diabetes mellitus history	1 (8.3)	11 (13.4)	1 (4.3)	22 (6.7)
Hypertension history	7 (58.3)	33 (40.2)	2 (8.7)	62 (19.0)
Cerebrovascular accident history	0 (0)	1 (1.2)	0 (0)	3 (0.9)
Arrhythmia history	0 (0)	2 (2.4)	2 (8.7)	15 (4.6)
Osteoarthritis history	0 (0)	4 (4.9)	0 (0)	10 (3.1)
Psychiatric drug user	2 (16.7)	3 (3.7)	1 (4.3)	17 (5.2)
Anaemia†	6 (50.0)	23 (28.0)	6 (26.1)*	29 (8.9)
Albumin (mg/dL)	4.38 (0.19)	4.46 (0.23)	4.57 (0.25)	4.57 (0.22)
QUS-C-derived BUA (dB/MHz)	52.17 (18.35)	63.84 (20.29)	73.35 (14.09)	70.94 (16.68)

Continuous variables, mean (SD), were analysed using an independent t test; categorical variables, n (%), were analysed using a χ^2 test.

* $p < 0.05$, ** $p < 0.01$.

†Anaemia in men if haemoglobin (Hb) < 13 g/dL, and in women if Hb < 12 g/dL.

BUA, broadband ultrasound attenuation; QUS-C, calcaneal quantitative ultrasound.

falls.^{14 15} In contrast, our epidemiological study was community based and included middle-aged (40–70 years old) participants in addition to participants with and without a history of falls, which might have yielded a lower incidence of falls.

Recall bias might also be a concern when trying to determine the incidence of falls during the study period. Different researchers have used different methods to periodically follow up participants and more frequently obtain a record of falls.¹⁴ In contrast, in the present study, the same trained professional asked participants about their falls in telephone interviews. A fall with injuries is clinically relevant and will easily be remembered for years. Because most falls do not result in injuries, only major falls will be reported by participants and is worth investigating, as in this study.

Consistent with other studies,^{12 13 15 16} men were negatively associated with falls in our ≥ 40 year old study population. Current smoking and current alcohol consumption are risk factors associated with frailty,⁷ which is highly associated with falls.¹⁶ Anaemia is common in the frail elderly,⁷ and is positively associated with the incidence of falls.³² Histories of using psychiatric medications¹⁴ and living alone^{3 14} are positively associated with the incidence of falls in the elderly. A history of falls is known to be an independent risk factor for future falls.³³ A history of fractures was a risk factor for future falls in a cross-sectional study.¹¹ All of these risk factors for falls are compatible with our findings. Why these risk factors were different in the ≥ 70 year old and the

40–70 year old groups might be because the number of cases for several variables was insufficient to allow us to draw any significant conclusions despite the consistent direction of the OR for most of the variables. However, a higher prevalence of insomnia³⁴ in elderly people increases their use of psychiatric medications and, in turn, their probability of falling.² The primary fall-inducing substance in the ≥ 70 year old group was psychiatric medication, and in the 40–70 year old group it was alcohol.³⁵ Therefore, in this study it is reasonable to use alcohol consumption as an independent risk factor for the incidence of falls in the 40–70 year old group and psychiatric drug use in the ≥ 70 year old group.

This is the first study to report that the QUS-C-derived BUA is a predictor of falls. After adjusting for the major factors of current smoking, psychiatric drug use and a history of falls, the QUS-C-derived BUA continued to be a significant independent factor for predicting the incidence of falls in elderly people. The OR of QUS-C-derived BUA was only 0.93 for our 443 participants, but its feasibility is practical for fall prevention in clinical practice. Because there were no significant demographic differences between the 1200 members of the randomly sampled cohort, our findings would be consistent even with a larger sample size. However, there was no significant correlation between QUS-C-derived BUA and the incidence of falls in the 40–70 year old group. It is possible that the incidence of falls in this age group was too low to be significant in a moderate sample size in this study. Nevertheless, our findings

Table 2 Logistic regression models of associated factors, including QUS-C, for incidence of falls in three different age groups

Variables	≥40 years old (n=443) OR (95% CI)	≥70 years old (n=94) OR (95% CI)	40–70 years old (n=349) OR (95% CI)
Nagelkerke R ² value	0.20	0.50	0.20
Age (years)	1.03 (0.99 to 1.08)	0.99 (0.78 to 1.26)	1.07 (0.99 to 1.16)
Sex (men=1, women=0)	0.15 (0.04 to 0.62)**	0.84 (0.04 to 6.36)	0.05 (0.01 to 0.44)**
Body height (cm)	1.04 (0.96 to 1.11)	1.10 (0.95 to 1.28)	1.04 (0.94 to 1.15)
Skeletal muscle mass (kg)	1.02 (0.94 to 1.11)	1.09 (0.87 to 1.37)	1.01 (0.91 to 1.12)
Current smoking (Yes=1, No=0)	4.97 (1.43 to 17.29)*	88.37 (2.66 to 2941.95)*	4.26 (0.86 to 21.07)
Current alcohol consumption (Yes=1, No=0)	3.13 (1.04 to 9.45)*	1.58 (0.08 to 33.18)	4.79 (1.21 to 18.99)*
Moderate exercise habit (Yes=1, No=0)	0.75 (0.30 to 1.87)	0.50 (0.03 to 7.73)	0.75 (0.26 to 2.15)
Living alone (Yes=1, No=0)	3.76 (1.18 to 11.93)*	19.47 (1.32 to 286.59)*	1.73 (0.18 to 16.90)
Psychiatric drug use (Yes=1, No=0)	1.56 (0.39 to 6.19)	46.80 (1.21 to 1806.38)*	0.61 (0.07 to 5.72)
Diabetes mellitus (Yes=1, No=0)	0.71 (0.15 to 3.43)	0.64 (0.04 to 11.75)	0.78 (0.09 to 7.06)
Hypertension (Yes=1, No=0)	0.99 (0.39 to 2.53)	1.44 (0.20 to 10.28)	0.40 (0.08 to 1.95)
Fracture history [†] (Yes=1, No=0)	2.02 (0.89 to 4.56)	2.76 (0.31 to 24.47)	2.88 (1.04 to 7.96)*
Fall history (Yes=1, No=0)	2.66 (1.20 to 5.89)*	4.06 (0.66 to 24.95)	2.74 (1.00 to 7.46)*
Anaemia (Yes=1, No=0) [†]	3.63 (1.49 to 8.84)**	6.70 (0.93 to 48.30)	2.95 (0.90 to 9.69)
Albumin (mg/dL)	0.61 (0.11 to 3.46)	0.22 (0.01 to 20.09)	1.40 (0.16 to 12.50)
QUS-C-derived BUA (dB/MHz)	1.00 (0.98 to 1.03)	0.93 (0.88 to 0.99)*	1.02 (0.99 to 1.06)

The incidence of falls was 29.0 per 1000 person-years in the ≥40 year old group, 41.0 per 1000 person-years in the ≥70 year old group, and 23.5 per 1000 person-years in the 40–70 year old group.

*p<0.05, **p<0.01.

[†]Anaemia in men if haemoglobin (Hb) <13 g/dL, and in women if Hb <12 g/dL.

BUA, broadband ultrasound attenuation; QUS-C, calcaneal quantitative ultrasound.

confirm a trivial but important role of QUS-C-derived BUA in predicting the incidence of future falls. QUS-C-derived BUA can be used to predict fall-related osteoporotic fractures,^{18–22} which might be partially mediated by its association with falls, as shown in Ou *et al*¹¹ and in the present study.

The mechanism that explains why a lower QUS-C-derived BUA yielded a higher incidence of falls is unclear. Deterioration of lower-limb stability is a predictor of future falls.³⁶ QUS-C-derived BUA has been proposed as a surrogate of muscle power and stability in the lower limbs, for example, quadriceps muscle strength.¹⁰ Bone status measured using QUS-C-derived BUA has also been positively associated with low gait speed in community-dwelling postmenopausal women.¹⁰ Poor mobility, however, is a risk factor for falls.³⁷ In our study, participants >70 years old did not even have moderate exercise habits; thus, the lower QUS-C-derived BUA might reflect lower-limb instability and relatively poorer mobility in elderly people. Hence, it is reasonable to claim an independent effect of QUS-C-derived BUA on the incidence of falls. However, additional investigations are required to confirm this finding. Practically, QUS-C-derived BUA is portable, cheap, non-radioactive and can be easily and safely used to prevent falls among community-dwelling individuals.

This study has some limitations. First, the participants were recruited from suburban and rural communities in central western Taiwan. The findings probably cannot be generalised to metropolitan or other communities.

Second, only QUS-C-derived BUA was used to determine the relationship with the incidence of falls. Theoretically, QUS-C-derived BUA from non-weight-bearing sites (radial or tibial) might have no relationship with falls. Whether the QUS-C-derived BUA measured using other brands or at different sites would yield similar findings is unknown and warrants further study.

Conclusion

We found that the incidence of falls in the 40–70 year old group was higher and had different associated risk factors than in the ≥70 year old group, and that QUS-C-derived BUA can be conveniently used to predict the incidence of falls that will lead to osteoporotic fractures in potential fallers >70 years old. Clinically, the more potential fallers who can be found before they fall, the more fall-prevention programmes can be efficiently applied. When using QUS-C-derived BUA to screen for osteoporosis, preventing falls might also be possible.

Author affiliations

¹Department of Family Medicine, Antai Medical Cooperation, Tien Sheng Memorial Hospital, Pingtung, Taiwan

²Department of Nursing, Meiho University, Pingtung, Taiwan

³College of Education, National Kaohsiung Normal University, Kaohsiung, Taiwan

⁴Departments of Family Medicine, National Cheng Kung University Hospital, Tainan, Taiwan

⁵Institutes of Gerontology, National Cheng Kung University College of Medicine, Tainan, Taiwan

⁶Departments of Internal Medicine, National Cheng Kung University Hospital, Tainan, Taiwan

⁷Department of Orthopedics, Tainan Municipal An-Nan Hospital of China Medical University, Tainan, Taiwan

⁸Institutes of Behavioral Medicine, National Cheng Kung University College of Medicine, Tainan, Taiwan

Acknowledgements We thank the staff working at the health promotion centre of the Dou-Liou branch of National Cheng Kung University Hospital for their administrative assistance, Pu-Hsian Guo for the SMM data, and Bill Franke for editing our English.

Contributors L-CO wrote the paper. C-HW wrote and revised the paper. L-CO and C-HW had the idea for the study and were involved in all aspects of this study. L-CO recruited the study participants. Y-FC, J-CC and C-SC helped interpret the data and made statistical suggestions. Z-JS, C-HW, T-HC and R-ML coordinated the study affairs and budget. All authors reviewed and approved the final version of this manuscript.

Funding National Cheng Kung University Hospital (NCKUH) (grant number: NCKUH-10102043).

Competing interests None declared.

Patient consent Obtained.

Ethics approval Institute Review Board of National Cheng Kung University Hospital (A-ER-100-347).

Provenance and peer review Not commissioned; externally peer reviewed.

Data sharing statement Extra data can be accessed via the Dryad data repository at <http://datadryad.org/> with the doi:10.5061/dryad.17tp1.

Open Access This is an Open Access article distributed in accordance with the Creative Commons Attribution Non Commercial (CC BY-NC 4.0) license, which permits others to distribute, remix, adapt, build upon this work non-commercially, and license their derivative works on different terms, provided the original work is properly cited and the use is non-commercial. See: <http://creativecommons.org/licenses/by-nc/4.0/>

REFERENCES

- Wu TY, Chie WC, Yang RS, *et al.* Factors associated with falls among community-dwelling older people in Taiwan. *Ann Acad Med Singap* 2013;42:320–7.
- Moyer VA, U.S. Preventive Services Task Force. Prevention of falls in community-dwelling older adults: U.S. Preventive Services Task Force recommendation statement. *Ann Intern Med* 2012;157:197–204.
- Elliott SA, Painter J, Hudson S. Living alone and fall risk factors in community-dwelling middle age and older adults. *J Community Health* 2009;34:301–10.
- Williams SB, Brand CA, Hill KD, *et al.* Feasibility and outcomes of a home-based exercise program on improving balance and gait stability in women with lower-limb osteoarthritis or rheumatoid arthritis: a pilot study. *Arch Phys Med Rehabil* 2012;91:106–14.
- Landi F, Liperoti R, Russo A, *et al.* Sarcopenia as a risk factor for falls in elderly individuals: results from the iSIRENTE study. *Clin Nutr* 2012;31:652–8.
- Kenkmann A, Price GM, Bolton J, *et al.* Health, wellbeing and nutritional status of older people living in UK care homes: an exploratory evaluation of changes in food and drink provision. *BMC Geriatr* 2010;10:28.
- Heuberger RA. The frailty syndrome: a comprehensive review. *J Nutr Gerontol Geriatr* 2011;30:315–68.
- Gerdhem P. Osteoporosis and fragility fractures. *Best Pract Res Clin Rheumatol* 2013;27:743–55.
- Bhatt T, Espy D, Yang F, *et al.* Dynamic gait stability, clinical correlates, and prognosis of falls among community-dwelling older adults. *Arch Phys Med Rehabil* 2011;92:799–805.
- Sakazaki T, Koike T, Yanagimoto Y, *et al.* Association between gait speed and bone strength in community-dwelling postmenopausal Japanese women. *Environ Health Prev* 2012;17:394–400.
- Ou LC, Sun ZJ, Chang YF, *et al.* Epidemiological survey of quantitative ultrasound in risk assessment of falls in middle-aged and elderly people. *PLOS ONE* 2013;8:e71053.
- Bischoff HA, Conzelmann M, Lindemann D, *et al.* Self-reported exercise before age 40: influence on quantitative skeletal ultrasound and fall risk in the elderly. *Arch Phys Med Rehabil* 2001;82:801–6.
- Stanaway FF, Cumming RG, Naganathan V, *et al.* Ethnicity and falls in older men: low rate of falls in Italian-born men in Australia. *Age Ageing* 2011;40:595–601.
- Kwan MM, Close JC, Wong AK, *et al.* Falls incidence, risk factors, and consequences in Chinese older people: a systematic review. *J Am Geriatr Soc* 2011;59:536–43.
- Wu TY, Chie WC, Yang RS, *et al.* Risk factors for single and recurrent falls: a prospective study of falls in community dwelling seniors without cognitive impairment. *Prev Med* 2013;57:511–17.
- de Vries OJ, Peeters GM, Lips P, *et al.* Does frailty predict increased risk of falls and fractures? A prospective population-based study. *Osteoporos Int* 2013;24:2397–403.
- Lee HD, Hwang HF, Lin MR. Use of quantitative ultrasound for identifying low bone density in older people. *J Ultrasound Med* 2010;29:1083–92.
- Kauppi M, Stenholm S, Impivaara O, *et al.* Fall-related risk factors and heel quantitative ultrasound in the assessment of hip fracture risk: a 10-year follow-up of a nationally representative adult population sample. *Osteoporos Int* 2014;25:1685–95.
- Marín F, González-Macías J, Díez-Pérez A, *et al.* Relationship between bone quantitative ultrasound and fractures: a meta-analysis. *J Bone Miner Res* 2006;21:1126–35.
- Youn T, Koval KJ, Kummer FJ, *et al.* Do all hip fractures result from a fall? *Am J Orthop* 1999;28:190–4.
- Ye F, Zhu SB, Wang X, *et al.* Operative treatment for proximal humeral fracture with rotator cuff tear in elderly patients. *Zhongguo Gu Shang* 2015;28:1111–13.
- Diaz-Garcia RJ, Oda T, Shauver MJ, *et al.* A systematic review of outcomes and complications of treating unstable distal radius fractures in the elderly. *J Hand Surg Am* 2011;36:824–35.
- NIH Consensus Development Panel on Osteoporosis Prevention, Diagnosis, and Therapy. Osteoporosis prevention, diagnosis, and therapy. *JAMA* 2001;285:785–95.
- Cosman F, de Beur SJ, LeBoff MS, *et al.* National Osteoporosis Foundation. Clinician's guide to prevention and treatment of osteoporosis. *Osteoporos Int* 2014;25:2359–81.
- Glüer CC, Eastell R, Reid DM, *et al.* Association of five quantitative ultrasound devices and bone densitometry with osteoporotic vertebral fractures in a population-based sample: the OPUS Study. *J Bone Miner Res* 2004;19:782–93.
- Guo PH, Sun ZJ, Ou LJ, *et al.* Epidemiological survey of the prevalence and associated risk factors of sarcopenia in middle-aged and old people. *Taiwan Geriatr Gerontol* 2013;8:27–45.
- Janssen I, Heymsfield SB, Baumgartner RN, *et al.* Estimation of skeletal muscle mass by bioelectrical impedance analysis. *J Appl Physiol* 2000;89:465–71.
- Silva CL, Lima-Costa MF, Firmo JO, *et al.* Hemoglobin level in older adults and the association with nutritional status and use of health services: the Bambuí Project. *Cad Saude Publica* 2012;28:2085–94.
- Corti MC, Guralnik JM, Salive ME, *et al.* Serum albumin level and physical disability as predictors of mortality in older persons. *JAMA* 1994;272:1036–42.
- Merom D, Mathieu E, Cerin E, *et al.* Social dancing and incidence of falls in older adults: a cluster randomised controlled trial. *PLoS Med* 2016;13:e1002112.
- Morrison A, Fan T, Sen SS, *et al.* Epidemiology of falls and osteoporotic fractures: a systematic review. *Clinicoecon Outcomes Res* 2013;5:9–18.
- Penninx BW, Pluijm SM, Lips P, *et al.* Late-life anemia is associated with increased risk of recurrent falls. *J Am Geriatr Soc* 2005;53:2106–11.
- Stalehoef PA, Diederiks JP, Knottnerus JA, *et al.* A risk model for prediction of recurrent falls in community-dwelling elderly: a prospective cohort study. *J Clin Epidemiol* 2002;55:1088–94.
- Thomas P, Chara T, Christos T, *et al.* Insomnia and its correlates in a representative sample of Greek population. *BMC Public Health* 2010;10:531.
- Wu LT, Blazer DG. Substance use disorders and psychiatric comorbidity in mid and later life: a review. *Int J Epidemiol* 2014;43:304–17.
- van Schooten KS, Pijnappels M, Rispens SM, *et al.* Daily-life gait quality as predictor of falls in older people: a 1-year prospective cohort study. *PLOS ONE* 2016;11:e0158623.
- Masumoto T, Yamada Y, Yamada M, *et al.* Fall risk factors and sex differences among community-dwelling elderly individuals in Japan. A Kameoka study. *Nihon Koshu Eisei Zasshi* 2015;62:390–401.