BMJ Open Protocol of a feasibility study for cognitive assessment of an ageing cohort within the Southeast Asia Community Observatory (SEACO), Malaysia

Devi Mohan,¹ Blossom C M Stephan,² Pascale Allotey,^{1,3} Carol Jagger,² Mark Pearce,² Mario Siervo,² Daniel D Reidpath^{1,3}

To cite: Mohan D, Stephan BCM, Allotey P, *et al.* Protocol of a feasibility study for cognitive assessment of an ageing cohort within the Southeast Asia Community Observatory (SEACO), Malaysia. *BMJ Open* 2017;**7**:e013635. doi:10.1136/bmjopen-2016-013635

► Prepublication history for this paper is available online. To view these files please visit the journal online (<http://dx.doi.org/10.1136/bmjopen-2016-013635>).

Received 27 July 2016
Revised 7 December 2016
Accepted 22 December 2016

For numbered affiliations see end of article.

Correspondence to

Dr Devi Mohan;
devi.mohan@monash.edu

ABSTRACT

Introduction: There is a growing proportion of population aged 65 years and older in low-income and middle-income countries. In Malaysia, this proportion is predicted to increase from 5.1% in 2010 to more than 15.4% by 2050. Cognitive ageing and dementia are global health priorities. However, risk factors and disease associations in a multiethnic, middle-income country like Malaysia may not be consistent with those reported in other world regions. Knowing the burden of cognitive impairment and its risk factors in Malaysia is necessary for the development of management strategies and would provide valuable information for other transitional economies.

Methods and analysis: This is a community-based feasibility study focused on the assessment of cognition, embedded in the longitudinal study of health and demographic surveillance site of the South East Asia Community Observatory (SEACO), in Malaysia. In total, 200 adults aged ≥50 years are selected for an in-depth health and cognitive assessment including the Mini Mental State Examination, the Montreal Cognitive Assessment, blood pressure, anthropometry, gait speed, hand grip strength, Depression Anxiety Stress Score and dried blood spots.

Discussion and conclusions: The results will inform the feasibility, response rates and operational challenges for establishing an ageing study focused on cognitive function in similar middle-income country settings. Knowing the burden of cognitive impairment and dementia and risk factors for disease will inform local health priorities and management, and place these within the context of increasing life expectancy.

Ethics and dissemination: The study protocol is approved by the Monash University Human Research Ethics Committee. Informed consent is obtained from all the participants. The project's analysed data and findings will be made available through publications and conference presentations and a data sharing archive. Reports on key findings will be made available as community briefs on the SEACO website.

INTRODUCTION

Cognitive ageing and dementia are global public health priorities. The number of people living with dementia globally is predicted to increase from 46.8 million in 2015 to 74.7 million in 2030 and 131.5 million in 2050; and more than 70% of them will be from the low-income and middle-income countries.¹ Malaysia is undergoing a rapid demographic transition. It is projected that, by 2050 the older aged population, defined as persons aged 65 years and older, will make up 15% of the country's population.² This change in age structure will lead to an unprecedented increase in the number of people living with cognitive impairment and dementia and will have significant personal, financial and social impacts.

Malaysia is a multiethnic country with wide cultural diversity.³ Also, it is a very fast growing and transitional economy. Indeed, by 2020 it is projected that Malaysia will transition to a high-income country. Owing to these reasons, the prevalence and risk factors of dementia and cognitive impairment may not be consistent with those reported in other world regions.

The prevalence estimates of dementia in developed countries range from 4.7% to 7.2%.⁴ Few prevalence studies of dementia and cognitive impairment have been conducted in Malaysia and most have been undertaken in small non-representative samples (table 1). No research has been conducted on incidence. Where assessed, the prevalence of cognitive impairment and dementia, typically defined using Mini Mental State Examination (MMSE) cut-off scores or the Elderly Cognitive Assessment

Table 1 Summary of findings from prevalence studies of dementia and cognitive impairment in Malaysia

First author (year)	Sample	N	Age	Neuropsychological assessment	Results
Krishnaswamy (1997) ⁵	3 urban settlements in Kuala Lumpur. No response rate given. Only Ethnic Malays were tested	522	≥60	2-stage screening. ECAQ (cut-off score ≤6) used for further testing using GSM for an AGECAQ diagnosis of dementia according to ICD-10 (in Kampung Baru only, N=400). Neurologist re-examined dementia patients a 3 months	6% dementia in people aged ≥65 and 12% in people aged ≥75 (sample size=323).
Sherina (2004) ⁶	Rural community of a village of Selangor state. 84.8% response rate	223	≥60	ECAQ (cut-off score ≤5)	22.4% cognitively impaired
Rashid (2006) ⁷	2 coastal villages in Kuala Muda in Kedah. No response rate given	92	≥60	ECAQ (cut-off score ≤4)	18.5% cognitive impairment
Al-Jawad (2007) ⁸	200-bed Government run residential home, Malaysia. 83.5% response rate	167	≥60	ECAQ (cut-off score ≤4, reflecting probable dementia)	36.5% dementia
Khaw (2009) ⁹	190-bed privately run care home, Penang Malaysia. 81.6% response rate	155	≥60	ECAQ (cut-off score ≤5)	12.3% dementia
Norlaily (2009) ¹⁰	Outpatient clinics of University Sains Malaysia Hospital (HUSM), east coast of Peninsular Malaysia. 0.100% response rate phase 1 and 83% phase 2 (N=39/47)	399	≥65	2-stage screening. Literate: MMSE (cut-off ≤17) and illiterate: ECAQ (cut-off ≤5). All impaired had a clinical interview based on DSM IV criteria	2.5% dementia
Hamid (2010) ¹¹	National, including all 13 Malaysian states and the Federal Territory of Kuala Lumpur. All community dwelling. 88% response rate	2980	≥60	GMS-B3 (AGECAT diagnosis of dementia)	14.3% dementia
Rashid (2012) ¹²	22 villages from Kedah, a north western state in Malaysia. 97.7% response rate	418	≥60	ECAQ (cut-off score ≤4, reflecting probable dementia)	11% cognitively impaired
Razali (2012) ¹³	Anteeds of the Medical and Nephrology Clinics at UKMMC. Response rate 96.9%	346	≥60	MMSE (MCI defined as impairment on delayed recall of verbal material, non-verbal material, or both, from the MMSE, with a total score of 21–28/30 on the MMSE)	64.7% MCI. 3.1% had MMSE ≤20
Arabi (2013) ¹⁴	2 primary care clinic in Cheras, Kuala Lumpur. 96.7% response rate	155	≥60	EDQ, MMSE ≤21	Prevalence of early dementia using the EDQ was 52.3%, and prevalence of dementia using MMSE was 15.3%.
Pu'un (2014) ¹⁵	Melanau population in Mikah (8 out of 32 villages), an indigenous population in Sarawak. 28% of elderly population screened	344	≥60	Two-stage screening. ECQA (cut-off score ≤5, reflecting probable dementia) further evaluation by clinical interview (using DSM-IV-TR criteria for dementia)	27.3% cognitive impairment and 10.5% dementia

AGECAT, Automated Geriatric Examination for Computer Assisted Taxonomy; BMI, body mass index; CVS, cardiovascular illnesses variable incorporating hypertension, diabetes mellitus, ischaemic heart disease, hyperlipidaemia and stroke; DSM, Diagnostic and Statistical Manual of Mental Disorders; DSM-IV-TR, Diagnostic and Statistical Manual of Mental Disorders, Fourth Edition (IV), Text Revision (TR); ECAQ, Elderly Cognitive Assessment Questionnaire; EDQ, Early Dementia Questionnaire; GSM, Geriatric Mental State Examination; ICD, International Classification of Diseases; MCI, Mild Cognitive Impairment; MMSE, Mini Mental State Examination.

Questionnaire (ECAQ), has been found to range from 2.5% to 36% (table 1).

Risk factors for cognitive impairment have included age, sex, educational attainment and ethnicity.^{5 6 8 9 12} However, the link between health and cognition has been poorly explored in the community-based settings. Results from studies that have explored health cognition associations have been mixed including no association between body mass index or hypertension and cognitive impairment¹² or dementia,⁹ an association between dementia and occupational pesticide exposure and history of stroke in a outpatients setting.¹⁰ In a community sample, cognitive impairment was not found to be associated with hypertension,⁷ while in another study dementia was found to be associated with cardiovascular illnesses defined as a combination variable including hypertension, diabetes mellitus, ischaemic heart disease, hyperlipidaemia and stroke.¹⁵ A more in-depth investigation of cardiovascular and metabolic risk factors in addition to assessment of other health and lifestyle factors, which are known risks for dementia; needs to be undertaken in the Malaysian population.

The main difficulties in conducting dementia studies in Malaysia have been limitations in assessment methods including a lack of availability of translations of commonly used cognitive screening tests, lack of normative data, literacy/educational attainment biases and lack of tests that are sensitive to ethnic differences (Malays, Chinese and Indians). This project will redress these through the development and piloting of a neuropsychological battery incorporating various cognitive assessment tools that have been validated in similar settings.

Knowing the burden of cognitive impairment and dementia and their risk factors in Malaysia will be necessary for the development of treatment and prevention strategies and to inform public health policies focused on an ageing population. Therefore, the objective of this study is to evaluate the feasibility of detailed cognitive and health assessment using a set of cognitive battery specifically designed to assess cognition in Malaysia. The long-term aim is to develop a comprehensive epidemiological research platform for investigating cognition and dementia in Malaysia, and, thereby, maximise the number of years free of cognitive impairment and dementia. This research platform will then provide a unique resource for studying ageing during economic transition and across ethnic and religiously diverse groups.

Objectives

- To assess the feasibility and acceptability of a tablet-based neuropsychological battery, to screen for cognitive impairment and dementia in individuals aged 50 years and over within a health and demographic surveillance site in Malaysia.
- To assess the feasibility and acceptability of collecting physical measurements including gait speed, hand

grip strength, single lead ECG and dried blood spots (DBS).

- To assess the feasibility of establishing an ageing cohort in order to monitor incidence of dementia, cognitive decline and other ageing-related health risks.

METHODS AND ANALYSIS

Participants

This study is embedded within the South East Asia Community Observatory (SEACO) population.¹⁶ SEACO is a health and demographic surveillance site (HDSS); a unique research platform in population health and well-being focused on a middle-income community in Segamat, Johor in the southern tip of the Peninsular Malaysia. The district of Segamat comprises urban, rural and plantation areas with an ethnic mix including Malays, Chinese and Indian, close to national proportions. SEACO runs its HDSS research platform in five subdistricts (mukims) of the Segamat district: Bekok, Chaah, Gemereh, Jabi and Sungai Segamat. In total, 38 228 individuals completed the baseline census (March 2012 to February 2013) that included information on sociodemographic status and self-reported health conditions. Data were collected by trained interviewers at the person's place of residence. Individuals have been reseen after a year (August 2013 to June 2014) for the health round, which collected a detailed health profile of all the consenting participants.

Individuals aged 50 years and over, who are living in Segamat Johor and have been assessed as part of SEACO in its previous health round form the study population. Nearly 9000 individuals meet these inclusion criteria. Baseline data from the census, update and health rounds are used to identify participants for inclusion. Persons are excluded if they cannot communicate effectively, have severe hearing loss or visual impairment (ie, are unable to hear normal conversation and/or not able to see the newsprint), are untraceable or are unable to give consent. From the eligible population, 200 samples are selected by stratified random sampling. The target is to recruit 200 participants stratified by age (50–59 and ≥60 years) and gender, with equal numbers in each group. Participants are also stratified based on their ethnicity (Malay, Chinese, Indian and Orang Asli) to ensure that each group is represented proportionately to the composition of the elderly population of SEACO as shown in figure 1. Stratified sampling is used in order, not to miss any of these strata in the relatively small sample, so that operational issues across all the groups are captured. The SEACO health round data base is used as the sampling frame.

Data collection

A home visit will be conducted to confirm eligibility, provide an information sheet, discuss any questions and to make an appointment for data collection. On the

Figure 1 Stratification grid used for selecting the feasibility study samples from the SEACO population. SEACO, South East Asia Community Observatory.

appointment date informed consent will be obtained. All interviews will be conducted in Bahasa Malay or Chinese according to the participant's choice. All data are collected on a Tablet (Samsung Galaxy Tab 3 V), unless otherwise specified. The assessment interview will take ~1.5–2 hours to complete. The ordering of the measures will be standardised for all participants and will include:

1. *Blood pressure (BP)*: resting BP will be measured in triplicate using an Omron electronic BP monitor. Participants will be resting in a seated position for at least 10 min before the measurement, and the three measurements will be taken 1 min apart.
2. *Anthropometry*: height and weight will be measured with a combined electronic scales and laser height measurement device. The participant will be barefoot and wearing light clothing.
3. *Neuropsychological battery*: the neuropsychological battery consists of four components:
 - 3.1. Subjective memory complaint: a single item will ask participants: Do you have problems with your memory? (Possible answers are: 'NO' OR 'YES')? If they answer yes they will be asked additional questions about memory loss in the previous year; and memory in comparison to other of same age
 - 3.2. Subjective memory symptom from the relative/caretaker: a close relative/caretaker will be asked a single question on problems with memory of the participant in the past year relative to previous years with possible answers: 'NO' OR 'YES'.
 - 3.3. MMSE: depending on the language chosen for interview by the respondent, the Malay (MMSE-7), or Chinese version will be used to screen for severe cognitive impairment. The Malay version was validated by Norlinah and colleagues among multiethnic elderly attending two hospitals in Kuala Lumpur.¹⁷ The Chinese version was validated among a Chinese population in Singapore.¹⁸ Scores range from 0 to 30,

with higher scores indicating better global cognitive functioning.

- 3.4. Montreal Cognitive Assessment tool (MoCA) will be used to screen for milder forms of cognitive impairment.¹⁹ The Singapore Chinese²⁰ or Bahasa Malay version²¹ will be used, based on the language of interview of the participants.

All the components of cognitive assessment will be administered using tablet, except MoCA, which will be administered in paper form. XLS Form will be used to create the questionnaire on the tablet, and this will be developed in consultation with the SEACO IT team. All MMSE questions, the tasks and the images (along with the rest of the questionnaire) will be incorporated into the form. Data collector will administer the tool and will enter the participant responses and scores for each question. The form will be programmed to calculate the total score (out of 30) for the participant once the data collector finishes administering the MMSE. This score will help the data collector to decide on need for referral.

4. *Functional performance*: two tests will be used to assess functional performance including: *handgrip strength* assessed in both arms (in duplicate) using a portable dynamometer; and, *gait speed*, which measures using a stop watch, the time it takes a participant to walk over a 4 m distance at their usual pace. The respondent will be asked to walk a distance of 6 m while the stop watch recording starts only at the end of the first 1 m and is stopped at the end of fifth meter of walk. In the field this is done using a foldable custom-made device colour coded for easy understanding. The participant will also be asked about the history of imbalance on walking.
5. *Health and lifestyle questionnaire*: participants will be asked to complete a questionnaire including the following:

- Information on self-rated health status: self-rated health will be measured by the question 'How would you rate your current health?' and the response will be captured in a five-point Likert scale of very poor, poor, neither poor nor good, good and very good.

- ▶ Global Activity Limitation Indicator (GALI)²² which underlies the European Healthy Life Years indicator includes a single item that asks ‘For the past 6 months at least, to what extent has you been limited because of a health problem in activities people usually do?’
 - ▶ Activities of daily living measured using Katz Index of Activities of Daily Living (adapted version).²³
 - ▶ Lifestyle factors including tobacco and alcohol use—tobacco use: the participant will be asked if he or she has ever smoked, history of current smoking, age of initiation, type of tobacco product used, number of cigarettes smoked and attempts to quit. Alcohol use: single item asking ‘Have you consumed alcohol in the past 30 days?’
 - ▶ Self-reported exhaustion—a single item asking ‘Have you felt that everything you did was an effort/you could not get going—for at least 3 days during the last week?’
 - ▶ Level of activity—two items asking: (1) ‘How often in the last week did you engage in moderate activities like gardening, cycling or going for a walk?’; and (2) ‘Do you engage in at least 150 min of moderate-intensity aerobic physical activity throughout the week?’
 - ▶ A history of diagnosed conditions such as stroke, ischaemic heart disease, transient ischaemic attack, heart failure, hypertension, diabetes (type 1 or type 2) and cancer.
 - ▶ Depression Anxiety Stress Scores.²⁴
6. *Dry blood spot collection and blood glucose testing*: four drops of blood will be collected on blood spot card (Whatman 903), under strict aseptic precaution. The DBS will be packed with desiccant packs and barcode labelled. The samples will be transferred to a freezer (−20°C) within a week of collection. The DBS will be used to determine glycated haemoglobin levels, lipid profile and C reactive protein levels. A drop of blood will also collected to check the random blood sugar using glucometer.
 7. *Single lead ECG for atrial fibrillation*: the AliveCor Mobile ECG records, displays, stores and transfers single channel rhythms. It records a 30 s tracing and gives the diagnosis as normal or possible atrial fibrillation. The ECG recording will be validated against a diagnosis by cardiologist who is blinded to the diagnosis given by the app.

Data collector training and supervision

SEACO is an ISO certified research platform for its operations and hence training and supervision are carried out in accordance with SEACO’s standard operating procedures. All data collectors are required to be fluent in the local language used for interview (Bahasa Malay or Mandarin) and have at least a working knowledge of English. To ensure data quality and consistency across interviewers, the field supervisor will undertake

random concurrent supervisory visits to observe practice. The data collectors will also be periodically observed by the project leader and the field manager. Based on the scoring done during these sessions, onsite training or retaining will be carried out as appropriate.

ETHICS AND DISSEMINATION

The study protocol was approved by Monash University Human Research Ethic Committee (CF15/4367—2015001885). Informed consent will be obtained from all eligible participants through signed or video consent. Separate consent will be obtained from the relative before interviewing them on the participants’ cognitive status. All participants will receive feedback from their interview which includes information about their height, weight, BP, MMSE score, glucometer blood sugar level and ECG diagnosis; with specific mention of the items that require a physician consultation and evaluation. The cut-offs used for referrals across the different measures are shown in [table 2](#). All data and samples collected in the study will be delinked prior to analysis to ensure that anonymity is maintained.

The project data and findings will be made available through publications and conference presentations. There is also a searchable data archive under development that would ensure data sharing. Reports will be made available as community briefs on the SEACO website.

DISCUSSION

What is unique about the study?

Malaysian economy is growing and it is predicted that by 2020 Malaysia will transition to becoming a high-income country. Further, Malaysia is a multiethnic country. Therefore, the proposed project will provide a unique resource for studying ageing during economic transition and across ethnic and religiously diverse groups. Risk factors and disease associations may not be consistent with those reported in other world regions where the focus has been on Caucasian populations in a higher income context, and with access to higher quality

Table 2 Cut-offs used for referral of participants in the study across the different measures

Measurements	Cut-off used for referral for further medical evaluation
BP	At least one of the three systolic BP recording ≥ 140 mm of Hg and/or at least one of the three diastolic BP recording ≥ 90 mm of Hg
MMSE score (out of a total of 30)	≤ 20
Glucometer random blood sugar	≥ 11.1 mmol/L
Single lead ECG diagnosis	Possible atrial fibrillation
BP, blood pressure; MMSE, Mini Mental State Examination.	

healthcare and state of the art health technologies for diagnostics and treatment. The study provides the opportunity to evaluate these in the Malaysian context which is more typical of countries within the South East Asian region. The study can also draw on the extensive sociodemographic and health data infrastructure as well as access to populations facilitated through the SEACO community engagement process. This will enhance the response and retention rates towards the establishment of a sustained cohort of elderly people for the long-term assessment of cognitive function in Malaysia.

Why is the study important?

Knowing the burden of cognitive impairment and dementia and risk factors for disease in Malaysia will be necessary for the development of treatment and prevention efforts but it is also important to view these alongside continued increases in life expectancy through population-level indicators such as healthy life expectancy. A similar non-Caucasian resource is the 10/66 Research Study, a multicountry study that explores dementia and ageing in low-income and middle-income countries.²⁵ While there are some data on variability in dementia prevalence and its risk factors (eg, level of educational attainment), questions about ethnic and cultural variation remain unanswered. Our study will therefore complement and extend the work undertaken by the 10/66 Dementia Research Group. It will also provide the opportunity to develop a range of cross-culturally validated tools. The study will provide a comprehensive epidemiological research platform for investigating cognition and dementia in Malaysia, and, in the longer term, to maximise the number of years free of cognitive impairment and dementia.

Author affiliations

¹Global Public Health, Jeffrey Cheah School of Medicine and Health Sciences, Monash University, Bandar Sunway, Malaysia

²Newcastle University Institute for Ageing & Institute of Health and Society, Newcastle University, UK

³South East Asia Community Observatory (SEACO), Monash University, Segamat, Malaysia

Acknowledgements The authors acknowledge the SEACO field operations team for their support in designing this study.

Contributors DM, BCMS and PA contributed to the drafting of this manuscript. CJ, MP, MS and DDR provided inputs in initial designing of the protocol and in final reviewing of the manuscript.

Funding Funding for SEACO is provided by the Research Offices of Monash University campuses in Australia and Malaysia; the Faculty of Medicine, Nursing and Health Sciences; the Jeffrey Cheah School of Medicine and Health Sciences; and the Faculty of Arts. Ethics approvals for the establishment of SEACO and for this protocol have been obtained from the Monash University Human Research Ethics Committee. This feasibility study is funded through a seed grant from the Monash-Newcastle Partnership Grants.

Competing interests None declared.

Ethics approval Monash University Human Research Ethics Committee.

Provenance and peer review Not commissioned; externally peer reviewed.

Data sharing statement The data generated from the study are encrypted and backed up by servers at Monash Sunway Campus and at Monash Clayton. Research staff have limited access to anonymised data. There is a searchable data archive under development that would ensure data sharing. It is anticipated that in the medium term, limited de-identified public use data sets will be made available for secondary analysis of the data.

Open Access This is an Open Access article distributed in accordance with the Creative Commons Attribution Non Commercial (CC BY-NC 4.0) license, which permits others to distribute, remix, adapt, build upon this work non-commercially, and license their derivative works on different terms, provided the original work is properly cited and the use is non-commercial. See: <http://creativecommons.org/licenses/by-nc/4.0/>

REFERENCES

1. Prince MJ. *World Alzheimer Report 2015: the global impact of dementia: an analysis of prevalence, incidence, cost and trends*. 2015.
2. Nations U. *World population ageing: 1950–2050*. New York: Department of Economic and Social Affairs PD. 2001. Contract No.: ST/ESA/SER.A/207.
3. Department of Statistics M. *Population distribution and basic demographic characteristic report 2010 Putrajaya*. DOSM, 2011 29 July 2011. Report No.
4. Prince M, Bryce R, Albanese E, *et al*. The global prevalence of dementia: a systematic review and metaanalysis. *Alzheimers Dement* 2013;9:63–75.e2.
5. Krishnaswamy S, Kadir K, Ali RA, *et al*. Prevalence of dementia among elderly Malays in an urban settlement in Malaysia. *Neurol J Southeast Asia* 1997;2:159–62.
6. Sherina M, Rampal L, Mustaqim A. Cognitive impairment among the elderly in a rural community in Malaysia. *Med J Malaysia* 2004;59:252–7.
7. Rashid AK, Narayan KA, Azizah AM. The prevalence of cognitive impairment and depression and their associated factors in an elderly population in two villagers in Kedah. *Malays J Public Health Med* 2006;6:29–37.
8. Al-Jawad M, Rashid A, Narayan K. Prevalence of undetected cognitive impairment and depression in residents of an elderly care home. *Med J Malaysia* 2007;62:375.
9. Khaw C, Teo C, Rashid A. Cognitive impairment and depression among residents of an elderly home in Penang, Malaysia. *Internet J Psychiatry* 2009;1:1–9.
10. Norlaili H, Azidah AK, Asrenee AR, *et al*. Proportion of dementia and its associated factors among elderly patients attending outpatient clinics of Universiti Sains Malaysia Hospital. *Med J Malaysia* 2009;64:140–5.
11. Hamid TA, Krishnaswamy S, Abdullah SS, *et al*. Sociodemographic risk factors and correlates of dementia in older Malaysians. *Dement Geriatr Cogn Disord* 2010;30:533–9.
12. Rashid AK, Azizah AM, Rohana S. Cognitive impairment among the elderly Malays living in rural Malaysia. *Med J Malaysia* 2012;67:186–9.
13. Razali R, Baharudin A, Jaafar NRN, *et al*. Factors associated with mild cognitive impairment among elderly patients attending medical clinics in Universiti Kebangsaan Malaysia Medical Centre. *Sains Malaysiana* 2012;41:641–7.
14. Arabi Z, Aziz NA, Aziz AFA, *et al*. Early Dementia Questionnaire (EDQ): a new screening instrument for early dementia in primary care practice. *BMC Fam Pract* 2013;14:49.
15. Pu'un BI, Othman Z, Drahtman I. Dementia among elderly Melanau: a community survey of an indigenous people in East Malaysia. *Int Med J* 2014;21:1–4.
16. South East Asia Community Observatory. 2016 (cited 21 July 2016). <http://www.seaco.asia>
17. Ibrahim NM, Shohaimi S, Chong HT, *et al*. Validation study of the Mini-Mental State Examination in a Malay-speaking elderly population in Malaysia. *Dement Geriatr Cogn Disord* 2009;27:247–53.
18. Ng TP, Niti M, Chiam PC, *et al*. Ethnic and educational differences in cognitive test performance on mini-mental state examination in Asians. *Am J Geriatr Psychiatry* 2007;15:130–9.
19. Nasreddine ZS, Phillips NA, Bedirian V, *et al*. The Montreal Cognitive Assessment, MoCA: a brief screening tool for mild cognitive impairment. *J Am Geriatr Soc* 2005;53:695–9.
20. Chen C, Dong Y, Merchant R, *et al*. The Montreal cognitive assessment (MoCA) is superior to the mini-mental state examination (MMSE) in detecting patient's with moderate cognitive impairment,

- no-dementia (CIND) and at high risk of dementia. *Alzheimers Dement* 2011;7:S240–1.
21. Sahathevan R, Mohd Ali K, Ellery F, *et al.* A Bahasa Malaysia version of the Montreal Cognitive Assessment: validation in stroke. *Int Psychogeriatr* 2014;26:781–6.
 22. Berger N, Van Oyen H, Cambois E, *et al.* Assessing the validity of the Global Activity Limitation Indicator in fourteen European countries. *BMC Med Res Methodol* 2015;15:1.
 23. Shelkey M, Wallace M. Katz Index of Independence in Activities of Daily Living (ADL) (try this: best practices in nursing care to older adults). *Medsurg Nurs* 2002;11:100–2.
 24. Henry JD, Crawford JR. The short-form version of the Depression Anxiety Stress Scales (DASS-21): construct validity and normative data in a large non-clinical sample. *Br J Clin Psychol* 2005;44:227–39.
 25. Group DR. The 10/66 Dementia Research Group. 2008–2015 (cited 21 July 2016). <http://www.alz.co.uk/1066/>