BMJ Open Month of birth and risk of autism spectrum disorder: a retrospective cohort of male children born in Israel

Hila Shalev,¹ Ido Solt,^{1,2} Gabriel Chodick^{3,4}

To cite: Shalev H, Solt I, Chodick G. Month of birth and risk of autism spectrum disorder: a retrospective cohort of male children born in Israel. *BMJ Open* 2017;**7**:e014606. doi:10.1136/bmjopen-2016-014606

► Prepublication history and additional material are available online. To view these files please visit the journal online (<http://dx.doi.org/10.1136/bmjopen-2016-014606>).

Received 13 October 2016
Revised 27 March 2017
Accepted 5 April 2017

ABSTRACT

Background Increased incidence and prevalence of autism spectrum disorder (ASD) over the last two decades have prompted considerable efforts to investigate its aetiological factors. We examined an association between month of birth and ASD incidence.

Methods In a retrospective cohort of male children born from January 1999 to December 2008 in a large health organisation in Israel (Maccabi Healthcare Services), ASD was followed from birth through December 2015.

Results Of 108 548 boys, 975 cases of ASD were identified. The highest rates (10.3 and 10.2 per 1000 male live births) were recorded for children born in May and August, respectively, and the lowest rates for February (7.6 per 1000 male live births). Among lower socioeconomic status households, boys born in August were more likely (OR=1.71; 95% CI 1.06 to 2.74) of being diagnosed with ASD than children born in January. Significantly higher rates were not observed for other months.

Conclusions In line with several previous studies, we found a modestly higher likelihood of autism occurrence among male children of lower socioeconomic levels born in August.

INTRODUCTION

Autism spectrum disorder (ASD) is a biologically based neurodevelopmental condition characterised by (A) deficits in social communication and social interaction, and (B) restricted repetitive patterns of behaviour, interests and activities.¹ The prevalence of ASD has been rising² globally, and in Israel in particular.³

Analyses of season of birth or conception can provide clues about causes of diseases with unknown aetiology as in ASD. Patterns of seasonal variation in births have been found in several disorders; schizophrenia, mood disorders, epilepsy, language disorders, attention deficit hyperactivity disorder and neurodevelopmental disabilities.^{4–9} An association between season of birth or conception suggests periodicity of an environmental aetiological factor active during the prenatal, perinatal or early postnatal phase of development.¹⁰ The seasonal effects observed in

Strength and limitations of this study

- Validated source of information of autism spectrum disorder (ASD) diagnosis.
- The use of systematic data that were collected prospectively on a large population.
- No information was available regarding ASD disease severity.
- Only boys were included, contrasting with other studies that included both sexes.
- Lack of information in our population on influenza infections and vaccines.

these disorders led researchers to examine whether the same would apply to ASD. The findings of such investigations are inconsistent; several studies claimed increased risk of ASD among individuals born or conceived in various seasons^{11–13} or specific months,^{14–21} while other reports found no seasonal patterns.^{14 22–24} Inconsistencies in the literature could be due, in part, to a wide range of study populations, geographical areas, case definitions, comparison groups and analytical approaches.

In one of the first studies on season of birth and ASD to be carried out in a region other than the USA and western Europe, a study in Israel¹⁸ found an increase in March and August births among 188 ASD cases when compared with the general population. The replication of seasonal patterns in a region with a climate and environment different from those of the countries previously studied supports the generalisability of the findings. However, a later examination of all ASD cases (n=211) among Jews in Israel over 5 years revealed no seasonal patterns in birth period. Such inconsistency can be explained by methodological differences, selection of comparison group, lack of statistical power and use of different diagnostic criteria.

The objective of the present study was to examine a relationship between autism and seasonality in a large population in Israel,

¹Faculty of Medicine, Technion, Haifa, Israel

²Rambam Health Care campus, Haifa, Israel

³Maccabi Healthcare Services, Tel Aviv, Israel

⁴Sackler Faculty of Medicine, Tel Aviv University, Tel Aviv, Israel

Correspondence to

Prof. Gabriel Chodick;
hodik_g@mac.org.il

using the computerised databases of a single health provider and its standardised coding system. Male to female ratio among patients with ASD is reported between 3 to 1 and 4 to 1,^{25–27} reflecting a disorder of potential differences in mechanisms and aetiology between the two sexes.^{28–31} We excluded the girls from this investigation as we were interested in controlling the impact of sex by restricting the sample to boys.

METHODS

Settings

The 1994 National Health Insurance Law mandates compulsory universal health coverage for all Israeli residents in a health maintenance organisation (HMO) of their choice. Israeli citizens are obliged to choose one of four non-for-profit HMO's in the Israeli medical system, who provide equivalent medical services that are based on national medical standards. For this review, we examined data on ASD diagnosis from the computer database of the second-largest HMO, Maccabi Healthcare Services (MHS), which provides care to 2 million people (31% under 18 years of age) (Israeli Central Bureau of Statistics).

Study population

Using the computerised databases of MHS, we identified a total of 147 171 male children who were born between January 1999 and December 2008 (approximately 20% of all 736 thousand live male births in Israel during the same years). Children's parents were identified using data on household members in MHS. Children with missing paternal data (mostly due to previously belonging to a different health provider) were excluded from the study. Comparison of excluded ($n=38\,875$) and eligible ($n=110\,093$) children showed no statistically significant ($p>0.05$) differences in mean birth weight (3271 g and 3315 g, respectively), mean gestational week (39.0 weeks and 39.1 weeks) or mean mother's age at birth (29.4 years and 30.5 years) and a similar distribution of socioeconomic status (SES). To avoid inaccurate paternal identification, an additional 1545 children were excluded because more than one man was registered in the child's household. Hence, the total study population was 108 548 boys of 83 452 households.

Case identification and matching

A case was defined as a child with a physician-recorded diagnosis of autistic disorder, Asperger's syndrome or pervasive developmental-not otherwise specified (PDD-NOS), based on the Diagnostic and Statistical Manual of Mental Disorders, Fifth Edition (American Psychiatric Association 1994). In MHS, children with suspected ASD are referred to child development centres for a multidisciplinary assessment for diagnosis validation. The MHS registry of patients with ASD and validation processes were previously described.³²

Other study variables

SES was assessed according to the poverty index of the child's address enumeration area, as defined by the 2008

national census.³³ The Poverty Index is based on several parameters including household income, educational qualifications, crowding, material conditions, and car ownership, census of population and housing. SES levels range between 1 (lowest) to 10 (highest). In this study SES below median (6) was considered as low SES. Calendar year and month of conception was estimated based on the child's date of birth and the length of gestation as reported in the birth certificate. We categorised season of birth (and conception) as winter comprising December, January and February; spring, March, April and May; summer, June, July and August; and fall, September, October and November. Month of conception was calculated by subtracting gestational week from the date of birth.

Statistical analyses

First, bivariate analyses were conducted between month of birth and other covariates previously reported to be associated with increased risk of autism.^{34 35} The covariates included gestational age in weeks as a continuous variable,³⁶ birth weight, older maternal or paternal age (>35 years), and SES level above median (>6). Gestational age was available for 97 264 individuals (90% of the study population). Missing gestational ages were imputed using multiple imputation (five imputations) with birth weight as a covariate. The Cochran-Armitage test was used to assess a significant trend in the association between month of birth and ASD.

Multivariable logistic regression analyses were performed to determine the OR and 95% CI for ASD by month of birth, using January as the reference month. For the purpose of comparison with previous studies, we fitted alternative logistic regression models, with season of birth, month and season of conception (rather than birth), first month of the second trimester (see online supplementary appendix table a) and third trimester (see online supplementary appendix table b) as the main independent variables. Additionally, to evaluate the potential effect modification by socioeconomic level, the study population was divided into two groups in further analyses: low-medium (1–6) and medium-high (7–10). Effect modification by month of birth and SES was investigated by including a cross-product interaction term between the exposure variable (birth in August $\times$ SES group). A p value for interaction³⁷ was computed to determine the statistical significance of adjusted ORs related to birth in August between high and low SES levels using WINPEPI statistical package.³⁸

In a sensitivity analysis, using birth month and season, we excluded one calendar year at a time to identify the effect of one specific year, with no significant results.

A two-tailed p value <0.05 was considered statistically significant. All data processing and statistical analyses were performed using IBM SPSS Statistics for Windows, V.22.0. (IBM Corp. Armonk, New York, USA).

RESULTS

The mean number of monthly births ranged from 8318 in February to 9654 in August. In bivariate analyses, number

of births, maternal age, paternal age, birth weight, gestational length and SES were not associated with the month of birth (table 1). Of the 108 548 boys born during the study period, 975 were diagnosed with ASD, representing an incidence rate of 9.0 per 1000 live births. Mean age at ASD diagnosis was 3.0 years (SE=0.07 years). The highest incidences of ASD were recorded among children born in May and August: 10.3 and 10.2 per 1000, respectively, while the lowest incidences were in January and February (7.6 and 8.1 per 1000, respectively) (table 2). The results by month of conception are shown in table 3.

In a multivariable model, birth months of August and May were associated with a non-significantly ($p=0.16$ and $p=0.14$, respectively) increased OR (table 4). When the data were stratified by household socioeconomic level, the birth month of August in low SES was associated with an OR of 1.71 (95% CI 1.06 to 2.74) of ASD. No other significant results were observed for other birth months or in children of higher SES (table 5 and see online supplementary tables a,b,c,d,e). There was a borderline effect modification between birth in August and SES level ($p=0.07$). Using similar multivariable analyses, we found no significant ($p>0.5$) association between ASD and season of birth (see online supplementary appendix table c), season of conception (see online supplementary appendix table d), or month of conception (supplementary file 1).

DISCUSSION

In this retrospective cohort of boys born in Israel during a period spanning 10 years, children of low SES together with birth in August were associated with an increased likelihood of being diagnosed with ASD. This is the one of the largest studies outside the USA to describe the relationship between ASD incidence and month of birth.

The literature with respect to season of birth and risk of ASD is inconsistent. Of the 15 studies (total of 31 777 births) summarised in table 6 (all conducted in the northern hemisphere) 5 studies (total $n=23\,355$) reported an increased risk of ASD among individuals born in August. March was significantly associated with increased risk in eight studies (total $n=3008$), and three studies (total $n=5839$) found no significant association between month of birth and ASD occurrence. No other specific birth month was repeatedly reported to be associated with ASD.

There are several mechanisms by which season of birth may affect the developing brain, either directly on the fetus and newborn or indirectly through the mother during pregnancy or the postpartum period. Seasonal environmental factors that could be associated with ASD include viral or other infections,^{39 40} nutritional factors⁴¹ and vitamin deficiencies.⁴² The outcome of exposure to prenatal viral infection depends on many factors: the variation in susceptibility of the maternal and fetal host, including the maternal immune status, the infecting virus, the strain of virus, the developmental stage of the

Table 1 Study population characteristics by month of birth

Maternal age	Paternal age				Birth weight, g				Gestational age				SES 1 to 10				Births	Month
	M (SD)	Med	25%	75%	M (SD)	Med	25%	75%	M (SD)	Med	25%	75%	M (SD)	Med	25%	75%		
30.1 (4.9)	30	27	33	36	3296 (525)	3320	3000	3630	39.1 (1.9)	39	38	40	6.1 (2.3)	6	4	8	9367	January
30.2 (4.9)	30	27	34	36	3296 (528)	3320	3000	3640	39.0 (2.0)	39	38	40	6.2 (2.3)	6	5	8	8318	February
30.2 (4.9)	30	27	34	36	3305 (519)	3340	3020	3650	39.0 (1.9)	39	38	40	6.1 (2.3)	6	4	8	9038	March
30.3 (4.9)	30	27	34	37	3310 (519)	3340	3030	3640	39.1 (1.9)	39	38	40	6.2 (2.3)	6	5	8	8465	April
30.3 (4.9)	30	27	34	37	3300 (532)	3330	3010	3640	39.1 (1.9)	39	38	40	6.3 (2.3)	6	5	8	8711	May
30.3 (5.0)	30	27	34	37	3306 (538)	3350	3010	3650	39.0 (2.0)	39	38	40	6.2 (2.3)	6	5	8	8523	June
30.2 (4.9)	30	27	34	37	3304 (528)	3330	3010	3650	39.0 (2.0)	39	38	40	6.1 (2.3)	6	4	8	9330	July
30.3 (4.9)	30	27	34	37	3312 (532)	3340	3020	3660	39.0 (1.9)	39	38	40	6.1 (2.3)	6	4	8	9654	August
30.3 (4.9)	30	27	34	37	3291 (537)	3320	3000	3640	39.0 (1.9)	39	38	40	6.1 (2.3)	6	4	8	9270	September
30.3 (5.0)	30	27	34	37	3302 (524)	3330	3020	3640	39.1 (1.9)	39	38	40	6.1 (2.3)	6	4	8	9521	October
30.3 (5.0)	30	27	34	37	3302 (532)	3330	3000	3640	39.0 (1.9)	39	38	40	6.1 (2.3)	6	4	8	9069	November
30.3 (5.0)	30	27	34	37	3288 (540)	3320	3000	3630	39.0 (2.0)	39	38	40	6.1 (2.3)	6	4	8	9282	December
30.3 (5)	30	27	33.9	33.0	3642.5	3330.8	3010	3301	39.0 (2.0)	39	38	40	6.1 (2.3)	6	4.33	8	9046	Mean
				(5.7)				(529.5)										

SES, socioeconomic status.

Table 2 Incidence rate and 95% confidence interval (CI) per 1000 male live births of autism spectrum disorder (ASD), by month of birth

Month of birth	ASD cases	Rate (per 1000)	95% CI*
January	76	8.1	6.44 10.09
February	63	7.6	5.87 9.62
March	74	8.2	6.48 10.21
April	74	8.7	6.92 10.90
May	90	10.3	8.36 12.62
June	78	9.2	7.29 11.35
July	89	9.5	7.71 11.67
August	98	10.2	8.29 12.30
September	80	8.6	6.89 10.67
October	85	8.9	7.18 10.97
November	84	9.3	7.44 11.40
December	84	9.0	7.27 11.14
Total	975	8.96	7.17 11.07

*See text for details.

fetus, the amount of virus reaching the fetus—particularly the central nervous system and immune system, the route of access, genetics and probably other factors.⁴³ Maternal viral infection has been cited as the ‘principal non-genetic cause of autism’.⁴⁴ Maternal infections (ie, rubella, cytomegalovirus, influenza), prolonged fever and maternal inflammation cause changes in a variety of inflammatory cytokines that have been associated with ASD.^{45 46} Depending on the location of immune activation, persistent viral infection could lead to chronically elevated cytokine levels that can be produced directly in the brain or gain access to the CNS by crossing an immature blood-brain barrier, altering CNS development.⁴⁷

A few studies found influenza infection during pregnancy to be a factor that increases risk for infantile autism^{39 48} especially in the first trimester.⁴⁶ Results from animal models also support an effect of maternal influenza infection on the neurodevelopment of the offspring.^{49 50} In this analysis, ASD was related significantly with month of birth but not with month or season of conception and therefore our results do not support this potential aetiology. This is in line with recently published results of a large cohort from the USA.⁵¹ Potential explanations can be proposed for the observed effect modification of increased risk of ASD among August born of lower SES. These may include higher likelihood

Table 3 Incidence rate and 95% confidence interval (CI) per 1000 male live births of autism spectrum disorder (ASD), by month of conception

Month of conception	ASD cases	Rate (per 1000)	95% CI*
January	87	9.2	7.4 11.3
February	74	8.7	6.9 11
March	77	8.3	6.6 10.4
April	72	8.0	6.3 10.1
May	77	8.4	6.7 10.5
June	68	7.8	6.1 9.9
July	72	8.2	6.4 10.3
August	89	10.2	8.2 12.5
September	91	10.9	8.8 13.4
October	79	8.4	6.6 10.4
November	99	10.8	8.8 13.1
December	87	9.1	7.3 11.2
Total	972	9.0	8.4 9.9

*See text for details.

Table 4 Crude and adjusted odds ratio (OR) and 95% confidence interval (CI) of autism spectrum disorder (ASD) for month of birth: multivariable logistic regression

Month of birth	OR crude	95% CI		OR adjusted*	95% CI	
January	1 (ref.)			1 (ref.)		
February	0.93	0.67	1.31	.92	.66	1.29
March	1.01	0.73	1.39	1.00	0.73	1.38
April	1.08	0.78	1.49	1.08	0.78	1.48
May	1.28	0.94	1.73	1.26	0.92	1.71
June	1.13	0.82	1.55	1.11	0.81	1.52
July	1.16	0.86	1.58	1.15	0.84	1.56
August	1.25	0.93	1.69	1.24	0.92	1.68
September	1.07	0.78	1.46	1.05	0.77	1.44
October	1.10	0.81	1.50	1.09	0.80	1.49
November	1.13	0.83	1.55	1.12	0.82	1.53
December	1.11	0.81	1.51	1.09	0.80	1.50

*Adjusted for child's age, father's age, mother's age, district, birth weight, gestational age with imputation (see text for details).

of maternal exposure to infections or particulate matter air pollution that have been connected to increased risk of ASD.⁵² A role of vitamin D deficiency in the development of autism has been proposed.⁴² Thus, seasonal fluctuations in vitamin D should also be considered. In the northern hemisphere, vitamin D levels are the lowest in winter time;^{53–55} however, a study of healthy volunteers from Israel found no differences in vitamin D deficiency rates between summer and winter.⁵⁶

Strengths of the current study include the investigation of a large and unselected population that is similar in its characteristics (mean maternal age, years of follow-up, etc) to a large cohort of a recently published US study that examined autism and seasonality,⁵¹ the cohort design,

the use of systematic data that were collected prospectively, validated source of information of ASD diagnosis, the availability of data on prematurity and the objective assessment of SES. Some limitations should be discussed. We had no information about the ASD disease severity; in one previous report,⁵⁷ high functioning children with ASD showed an increase in summer births compared with the low functioning children with ASD. In addition, SES assessment was conducted using data on current residential area, which is not necessarily the residential area during the time of pregnancy, nor necessarily identical to individual SES. The children who were born at the end of the year were younger than those born earlier in the year, but this is negligible because they were all at

Table 5 Crude and adjusted odds ratios (ORs) and 95% confidence interval (CI) from multivariable logistic regression of autism spectrum disorder (ASD) for month of birth, by socioeconomic level (SES)

Month of birth	SES low						SES high					
	OR crude	95% CI		OR adjusted*	95% CI		OR crude	95% CI		OR adjusted*	95% CI	
January	1 (ref.)			1 (ref.)			1 (ref.)			1 (ref.)		
February	1.01	0.58	1.75	1.01	0.58	1.75	0.89	0.58	1.35	0.88	0.57	1.34
March	1.34	0.80	2.22	1.34	0.81	2.22	0.82	0.54	1.25	0.81	0.53	1.24
April	1.30	0.77	2.19	1.32	0.78	2.22	0.93	0.62	1.40	0.93	0.62	1.40
May	1.31	0.78	2.20	1.30	0.78	2.19	1.19	0.81	1.74	1.17	0.80	1.72
June	1.26	0.75	2.12	1.25	0.74	2.11	1.01	0.67	1.51	0.99	0.66	1.49
July	1.37	0.83	2.25	1.36	0.82	2.25	1.01	0.68	1.49	0.99	0.67	1.48
August	1.70	1.06	2.73	1.71	1.07	2.76	1.00	0.67	1.48	0.99	0.66	1.46
September	1.32	0.80	2.19	1.32	0.80	2.18	0.92	0.61	1.38	0.91	0.60	1.36
October	1.24	0.75	2.05	1.24	0.75	2.07	1.03	0.69	1.52	1.01	0.68	1.50
November	1.45	0.88	2.37	1.46	0.89	2.40	0.96	0.64	1.44	0.93	0.62	1.40
December	1.49	0.91	2.43	1.51	0.92	2.46	0.89	0.59	1.34	0.88	0.58	1.32

*Adjusted for child's age, father's age, mother's age, district, birth weight, age with imputation (see text for details).

Table 6 Summary of findings in previous studies investigating the relationship between seasonal variations of birth and autism

Ref.	Study period	Place	ASD cases	Months of increased risk of ASD
11	1990–2002	California	19238	November, December, August, September, October
14	1992–2000	California	2940	August*
14	1997–1999	California	3897	No association
22	1947–1980	UK	1631	No association
15	1983–2002	Maryland	1068	Spring
16	1981***	North Carolina	810	March, August
23	1947–1992	Child dev. clinic	620	No association**
17	1945–1980	Denmark	328	March
24	1980–1985	Israel	211	No association
18	1964–1986	Israel	188	March and August
12	1980s	Ontario , Canada	179	Summer and spring
19	1977–1985	USA	175	March
20	1962–1984	Sweden	174	March
13	1972–1978	Japan	132	Spring–summer (April, May, June)
21	1990–1992	UK	86	Spring

*Conception in November (for years 1994–1996).

**($p < 0.1$ for middle of August for verbal ASD).

***Year of publication.

ASD, autism spectrum disorder.

the age of at least 6 years when the study was done and most autistic children are diagnosed by this age.⁵⁸ Our analysis included only boys and therefore generalisability to both sexes is limited. Data were unavailable regarding important risk factors such as family history, viral infections and environmental factors. Further investigations on whether the observed seasonal patterns interplay with genetic and environmental risk factors in the development of ASD may provide clues regarding the aetiology of these complex disorders.

CONCLUSIONS

In line with several previous studies, we found modestly higher likelihood of ASD occurrence among male children of lower SES levels born in August. Month of birth may indicate seasonal causal factors that could be associated with autism. The fact that these patterns appear to differ by SES may also be aetiologically significant.

Correction notice This article has been corrected since it was published online. Data sharing statement has been updated.

Acknowledgements This study is part of the MD thesis by HS.

Contributors HS and GC contributed substantially to the conception of the work, analysis, interpretation of data for the work, drafting the work and final approval of the version to be published. IS contributed substantially to the acquisition, interpretation of data for the work, revising it critically for important intellectual content and final approval of the version to be published.

Competing interests None declared.

Provenance and peer review Not commissioned; externally peer reviewed.

Data sharing statement No Additional data are available.

Open access This is an open access article distributed in accordance with the terms of the Creative Commons Attribution (CC BY 4.0) license, which permits others to distribute, remix, adapt and build upon this work, for commercial use, provided the original work is properly cited. See: <http://creativecommons.org/licenses/by/4.0/>

© Article author(s) (or their employer(s) unless otherwise stated in the text of the article) 2017. All rights reserved. No commercial use is permitted unless otherwise expressly granted.

REFERENCES

1. American Psychiatric Association. *Diagnostic and statistical manual of mental disorders*. Arlington, VA: American Psychiatric Association, 2013.
2. Newschaffer CJ, Croen LA, Daniels J, et al. The epidemiology of autism spectrum disorders. *Annu Rev Public Health* 2007;28:235–58.
3. Senecky Y, Chodick G, Diamond G, et al. Time trends in reported autistic spectrum disorders in Israel, 1972–2004. *Isr Med Assoc J* 2009;11:30–3.
4. Torrey EF, Miller J, Rawlings R, et al. Seasonal birth patterns of neurological disorders. *Neuroepidemiology* 2000;19:177–85.
5. Mortensen PB, Pedersen CB, Westergaard T, et al. Effects of family history and place and season of birth on the risk of schizophrenia. *N Engl J Med* 1999;340:603–8.
6. Torrey EF, Miller J, Rawlings R, et al. Seasonality of births in schizophrenia and bipolar disorder: a review of the literature. *Schizophr Res* 1997;28:1–38.
7. Mick E, Biederman J, Faraone SV. Is season of birth a risk factor for attention-deficit hyperactivity disorder? *J Am Acad Child Adolesc Psychiatry* 1996;35:1470–6.
8. Liederma J, Flannery KA. Fall conception increases the risk of neurodevelopmental disorder in offspring. *J Clin Exp Neuropsychol* 1994;16:754–68.
9. Livingston R, Adam BS, Bracha HS. Season of birth and neurodevelopmental disorders: summer birth is associated with dyslexia. *J Am Acad Child Adolesc Psychiatry* 1993;32:612–6.
10. Castrogiovanni P, Iapichino S, Pacchierotti C, et al. Season of birth in psychiatry. A review. *Neuropsychobiology* 1998;37:175–81.

11. Zerbo O, Iosif AM, Delwiche L, *et al.* Month of conception and risk of autism. *Epidemiology* 2011;22:469–75.
12. Konstantareas MM, Hauser P, Lennox C, *et al.* Season of birth in infantile autism. *Child Psychiatry Hum Dev* 1986;17:53–65.
13. Tanoue Y, Oda S, Asano F, *et al.* Epidemiology of infantile autism in southern Ibaraki, Japan: differences in prevalence in birth cohorts. *J Autism Dev Disord* 1988;18:155–66.
14. Mazumdar S, Liu KY, Susser E, *et al.* The disappearing seasonality of autism conceptions in California. *PLoS One* 2012;7:e41265.
15. Lee LC, Newschaffer CJ, Lessler JT, *et al.* Variation in season of birth in singleton and multiple births concordant for autism spectrum disorders. *Paediatr Perinat Epidemiol* 2008;22:172–9.
16. Bartlik BD. Monthly variation in births of autistic children in North Carolina. *J Am Med Womens Assoc* 1981;36:363–8.
17. Mouridsen SE, Nielsen S, Rich B, *et al.* Season of birth in infantile autism and other types of childhood psychoses. *Child Psychiatry Hum Dev* 1994;25:31–43.
18. Barak Y, Ring A, Sulkes J, *et al.* Season of birth and autistic disorder in Israel. *Am J Psychiatry* 1995;152:798–800.
19. Stevens MC, Fein DH, Waterhouse LH. Season of birth effects in autism. *J Clin Exp Neuropsychol* 2000;22:399–407.
20. Gillberg C. Do children with autism have March birthdays? *Acta Psychiatr Scand* 1990;82:152–6.
21. Hebert KJ, Miller LL, Joinson CJ. Association of autistic spectrum disorder with season of birth and conception in a UK cohort. *Autism Res* 2010;3:185–90.
22. Bolton P, Pickles A, Harrington R, *et al.* Season of birth: issues, approaches and findings for autism. *J Child Psychol Psychiatry* 1992;33:509–30.
23. Landau EC, Cicchetti DV, Klin A, *et al.* Season of birth in autism: a fiction revisited. *J Autism Dev Disord* 1999;29:385–93.
24. Kolevzon A, Weiser M, Gross R, *et al.* Effects of season of birth on autism spectrum disorders: fact or fiction? *Am J Psychiatry* 2006;163:1288–90.
25. Biederman J. Attention-deficit/hyperactivity disorder: a selective overview. *Biol Psychiatry* 2005;57:1215–20.
26. Biederman J, Faraone SV. Attention-deficit hyperactivity disorder. *Lancet* 2005;366:237–48.
27. Rapley MD. Clinical practice. attention deficit-hyperactivity disorder. *N Engl J Med* 2005;352:165–73.
28. Alaerts K, Swinnen SP, Wenderoth N. Sex differences in autism: a resting-state fMRI investigation of functional brain connectivity in males and females. *Soc Cogn Affect Neurosci* 2016;11:1002–16.
29. Werling DM, Geschwind DH. Sex differences in autism spectrum disorders. *Curr Opin Neurol* 2013;26:146–53.
30. Werling DM. The role of sex-differential biology in risk for autism spectrum disorder. *Biol Sex Differ* 2016;7:58.
31. Mitra I, Tsang K, Ladd-Acosta C, *et al.* Pleiotropic Mechanisms Indicated for Sex Differences in Autism. *PLoS Genet* 2016;12:e1006425.
32. Davidovitch M, Hemo B, Manning-Courtney P, *et al.* Prevalence and incidence of autism spectrum disorder in an Israeli population. *J Autism Dev Disord* 2013;43:785–93.
33. Israel Central Bureau of Statistics. *Statistical Abstracts of Israel 2011: Census of Population and Housing Publication*. Jerusalem: Israel Bureau of Statistics, 2011.
34. Roberts EM, English PB, Grether JK, *et al.* Maternal residence near agricultural pesticide applications and autism spectrum disorders among children in the California Central Valley. *Environ Health Perspect* 2007;115:1482–9.
35. Croen LA, Grether JK, Selvin S. Descriptive epidemiology of autism in a California population: who is at risk? *J Autism Dev Disord* 2002;32:217–24.
36. Leavey A, Zwaigenbaum L, Heavner K, *et al.* Gestational age at birth and risk of autism spectrum disorders in Alberta, Canada. *J Pediatr* 2013;162:361–8.
37. Altman DG, Bland JM. Interaction revisited: the difference between two estimates. *BMJ* 2003;326:219.
38. Abramson JH. WINPEPI updated: computer programs for epidemiologists, and their teaching potential. epidemiologic perspectives & innovations [computer program]. 2011;8.
39. Deykin EY, MacMahon B. Viral exposure and autism. *Am J Epidemiol* 1979;109:628–38.
40. Lee BK, Magnusson C, Gardner RM, *et al.* Maternal hospitalization with infection during pregnancy and risk of autism spectrum disorders. *Brain Behav Immun* 2015;44:100–5.
41. Lyall K, Schmidt RJ, Hertz-Picciotto I. Maternal lifestyle and environmental risk factors for autism spectrum disorders. *Int J Epidemiol* 2014;43:443–64.
42. Grant WB, Soles CM. Epidemiologic evidence supporting the role of maternal vitamin D deficiency as a risk factor for the development of infantile autism. *Dermatoendocrinol* 2009;1:223–8.
43. Blattner RJ. The role of viruses in congenital defects. *Am J Dis Child* 1974;128:781–6.
44. Ciaranello AL, Ciaranello RD. The neurobiology of infantile autism. *Annu Rev Neurosci* 1995;18:101–28.
45. Ornoy A, Weinstein-Fudim L, Ergaz Z. Prenatal factors associated with autism spectrum disorder (ASD). *Reprod Toxicol* 2015;56:155–69.
46. Atladóttir HÓ, Henriksen TB, Schendel DE, *et al.* Autism after infection, febrile episodes, and antibiotic use during pregnancy: an exploratory study. *Pediatrics* 2012;130:e1447–e1454.
47. Muñoz-Fernández MA, Fresno M. The role of tumour necrosis factor, interleukin 6, interferon-gamma and inducible nitric oxide synthase in the development and pathology of the nervous system. *Prog Neurobiol* 1998;56:307–40.
48. Zhang X, Lv CC, Tian J, *et al.* Prenatal and perinatal risk factors for autism in China. *J Autism Dev Disord* 2010;40:1311–21.
49. Fatemi SH, Earle J, Kanodia R, *et al.* Prenatal viral infection leads to pyramidal cell atrophy and macrocephaly in adulthood: implications for genesis of autism and schizophrenia. *Cell Mol Neurobiol* 2002;22:25–33.
50. Shi L, Fatemi SH, Sidwell RW, *et al.* Maternal influenza infection causes marked behavioral and pharmacological changes in the offspring. *J Neurosci* 2003;23:297–302.
51. Zerbo O, Qian Y, Yoshida C, *et al.* Association between Influenza infection and vaccination during pregnancy and risk of Autism Spectrum Disorder. *JAMA Pediatr* 2017;171:e163609.
52. Raz R, Roberts AL, Lyall K, *et al.* Autism spectrum disorder and particulate matter air pollution before, during, and after pregnancy: a nested case-control analysis within the nurses' Health Study II Cohort. *Environ Health Perspect* 2015;123:264–70.
53. Kasahara AK, Singh RJ, Noymer A. Vitamin D (25OHD) Serum Seasonality in the United States. *PLoS One* 2013;8:e65785.
54. Kashi Z, Saeedian F, Akha O, *et al.* Vitamin D deficiency prevalence in summer compared to winter in a city with high humidity and a sultry climate. *Endokrynol Pol* 2011;62:249–51.
55. Costanzo PR, Constanzo PR, Elías NO, *et al.* (Ultraviolet radiation impact on seasonal variations of serum 25-hydroxy-vitamin D in healthy young adults in Buenos Aires). *Medicina* 2011;71:336–42.
56. Oren Y, Shapira Y, Agmon-Levin N, *et al.* Vitamin D insufficiency in a sunny environment: a demographic and seasonal analysis. *Isr Med Assoc J* 2010;12:751–6.
57. Marvin K. Season of Birth Effects in Autism Spectrum Disorders. *Honors Scholar Theses* 2008.
58. Hertz-Picciotto I, Delwiche L. The rise in autism and the role of age at diagnosis. *Epidemiology* 2009;20:84–90.

APPENDIX

Table A Crude and adjusted odd ratio (OR) and 95% confidence interval from multivariable logistic regression of autistic spectrum disorder (ASD) for first month of second trimester, by socioeconomic status (SES)

Month of second trimester	OR crude	95% CI		OR adjusted*	95% CI		OR crude	95% CI		OR adjusted*	95% CI	
January	1 (ref.)			1 (ref.)			1 (ref.)			1 (ref.)		
February	0.83	0.51	1.35	0.83	0.51	1.36	1.03	0.67	1.57	1.03	0.67	1.57
March	1.08	0.69	1.69	1.08	0.69	1.69	0.87	0.57	1.35	0.86	0.56	1.33
April	0.89	0.55	1.43	0.89	0.55	1.43	1.12	0.74	1.69	1.11	0.73	1.67
May	0.68	0.41	1.12	0.68	0.41	1.13	0.99	0.65	1.51	0.98	0.64	1.49
June	0.87	0.53	1.41	0.87	0.54	1.41	1.09	0.72	1.64	1.07	0.71	1.63
July	1.00	0.62	1.59	0.98	0.61	1.56	1.02	0.67	1.56	1.04	0.68	1.59
August	0.74	0.44	1.23	0.72	0.43	1.20	0.97	0.63	1.49	0.97	0.63	1.49
September	0.94	0.58	1.54	0.93	0.57	1.51	0.86	0.55	1.35	0.87	0.55	1.36
October	0.84	0.51	1.37	0.83	0.51	1.37	1.10	0.73	1.66	1.11	0.73	1.67
November	0.84	0.51	1.40	0.83	0.50	1.38	1.25	0.84	1.87	1.24	0.83	1.86
December	0.83	0.51	1.36	0.83	0.51	1.36	1.10	0.73	1.65	1.09	0.72	1.64

*Adjusted for child's age, father's age, mother's age, district, birth weight, age with imputation (see text for details).

Table B Crude and adjusted odd ratio (OR) and 95% confidence interval from multivariable logistic regression of autistic spectrum disorder (ASD) for first month of third trimester, by socioeconomic status (SES)

Month of second trimester	OR crude	95% CI		OR adjusted*	95% CI		OR crude	95% CI		OR adjusted*	95% CI	
January	1 (ref.)			1 (ref.)			1 (ref.)			1 (ref.)		
February	1.04	0.61	1.77	1.04	0.61	1.78	1.22	0.81	1.82	1.21	0.81	1.81
March	0.97	0.57	1.63	0.97	0.58	1.64	0.96	0.63	1.46	0.95	0.63	1.45
April	1.19	0.73	1.95	1.20	0.74	1.97	1.13	0.75	1.68	1.12	0.75	1.68
May	0.86	0.51	1.46	0.87	0.52	1.48	0.86	0.56	1.32	0.86	0.56	1.31
June	1.26	0.77	2.04	1.27	0.78	2.06	0.84	0.55	1.30	0.83	0.54	1.27
July	0.99	0.60	1.65	1.01	0.61	1.67	1.07	0.71	1.61	1.05	0.70	1.58
August	0.78	0.45	1.33	0.79	0.46	1.36	0.99	0.66	1.51	0.98	0.64	1.48
September	1.12	0.68	1.86	1.12	0.68	1.86	1.12	0.75	1.69	1.12	0.74	1.69
October	0.84	0.49	1.44	0.83	0.48	1.42	0.83	0.53	1.29	0.83	0.54	1.30
November	0.92	0.53	1.57	0.91	0.53	1.56	1.00	0.65	1.53	1.00	0.65	1.54
December	1.07	0.64	1.80	1.08	0.64	1.81	0.81	0.52	1.26	0.81	0.52	1.26

*Adjusted for child's age, father's age, mother's age, district, birth weight, age with imputation (see text for details).

Table C Crude and adjusted odd ratio (OR) and 95% confidence interval from multivariable logistic regression of autistic spectrum disorder (ASD) for season of birth, by socioeconomic status (SES)

Season of birth	SES median-high						SES median-low					
	OR crude		95% CI		OR adjusted		95% CI		OR crude		95% CI	
	OR crude	95% CI	OR adjusted	95% CI	OR crude	95% CI	OR adjusted	95% CI	OR crude	95% CI	OR adjusted	95% CI
Winter	1 (ref.)		1 (ref.)		1 (ref.)		1 (ref.)		1 (ref.)		1 (ref.)	
Spring	1.12	0.84	1.50	1.12	0.84	1.50	1.06	0.83	1.34	1.06	0.83	1.34
Summer	1.24	0.94	1.63	1.23	0.93	1.63	1.08	0.85	1.37	1.08	0.85	1.37
Fall	1.14	0.86	1.51	1.14	0.86	1.51	1.05	0.82	1.33	1.04	0.82	1.32

*Adjusted for child's age, father's age, mother's age, district, birth weight, age with imputation.

Table D Crude and adjusted odd ratio (OR) and 95% confidence interval from multivariable logistic regression of autistic spectrum disorder (ASD) for season of conception, by socioeconomic status (SES)

Season of conception	SES median-high						SES median-low					
	OR crude		95% CI		OR adjusted*		95% CI		OR crude		95% CI	
	OR crude	95% CI	OR adjusted*	95% CI	OR crude	95% CI	OR adjusted*	95% CI				
Winter	1 (ref.)		1 (ref.)		1 (ref.)		1 (ref.)					
Spring	1.00	0.75 1.34	0.99	0.74 1.33	0.96	0.75 1.22	0.96	0.75 1.23				
Summer	1.00	0.74 1.35	0.99	0.73 1.33	0.96	0.75 1.22	0.97	0.76 1.24				
Fall	1.09	0.82 1.45	1.08	0.82 1.44	1.05	0.83 1.33	1.06	0.83 1.34				

*Adjusted for child's age, father's age, mother's age, district, birth weight, age with imputation.

Table E Crude and adjusted odd ratio (OR) and 95% confidence interval from multivariable logistic regression of autistic spectrum disorder (ASD) for month of conception, by socioeconomic status (SES)

	SES 1-6						SES 7-10					
Month of conception	OR crude	95% CI		OR adjusted*	95% CI		OR crude	95% CI		OR adjusted*	95% CI	
January	1 (ref.)			1 (ref.)			1 (ref.)			1 (ref.)		
February	0.89	0.52	1.50	0.89	0.52	1.51	1.09	0.72	1.65	1.08	0.72	1.63
March	1.13	0.70	1.85	1.14	0.70	1.86	0.84	0.55	1.30	0.84	0.54	1.29
April	0.90	0.53	1.53	0.89	0.53	1.52	1.06	0.71	1.60	1.09	0.72	1.64
May	1.05	0.63	1.75	1.03	0.62	1.71	0.94	0.62	1.44	0.95	0.62	1.45
June	1.19	0.72	1.96	1.17	0.71	1.93	0.74	0.46	1.16	0.75	0.47	1.18
July	0.79	0.45	1.38	0.79	0.45	1.38	0.96	0.63	1.45	0.97	0.64	1.47
August	1.11	0.67	1.83	1.09	0.66	1.81	1.13	0.76	1.68	1.13	0.76	1.68
September	1.29	0.79	2.10	1.28	0.78	2.08	1.09	0.73	1.64	1.10	0.73	1.65
October	0.95	0.57	1.58	0.96	0.58	1.59	0.88	0.58	1.34	0.89	0.58	1.35
November	1.14	0.70	1.86	1.14	0.70	1.86	1.16	0.78	1.72	1.16	0.78	1.72
December	1.18	0.74	1.91	1.18	0.74	1.91	0.90	0.59	1.36	0.89	0.59	1.36

*Adjusted for child's age, father's age, mother's age, district, birth weight, age with imputation. see text for details