

BMJ Open Does the impact of a plant-based diet during pregnancy on birth weight differ by ethnicity? A dietary pattern analysis from a prospective Canadian birth cohort alliance

Michael A Zulyniak,^{1,2} Russell J de Souza,³ Mateen Shaikh,³ Dipika Desai,⁴ Diana L Lefebvre,¹ Milan Gupta,^{1,5} Julie Wilson,⁶ Gita Wahi,^{3,7} Padmaja Subbarao,⁸ Allan B Becker,⁹ Piush Mandhane,¹⁰ Stuart E Turvey,¹¹ Joseph Beyene,³ Stephanie Atkinson,⁷ Katherine M Morrison,⁷ Sarah McDonald,³ Koon K Teo,^{1,4} Malcolm R Sears,¹ Sonia S Anand,^{1,3,4} for the NutriGen Alliance investigators

To cite: Zulyniak MA, de Souza RJ, Shaikh M, *et al.* Does the impact of a plant-based diet during pregnancy on birth weight differ by ethnicity? A dietary pattern analysis from a prospective Canadian birth cohort alliance. *BMJ Open* 2017;7:e017753. doi:10.1136/bmjopen-2017-017753

► Prepublication history and additional material for this paper are available online. To view these files, please visit the journal online (<http://dx.doi.org/10.1136/bmjopen-2017-017753>).

Received 19 May 2017
Revised 29 September 2017
Accepted 2 October 2017



CrossMark

For numbered affiliations see end of article.

Correspondence to
Professor Sonia S Anand;
anands@mcmaster.ca

ABSTRACT

Objective Birth weight is an indicator of newborn health and a strong predictor of health outcomes in later life. Significant variation in diet during pregnancy between ethnic groups in high-income countries provides an ideal opportunity to investigate the influence of maternal diet on birth weight.

Setting Four multiethnic birth cohorts based in Canada (the NutriGen Alliance).

Participants 3997 full-term mother–infant pairs of diverse ethnic groups who had principal component analysis-derived diet pattern scores—plant-based, Western and health-conscious—and birth weight data.

Results No associations were identified between the Western and health-conscious diet patterns and birth weight; however, the plant-based dietary pattern was inversely associated with birth weight ($\beta=-67.6$ g per 1-unit increase; $P<0.001$), and an interaction with non-white ethnicity and birth weight was observed. Ethnically stratified analyses demonstrated that among white Europeans, maternal consumption of a plant-based diet associated with lower birth weight ($\beta=-65.9$ g per 1-unit increase; $P<0.001$), increased risk of small-for-gestational age (SGA; OR=1.46; 95% CI 1.08 to 1.54; $P=0.005$) and reduced risk of large-for-gestational age (LGA; OR=0.71; 95% CI 0.53 to 0.95; $P=0.02$). Among South Asians, maternal consumption of a plant-based diet associated with a higher birth weight ($\beta=+40.5$ g per 1-unit increase; $P=0.01$), partially explained by cooked vegetable consumption.

Conclusions Maternal consumption of a plant-based diet during pregnancy is associated with birth weight. Among white Europeans, a plant-based diet is associated with lower birth weight, reduced odds of an infant born LGA and increased odds of SGA, whereas among South Asians living in Canada, a plant-based diet is associated with increased birth weight.

INTRODUCTION

Birth weight is an indicator of infant health and a strong predictor of future health

Strengths and limitations of this study

- Prospective data collection, large sample size, adequate power to allow for ethnic comparisons and detailed measurement of diet using ethnic-specific Food Frequency Questionnaire are the strengths of the study.
- Limitations include self-reported retrospective food intake, the inability to adjust for differences in cooking methods and residual confounding, which is a potential bias in all cohort studies.
- To minimise potential bias, a sensitivity analysis was conducted, which excluded mothers who may have deliberately altered their diet due to a medical condition.

outcomes.¹ Infants born small (birth weight <10th percentile) or large (birth weight \geq 90th percentile) for sex and gestational age are at increased risk of future health complications, including asthma,² obesity³ and cardiovascular disease.⁴ High-income countries generally have similar proportions of babies who weigh <2500 g at birth, but there is greater variation (up to 10%) in the proportion of infants born >4000 g.^{5–7} However, such population figures often mask important differences in the distribution of birth weight between ethnic groups. In the USA, white Europeans have more high birth weight (9.6%) than low birth weight infants (7.0%); African-Americans have more low birth weight (13.1%) than high birth weight infants (4.3%); and Hispanics have an equal proportion (7.1% high birth weight and 7.1% low birth weight infants).⁵ Furthermore, we recently reported that newborns in Canada

of South Asian ancestry (those who originate from the Indian subcontinent) are of lower birth weight, and that Aboriginal newborns are of higher birth weight compared with full-term newborns of white European ancestry in Canada.⁸ This provides additional evidence that among full-term newborns, ethnic differences in birth weight distribution exist. Numerous factors influence birth weight, including gestational weight gain, maternal prepregnancy weight, maternal height, gestational diabetes and fetal smoke exposure.^{9 10} It has been postulated that maternal dietary intake also influences birth weight.^{11 12} The investigation of specific food items and nutrients has advanced our understanding of specific nutrient deficiency syndromes (eg, neural tube defects) and facilitated the identification of particularly harmful food components (eg, trans fats). However, single-nutrient studies may be misleading because they fail to capture the complex interplay between foods and nutrients consumed as meals over long periods of time. To overcome this, the empirical derivation of dietary patterns¹³ has been proposed as a method to characterise diet that more accurately reflects how we consume foods and nutrients, and these patterns can be assessed for their associations with health and disease.

In this study, we investigated the association between maternal diet and birth weight in a multiethnic cohort using the dietary pattern analysis approach.

METHODS

Study population

The NutriGen Alliance¹³ is a consortium of four birth cohorts in Canada investigating the contribution of nutritional, genetic and epigenetic factors to the health of pregnant women and their children—(1) Canadian Healthy Infant Longitudinal Development¹⁴ study (CHILD); (2) Family Atherosclerosis Monitoring In earLY life (FAMILY) study¹⁵; (3) SouTh Asian birth cohORT (START)¹⁶; and (4) Aboriginal Birth Cohort (ABC).¹⁷ Of these, ABC is still recruiting. To date (October 2016), 5018 pregnant women have provided comprehensive clinical and dietary data and 4556 (≈90.8%) have provided birth data. From this group, 559 women were excluded from the present analysis because they (1) reported an implausible energy intake (<500 or ≥6500 kcal/day), (2) were non-singleton pregnancies, (3) delivered preterm (<36 weeks) and/or (4) reported not knowing their ethnic origin. Preterm infants were excluded to avoid confounding by pregnancy or neonatal complications (eg, trauma, stress, infections or placenta previa). The remaining 3997 full-term infant–mothers represented multiple maternal self-reported ethnicities—white European (n=2367), South Asian (n=884), East/South-East Asian (n=335), Aboriginal (n=190), African (n=60) and women who reported another ethnicity (n=141, eg, Egyptian, Haitian and others) (online supplementary efigure 1). Informed written consent was obtained from each participant within each study.

Dietary assessment

Dietary information during pregnancy was collected in each cohort 24–28 weeks' gestation using a validated semiquantitative Food Frequency Questionnaire (FFQ). The CHILD cohort used a version of the Fred Hutchinson Cancer Center tool.¹⁸ The FAMILY, START and ABC cohorts used ethnic-specific FFQs developed for the Study of Health Assessment and Risk in Ethnic groups.^{13 16 17}

Diet pattern analysis

Detailed methods of FFQ harmonisation and the statistical derivation of dietary patterns within the NutriGen cohort have been previously described.¹³ We performed principal component analysis (PCA) to identify three orthogonal dietary patterns—'plant-based', 'western' and 'health-conscious' (online supplementary efigure 1). This approach retains the information of the quantitative data from the original FFQ but shifts the focus from individual food components to food combinations. An empirically derived measure of adherence (ie, loading scores) for each person to each of the individual dietary patterns is calculated in the PCA. The PCA loading scores for each individual were adjusted to the mean total population caloric intake (2500 kcal/day) using the residual method.¹⁹

Clinical parameters

The primary outcome of this analysis was newborn birth weight. Birth weight was collected from the birth chart or measured in duplicate by trained staff perinatally (median <24 hours; 95% ≤48 hours). Due to the ethnic diversity of the study population, sex-specific and ethnic-specific birth weight cut points were used to define infants born small (SGA <10th percentile) or large (LGA ≥90th percentile) for gestational age, which demonstrate greater accuracy to identify risk in multiethnic cohorts.²⁰ Gestational age was determined by ultrasound at the initial visit. Data on all clinical parameters are provided in table 1.

Statistical analysis

The distribution of exposures and covariates was summarised using JMP (V.9.0.1) as means (SD) for continuous variables, or counts (%) for categorical variables; between groups, differences were assessed using analysis of variance (continuous variables) or χ^2 test (categorical variables).

Prepregnancy weights from the CHILD cohort were available for ≈73% of women. To impute missing weights, we constructed a multivariable model with white Europeans recruited in CHILD and FAMILY. The correlation between the value predicted by this equation and observed prepregnancy weight was 0.42.

Due to the numerous biological, sociological and environmental variables that associate with birth weight, we aimed to identify a parsimonious set of covariates. All variables listed in table 1 significant by association with birth weight as the response variable in a simple linear regression ($\alpha \leq 0.10$) were entered into a forward stepwise selection procedure. To account for perinatal reductions in weight,²¹ timing of birth weight measurement was

Table 1 Participant characteristics

Variable	Overall	East/South-East				P value*
		White European	South Asian	Asian	Aboriginal	
n	3997	2367	884	335	190	60
Maternal						
Age at enrolment (years)	31.6 (4.7)	32.1 (4.6)	30.2 (4.0)	33.1 (4.2)	28.0 (6.0)	31.2 (5.4)
Height (cm)	164.4 (6.9)	166.0 (6.5)	162.2 (6.4)	160.2 (6.6)	165.3 (6.9)	163.7 (7.7)
Prepregnancy weight (kg)	66.8 (13.9)	69.1 (14.1)	62.4 (11.9)	58.7 (9.2)	75.4 (16.9)	69.7 (14.2)
Prepregnancy body mass index (kg/m ²)	24.7 (4.8)	25.1 (5.0)	23.7 (4.4)	23.0 (3.3)	27.6 (6.2)	26 (4.8)
Final pregnancy weight (kg)	81.4 (15.7)	84.2 (16.2)	76.5 (12.7)	72.4 (12.5)	91.3 (22.7)	83.1 (14.8)
Primiparous (%)	49.8	51.3	41.8	59.7	40.8	45.0
Gestational diabetes (%)	11.3	7.4	24.2	10.8	4.2	5.0
Years lived in Canada	23.4 (12.2)	29.2 (8.3)	8.4 (7.8)	20.4 (12.5)	28.5 (5.8)	19.4 (12.6)
Postsecondary education (%)	85.1	86.8	84.7	95.2	52.9	85.0
Smoking during pregnancy (%)						
Never smoked	77.1	70.4	99.6	87.7	40.6	88.3
Quit before pregnancy	16.2	21.8	0.2	9.9	25.7	5.0
Quit during pregnancy	3.6	4.3	0.2	1.8	15.5	1.7
Currently smoking	3.1	3.5	0.0	0.6	18.2	5.0
Household income (%)						
Total income ≥\$60K	77.2	88.9	43.5	87.2	48.1	68.3
Multivitamin use (%)	85.8	89.3	71.8	95.9	91.4	83.4
Vegetarianism (%)	12.3	3.5	37.1	0.0	0.0	0.0
Newborn						
Sex (% male)	51.8	52.3	49.1	52.2	46.8	53.2
Gestational age (weeks)	39.6 (1.1)	39.7 (1.1) ^a	39.4 (1.1) ^b	39.4 (1.1) ^{bc}	39.6 (1.1) ^{bc}	39.5 (1.1)
Birth weight (g)	3432 (466)	3493 (457) ^a	3265 (438) ^b	3330 (448) ^c	3567 (469) ^d	3347 (468)
Birth length (cm)	51.3 (2.5)	51.4 (2.5) ^{ab}	51.3 (2.5) ^{ab}	51.0 (2.4) ^b	51.6 (3.2) ^a	51.4 (2.4)
LGA (n)	370 (9.8%)	224 (9.9%)	87 (9.9%)	30 (9.2%)	18 (9.6%)	10 (17%)
SGA (n)	348 (9.2%)	219 (9.6%)	83 (9.4%)	29 (8.9%)	14 (7.5%)	1 (1.7%)

*Continuous measures (mean and SD) were compared using analysis of variance, and categorical variables were compared using χ^2 . Non-matching superscript letters indicate significant difference ($P < 0.05$) between ethnic groups. Data on parity, prepregnancy weight, smoking history, ethnicity, postgraduate education, marital status, employment status and total household income were self-reported. Maternal age, height and gestational age of offspring at birth were obtained from participant medical records. Last measured maternal pregnancy weight was obtained from medical records at time of birth (ABC, FAMILY and START) or using a combination of medical records or maternal recollection (CHILD). Gestational diabetes was determined either through (1) self-reported by the mother or recorded from the medical chart (CHILD), or (2) diagnosed using an oral glucose tolerance test, self-reported by the mother or recorded from the medical chart (ABC, FAMILY and START).
 ABC, Aboriginal Birth Cohort; CHILD, Canadian Healthy Infant Longitudinal Development study; FAMILY, Family Atherosclerosis Monitoring in early life; LGA, ethnic-specific large-for-gestational age (birth weight ≥90th percentile for gestational age and sex); SGA, ethnic-specific small-for-gestational age (birth weight <10th percentile for gestational age and sex); START, South Asian birth cohort.

assessed as a covariate. Covariates significant at $\alpha < 0.05$ in the stepwise multivariable model were retained as covariates in the final model.

Multivariable linear regression was used to assess the association between maternal dietary patterns and covariates on infant birth weight. The main effects, ethnicity and each of the dietary patterns were included in the initial model. Interaction terms between a dietary pattern and ethnicity were added if they were both significant at $P < 0.05$. To determine which foods contributed to the observed diet and birth weight association, food groups were added one at a time to the full dietary pattern–birth weight multivariable model. If the association between the dietary pattern and birth weight was rendered non-significant by the addition of the food group, we deemed it to be an important explanatory variable.

For ethnic-specific analyses, we had adequate power to investigate the association between plant-based diet and birth weight within the two largest ethnic groups—white Europeans (>95% power) and South Asians ($\approx 80\%$ power)—but <40% power to do so in the other ethnic groups. Therefore, stratified analyses by ethnic group were limited to white Europeans and South Asians.

RESULTS

Demographic and clinical parameters

Maternal demographic and clinical parameters for each of the six ethnic groups are presented in [table 1](#).

Birth weight

There were significant differences in birth weight by ethnic group ($P < 0.001$). Aboriginal newborns had the highest birth weight, followed by white Europeans, other ethnicity, African origin, East/South-East Asians and South Asians ([table 1](#)).

Effect of dietary patterns on birth weight

In the fully adjusted main-effects multivariable model ([table 2](#), model 1), non-white ethnicity and plant-based diet were significantly associated, while the Western and health-conscious diet patterns were not. After removal of non-significant dietary patterns, an interaction between the plant-based diet and non-white ethnicity was tested and found to be significant ([table 2](#), model 2).

Stratified analyses

Multivariable models, stratified by ethnicity (white European and South Asian) and adjusted for the same covariates as the main model (except for smoking in South Asians, as its prevalence was <0.5%), were built, treating dietary pattern scores as either continuous ([table 3](#)) or dichotomous (fourth vs first quartile) variables ([table 3](#)). In white Europeans, an inverse association was observed between plant-based diet and birth weight. Among South Asians, a positive association was observed between the plant-based diet and birth weight ([figure 1](#)).

A sensitivity analysis, which excluded mothers who were diagnosed with gestational diabetes and/or hypertension, resulted in dietary associations that were largely unchanged,

Table 2 Final multivariable regression of the association and interaction between diet patterns in white and non-whites with birth weight (g)

Variables	Model 1		Model 2	
	β	P value	β	P value
Intercept	-4551.9	<0.001	-4560.8	<0.0001
Prepregnancy weight (kg)	5.5	<0.001	5.4	<0.001
Maternal height (cm)	8.4	<0.001	8.5	<0.001
Parity (number of children)	67.6	<0.001	66.0	<0.001
Gestational age (weeks)	158.4	<0.001	158.2	<0.001
Smoking status	-21.9	0.04	-23.6	0.02
Infant sex (female=1)	-118.6	<0.001	-118.8	<0.001
Non-white ethnicity	-33.5	0.04	-28.1	0.08
Plant-based diet	-34.6	<0.001	-67.6	<0.001
Western diet	-12.7	0.35	ns	ns
Health-conscious diet	6.0	0.56	ns	ns
Plant-based diet \times non-white	NA	NA	40.5	0.03

Model 1 included all covariates, non-White ethnicity indicator, three diet patterns identified using PCA: plant-based, Western and health-conscious diet. Model 2 tests interactions between non-white ethnicity and significant diet patterns after removal of non-significant variables from model 1 ($n=3997$). Non-significant variables removed from the model are denoted as 'ns'. Smoking status was ordinal and input as either 0=never smoked, 1=quit smoking pre-pregnancy, 2=quit smoking during pregnancy or 3=current smoking. Parity was ordinal and reported as having 0, 1, 2, 3, 4, 5 or 6 (or more) children. Non-whites included individuals who self-reported as being of South Asian, East/South-East Asian or Aboriginal descent.

NA, not applicable; PCA, principal component analysis.

Table 3 Multivariable regression of plant-based diet pattern (as a continuous measure) and birth weight stratified by ethnicity: white European and South Asians

Variables	White Europeans		South Asians	
	β	P value	β	P value
Intercept	-4723.7	<0.001	-4248.1	<0.001
Prepregnancy weight (kg)	5.1	<0.001	5.3	<0.001
Maternal height (cm)	9.3	<0.001	6.0	0.01
Parity (number of children)	73.8	<0.001	43.5	0.01
Gestational age (weeks)	159.3	<0.001	155.6	<0.001
Smoking status	-37.6	0.002	ns	ns
Infant sex (female=1)	-127.7	<0.001	-98.3	<0.001
Plant-based (continuous)	-65.9	<0.001	40.5	0.01

White European $r^2=0.246$, South Asian $r^2=0.178$. The same covariates were included for each ethnic-specific model, but non-significant covariates (eg, 'smoking' in South Asians) were removed in the final model (denoted as 'ns'). Smoking status was coded as follows: 0=never smoked, 1=quit smoking prepregnancy, 2=quit smoking during pregnancy or 3=currently smoking. Parity was reported as having 0, 1, 2, 3, 4, 5 or 6 (or more) children. White Europeans (n=2367) and South Asians (n=884).

with a negative association remaining in white Europeans ($\beta=-64.5$ g per 1-unit increase in score; n=2038; P=0.002) and a positive association in South Asians ($\beta=38.3$ g per 1-unit increase in score; n=728; P=0.03).

In white Europeans, a 1-unit increase in plant-based diet score was associated with $\approx 50\%$ increase in odds of SGA and a $\approx 30\%$ decrease in odds of LGA (table 4). In South Asians, a 1-unit increase in plant-based diet score presented a non-significant reduction in odds of SGA (P=0.428) and non-significant increase in odds of LGA (P=0.249) (table 4).




Figure 1 Multivariable regression of plant-based diet pattern and birth weight stratified by ethnicity. Association between maternal adherence to a plant-based diet, whereby a higher score reflects greater adherence to the dietary pattern, and birth weight in white Europeans (dashed line; n=2367) and South Asians (solid line; n=884) after accounting for covariates: prepregnancy weight, maternal height, maternal smoking status, number of children, and infant gestational age and sex. Dotted line=95% CI.

Validity of the plant-based diet score

The classification of individuals as plant-based consumers using the harmonised FFQ/PCA was validated against a classification using preharmonisation FFQ data. Using the raw number of servings obtained by each study-specific FFQ, the sum of servings for five plant-based foods groups common across all FFQs—that is, total dairy (an important protein source for lacto-vegetarians), vegetables, legumes, fruit, and breads and rice—was calculated. There was a high degree of agreement between preharmonised and postharmonised scores, with $>90\%$ of the same plant-based consumers being classified in the fourth quartile using either metric (P<0.001).

Diet comparison

The food groups among individuals in the fourth quartile of the plant-based diet were compared between white Europeans and South Asians to understand the ethnic-specific structure of plant-based diets. For white Europeans, the plant-based diet was characterised by high intakes of fruit, nuts and seeds, convenience foods, sweet drinks, sweets and eggs, and moderate intakes of fish, poultry and red meat, while the South Asian plant-based diet was characterised by higher intakes of South Asian breads (eg, roti, naan), rice, legumes, raw vegetables and cooked vegetables (online supplementary table 2). The use of multivitamins was comparable among high consumers of the plant-based diet in white Europeans (79%) and South Asians (78%), but differed in the low consumers of the plant-based diet, with white Europeans (94%) more likely to take multivitamins than South Asians (66%).

Among white Europeans, higher levels of postsecondary education, employment and marriage were observed in the fourth versus the first quartile of the plant-based diet, while South Asians in the fourth quartile were more likely to have recently immigrated to Canada, live with extended family in their household and were less likely to

Table 4 Multivariable regression of plant-based diet pattern (as a categorical measure) and birth weight stratified by ethnicity: white European and South Asians

Variables	White Europeans		South Asians	
	β	P value	β	P value
Intercept	-5001.5	<0.001	-2482.3	<0.001
Prepregnancy weight (kg)	4.1	<0.001	6.8	<0.001
Maternal height (cm)	9.4	<0.001	ns	ns
Parity (number of children)	77.1	<0.001	ns	ns
Gestational age (weeks)	168.6	<0.001	135.6	<0.001
Smoking status	ns	ns	ns	ns
Infant sex (female=1)	-130.4	<0.001	-122.8	0.002
Plant-based (fourth vs first quartile)	-81.2	0.002	70.5	0.07

White European $r^2=0.246$, South Asian $r^2=0.155$. The same covariates were included for each ethnic-specific model, but non-significant covariates (eg, 'smoking' in South Asians) were removed in the final model (denoted as 'ns'). Smoking status was coded as follows: 0=never smoked, 1=quit smoking pre-pregnancy, 2=quit smoking during pregnancy or 3=currently smoking. Parity was reported as having 0, 1, 2, 3, 4, 5 or 6 (or more) children. White Europeans (n=2367) and South Asians (n=884).

have postsecondary education compared with quartile 1 (online supplementary table 2).

Food groups

Post-hoc analyses were conducted separately in white Europeans and South Asians, and identified 15 mutually exclusive food groups (online supplementary table 2) that showed a high degree of variance between the individuals in the first and fourth quartiles of the plant-based diet. Only the addition of 'cooked vegetables' to the model reduced the magnitude of the plant-based dietary pattern association in South Asians ($\beta=37.9$ g; $P=0.041$) (online supplementary table 3). No other foods influenced the association of plant-based diet and birth weight in either ethnic group. Multivitamin use had no influence on the association.

Effect of total energy intake on birth weight

To investigate the effect of maternal energy intake on birth weight, we evaluated the association between total energy intake and birth weight in our population of white Europeans and South Asians with and without an interaction term for non-white ethnicity. Without ethnicity or its interaction term, we note there is a significant association between total maternal energy intake and infant birth weight, whereby a 100-calorie increase in total energy associated with 4 g (2–7 g) increase in birth weight ($P<0.01$). When ethnicity and its interaction term (energy intake \times non-white ethnicity) were included in the model, total energy intake and the interaction term were no longer significant ($P>0.05$), whereas ethnicity remained so ($P<0.01$), indicating that a South Asian newborn was estimated to weigh 231 g less (-266 to -197 g) than a white European newborn.

DISCUSSION

Among women living in a high-income country, ethnicity was a predictor of birth weight, independent of caloric

intake, in white Europeans and South Asians. Foods and diet differ substantially between ethnic groups and may contribute to the association. A plant-based diet in pregnancy is associated with newborn birth weight, the effects of which are moderated by ethnicity. In white European women, consumption of a plant-based diet is associated with lower birth weight, an increased risk of an SGA infant and a reduced risk of an LGA infant, while in South Asian women the plant-based diet is associated with higher birth weight but had no significant effect on the risk of having an SGA or LGA infant.

Our findings are consistent with a recent systematic review that failed to find a consistent association of vegetarian diets on birth weight; five studies reported vegetarian mothers had lower birth weight babies, and two studies reported higher birth weights.²² The interpretation of the studies was complicated by (1) the numerous ethnic groups investigated, (2) differences in the foods that characterised the vegetarian and non-vegetarian (ie, control) groups within each of the reported studies, and (3) variability in the socioeconomic status (SES) of individuals on a vegetarian diet. Similar, in our study, birth weight was not associated with self-reported vegetarianism, which may be partly explained by the high degree of variability in the foods that comprise a plant-based diet across ethnic groups. This suggests that 'vegetarianism' may be too crude a descriptor of diet to allow for discovery of true associations with birth weight, because the definition of this term is qualitatively different across ethnic groups.

Studies investigating the effect of a plant-based diet in ethnically homogeneous populations are sparse. No study in pregnant South Asian women living in a high-income country has reported on dietary patterns and birth weight. However, our findings in white Europeans are consistent with previous work indicating that smoking is associated with reduced birth weight,²³ an increased risk of SGA²⁴ and reduced risk of LGA,²⁵ and that increased parity, maternal

weight and height are associated with a reduced risk of SGA²⁶ and increased risk of LGA.²⁵ Concerning diet, a previous study in white Europeans (n=38)²⁷ reported that infants born to mothers whose diets were classified as 'plant-based' were lighter (3310g; 95% CI 3080 to 3350g) than infants born to 'omnivore' mothers (3480g; 95% CI 3350 to 3620g). The Danish National Birth Cohort (n=44612) reported that mothers adhering to a plant-based dietary pattern were at reduced risk of having an SGA infant (OR=0.74; 95% CI 0.64 to 0.86) compared with mothers adhering to a Western diet pattern.²⁸ However, this observation may be confounded by the high prevalence of smoking during pregnancy in the Western diet (38%) versus plant-based diet group (14%).

Our investigation to determine which of the individual food groups characteristic of the plant-based diet may be driving the association revealed that controlling for cooked vegetables reduced the magnitude and significance of the association between the plant-based diet and birth weight in South Asians but not among white Europeans. In South Asians, cooked vegetables were independently associated with birth weight (after removing plant-based dietary pattern from the model), whereas multivitamin use was not. This suggests that consumption of 'cooked vegetables' may be a driver of the observed association between the plant-based diet and increased birth weight in South Asians.

Prior cohort studies from India have shown that lower levels of serum vitamin B₁₂²⁹ and folate in pregnancy are associated with low birth weight.³⁰ In a randomised controlled trial, prenatal supplementation with vitamins A, D, E, C and B, as well as iron, zinc, copper, selenium and iodine, reduced the risk of low birth weight (Risk Ratio (RR) =0.88; 95% CI 0.85 to 0.91).³¹ In the slum-dwelling population of Mumbai (India), micronutrient-rich snacks (green leafy vegetables, fruit and milk) consumed before or during pregnancy increase infant birth weight (+48g; 95% CI 1 to 96g) compared with a low-micronutrient snack (potato and onion).³² However, the dietary context in Canada differs substantially from that of low-income countries. Since 1998 flour and cereals have been fortified with folate.³³ This, coupled with relatively high rates of multivitamin supplementation in pregnancy, makes folate and vitamin B₁₂ deficiency in pregnancy unlikely among Canadians. In addition, women who avoid meat products yet consume milk, yoghurt, cheese and eggs usually attain adequate vitamin B₁₂. Thus, even in the context of virtually no dietary folate and vitamin B₁₂ deficiency, high consumption of cooked vegetables as part of a traditional Indian diet is associated with higher birth weight compared with a lower plant-based/higher animal protein diet.

We can only speculate about the mechanisms underlying the association with cooked vegetables as we did not collect detailed information on cooking methods. In our study, South Asian women reported consuming a higher percentage of energy from fat than did white Europeans. Differences in food preparation methods can significantly alter the chemical and nutritional

composition of dishes.³⁴ This was recently examined using spectrophotometry to determine the chemical composition of fresh and fried portions of ovo-vegetarian dishes.³⁵ Fried ovo-vegetarian dishes prepared in oil were lower in protein, fibre and carbohydrate content but higher in caloric and lipid content per 100 g serving, compared with fresh dishes.³⁵ This aligns with work in pregnant South Asian women residing in East London (UK), who had a higher proportion of energy from fat (40% energy) compared with other ethnic groups (white European, African and African-Caribbean).³⁶ Further, a prospective birth study of 797 rural Indian women, reported that, although protein and carbohydrate intake at 18 weeks of gestation was unrelated to birth measurements, fat intake was associated with newborn length and adiposity.³⁷ This suggests that South Asian cooking methods may alter the macronutrient composition of food by increasing the proportion of total fat relative to other nutrients, and offers a possible explanation regarding the difference in the association between the plant-based diet and birth weight in South Asians and white Europeans.

Although markers of SES were not significantly associated with birth weight, understanding differences in sociodemographic factors between high and low consumers of the plant-based diet is informative to understand the context in which these diets are consumed. In the Avon Longitudinal Study of Parents and Children (ALSPAC) birth cohort (n=11 833), SES was not associated with birth weight but was associated with food choices, whereby women of high SES consumed more healthy foods and less processed foods than women of a lower SES.²³ In the present study, white European women who adhered most strongly to the plant-based diet represented the highest socioeconomic group compared with less adherent white European women. Interestingly this group consumed multivitamin supplements less frequently, and it is possible they perceived their 'healthy diet consumption' negated the need for the recommended daily supplement in pregnancy. In contrast, South Asian women who adhered most strongly to the plant-based diet were more recent immigrants to Canada, had lower income and were living with more extended family members, indicating more traditional cultural practices compared with the South Asian women who were low adherers to the plant-based diet. More recent immigrants with a high adherence to a plant-based diet also had greater multivitamin use compared with South Asians who were longer settled in Canada. The relationship between immigration and diet is complex due to the numerous and varied factors that influence acculturation within each ethnic group and country.³⁸⁻⁴⁰ Thus, policy recommendations for dietary guidelines should consider the socioeconomic profile of the population. Although recent evidence suggests that plant-based diets may improve health and reduce risk of disease,^{41 42} our observations suggest that the food composition of the plant-based diets matters. Emphasis should likely be placed on whole foods, minimally processed and non-refined items. In light of this,

dietary counselling and antenatal practitioners should tailor dietary guidance to match the SES and ethnicity of the patient whenever possible.

Our study has several strengths, including prospective data collection, large sample size, adequate power to allow for comparisons between ethnic groups, and detailed measurement of diet using validated, ethnic-specific FFQs. There are some limitations to our analyses, including self-reported retrospective food intake, the inability to account for differences in cooking methods and residual confounding, which is a potential bias in all cohort studies. To minimise this potential bias, we conducted a sensitivity analysis and excluded mothers who may have altered their diet due to a medical condition, such as diagnosed gestational diabetes or hypertension in pregnancy. After these exclusions, our results remained consistent.

CONCLUSION

Maternal consumption of a plant-based diet during pregnancy is associated with infant birth weight. Among white Europeans a plant-based diet is associated with lower birth weight, reduced odds of LGA and increased odds of SGA babies, whereas among South Asians living in Canada, a plant-based diet is associated with higher birth weight. Future studies are necessary to replicate these results and to elucidate potential mechanisms that underlie these ethnic-specific associations.

Author affiliations

¹Department of Medicine, McMaster University, Hamilton, Ontario, Canada

²School of Food Science and Nutrition, University of Leeds, Leeds, UK

³Department of Health Research Methods, Evidence, and Impact, McMaster University, Hamilton, Ontario, Canada

⁴Population Health Research Institute, Hamilton Health Sciences and McMaster University, Hamilton, Ontario, Canada

⁵Canadian Cardiovascular Research Network, Brampton, Ontario, Canada

⁶Six Nations Health Services, Ohsweken, Ontario, Canada

⁷Department of Paediatrics, McMaster University, Hamilton, Ontario, Canada

⁸Hospital for Sick Children and Department of Paediatrics, University of Toronto, Toronto, Ontario, Canada

⁹Department of Immunology, Faculty of Medicine, University of Manitoba, Winnipeg, Manitoba, Canada

¹⁰Department of Pediatrics, Faculty of Medicine and Dentistry, University of Alberta, Edmonton, Alberta, Canada

¹¹Department of Paediatrics, Faculty of Medicine, University of British Columbia, BC Children's Hospital, Vancouver, British Columbia, Canada

Contributors SSA, ABB, JB, RJdS, DD, DLL, JW, PS, PM, SET, SA, MG, SM, KMM, MRS, KKT and GW conceptualised the study and acquired the data. MAZ, RJdS and SSA designed the study. MAZ, RJdS, MRS and SSA conceptualised and designed statistical modelling, conducted and interpreted the analysis, and drafted the manuscript. MAZ, RJdS, JB, MRS and SSA checked and validated statistical and mathematical assumptions. MAZ, RJdS, MRS and SSA prepared the final manuscript. All authors critically reviewed and approved the final manuscript.

Funding This work was supported by funding by a Canadian Institute for Health Research (CIHR) Grant in Food & Health Population Health Research grant [RFA# 201301FH6; 2013-2018; SAA, ABB, JB, RJdS, JAD, MG, SDM, AM, DM, KMM, GP, MRS, PJ, MGS, KKT, GW] and a CIHR RCT Fellowship grant [MTP201410; MAZ]. START data was collected as part of a bilateral ICMR /CIHR funded program [INC-109205] and HSF Canada Grant in Aid [NA7283]. CIHR provided funds for CHILD [AEC85761; ABB, PM, MRS, PS, SET] with additional funding from AllerGen NCE Inc. for Nutrigen [SSA] and the CHILD Study (MRS). CIHR and HSF Canada provided

funds for FAMILY [KKT], and ABC [SSA] data collection. MAZ was the recipient of a Young Investigator Award from the South Asian Network Supporting Awareness and Research (SANSAR). SDM holds a Canada Research Chair Tier II in Maternal and Child Obesity Prevention and Intervention. MRS holds the Astra Zeneca chair in Respiratory Epidemiology. SSA holds a Canada Research Chair Tier 1 in Ethnic Diversity and Cardiovascular Disease and a Heart and Stroke Foundation/Michael G De Groote Chair in Population Health Research.

Competing interests None declared.

Ethics approval McMaster University. Ethical approval was obtained independently for all studies from the Hamilton Integrated Research Ethics Board—CHILD (REB 07–2929), FAMILY (REB 02–060), START (REB 10–640) and ABC (REB 12–152). Additional approval for ABC was obtained from the Six Nations Band Council.

Provenance and peer review Not commissioned; externally peer reviewed.

Data sharing statement Data sharing is currently not available for most data due to ethical restrictions.

Open Access This is an Open Access article distributed in accordance with the Creative Commons Attribution Non Commercial (CC BY-NC 4.0) license, which permits others to distribute, remix, adapt, build upon this work non-commercially, and license their derivative works on different terms, provided the original work is properly cited and the use is non-commercial. See: <http://creativecommons.org/licenses/by-nc/4.0/>

© Article author(s) (or their employer(s) unless otherwise stated in the text of the article) 2017. All rights reserved. No commercial use is permitted unless otherwise expressly granted.

REFERENCES

- Mikolajczyk RT, Zhang J, Betran AP, *et al*. A global reference for fetal-weight and birthweight percentiles. *Lancet* 2011;377:1855–61.
- Mebratu TF, Feltbower RG, Parslow RC. Effects of birth weight and growth on childhood wheezing disorders: findings from the Born in Bradford Cohort. *BMJ Open* 2015;5:e009553.
- Nordman H, Voutilainen R, Laitinen T, *et al*. Growth and cardiovascular risk factors in prepubertal children born large or small for gestational age. *Horm Res Paediatr* 2016;85:11–17.
- Kernell K, Sydsjö G, Bladh M, *et al*. Congenital heart disease in men - birth characteristics and reproduction: a national cohort study. *BMC Pregnancy Childbirth* 2014;14:187.
- Martin JA, Hamilton BE, Osterman MJ, *et al*. Births: final data for 2013. National vital statistics reports: from the centers for disease control and prevention, national center for health statistics. *National Vital Statistics System* 2015;64:1.
- National Health Services. *Hospital Episode Statistics: National Health Service*.
- Statistics Canada. *CANSIM. Table 102-4509. Catalogue 8400210-X*.
- Oliveira AP, Kalra S, Wahi G, *et al*. Maternal and newborn health profile in a first nations community in Canada. *J Obstet Gynaecol Can* 2013;35:905–13.
- Küpers LK, Xu X, Jankipersadsing SA, *et al*. DNA methylation mediates the effect of maternal smoking during pregnancy on birthweight of the offspring. *Int J Epidemiol* 2015;44:1224–37.
- Kim SY, Sharma AJ, Sappenfield W, *et al*. Association of maternal body mass index, excessive weight gain, and gestational diabetes mellitus with large-for-gestational-age births. *Obstet Gynecol* 2014;123:737–44.
- Knudsen VK, Heitmann BL, Halldorsson TI, *et al*. Maternal dietary glycaemic load during pregnancy and gestational weight gain, birth weight and postpartum weight retention: a study within the danish national birth cohort. *Br J Nutr* 2013;109:1471–8.
- Dwarkanath P, Barzilay JR, Thomas T, *et al*. High folate and low vitamin B-12 intakes during pregnancy are associated with small-for-gestational age infants in South Indian women: a prospective observational cohort study. *Am J Clin Nutr* 2013;98:1450–8.
- de Souza RJ, Zullyniak MA, Desai D, *et al*. Harmonization of food-frequency questionnaires and dietary pattern analysis in 4 ethnically diverse birth cohorts. *J Nutr* 2016;146:2343–50.
- Subbarao P, Anand SS, Becker AB, *et al*. The Canadian Healthy Infant Longitudinal Development (CHILD) Study: examining developmental origins of allergy and asthma. *Thorax* 2015;70:998–1000.
- Morrison KM, Atkinson SA, Yusuf S, *et al*. The Family Atherosclerosis Monitoring In early life (FAMILY) study: rationale, design, and baseline data of a study examining the early determinants of atherosclerosis. *Am Heart J* 2009;158:533–9.

16. Anand SS, Vasudevan A, Gupta M, *et al.* Rationale and design of South Asian Birth Cohort (START): a Canada-India collaborative study. *BMC Public Health* 2013;13:79.
17. Wahi G, Wilson J, Miller R, *et al.* Aboriginal birth cohort (ABC): a prospective cohort study of early life determinants of adiposity and associated risk factors among Aboriginal people in Canada. *BMC Public Health* 2013;13:608.
18. Fred Hutchinson Cancer Research Center. Food Frequency Questionnaires (FFQ). [sharedresources.fredhutch.org. http://sharedresources.fredhutch.org/services/food-frequency-questionnaires-ffq](http://sharedresources.fredhutch.org/services/food-frequency-questionnaires-ffq) (accessed 24 Jun2016).
19. Willett WC, Stampfer M, Willett W, eds. *Implications of total energy intake for epidemiologic analyses*: Nutritional epidemiology, 1998.
20. Hanley GE, Janssen PA. Ethnicity-specific birthweight distributions improve identification of term newborns at risk for short-term morbidity. *Am J Obstet Gynecol* 2013;209:428.e1–428.e6.
21. Wright CM, Parkinson KN. Postnatal weight loss in term infants: what is normal and do growth charts allow for it? *Arch Dis Child Fetal Neonatal Ed* 2004;89:F254–F257.
22. Piccoli GB, Clari R, Vigotti FN, *et al.* Vegan-vegetarian diets in pregnancy: danger or panacea? a systematic narrative review. *BJOG* 2015;122:623–33.
23. Rogers I, Emmett P, Baker D, *et al.* Financial difficulties, smoking habits, composition of the diet and birthweight in a population of pregnant women in the South West of England. ALSPAC Study Team. Avon longitudinal study of pregnancy and childhood. *Eur J Clin Nutr* 1998;52:251–60.
24. McCowan LM, Dekker GA, Chan E, *et al.* Spontaneous preterm birth and small for gestational age infants in women who stop smoking early in pregnancy: prospective cohort study. *BMJ* 2009;338:b1081–1.
25. Surkan PJ, Hsieh CC, Johansson AL, *et al.* Reasons for increasing trends in large for gestational age births. *Obstet Gynecol* 2004;104:720–6.
26. Thompson JM, Clark PM, Robinson E, *et al.* Risk factors for small-for-gestational-age babies: The Auckland birthweight collaborative study. *J Paediatr Child Health* 2001;37:369–75.
27. Sanders TA, Reddy S. The influence of a vegetarian diet on the fatty acid composition of human milk and the essential fatty acid status of the infant. *J Pediatr* 1992;120:S71–S77.
28. Knudsen VK, Orozova-Bekkevold IM, Mikkelsen TB, *et al.* Major dietary patterns in pregnancy and fetal growth. *Eur J Clin Nutr* 2008;62:463–70.
29. Gammon CS, von Hurst PR, Coad J, *et al.* Vegetarianism, vitamin B12 status, and insulin resistance in a group of predominantly overweight/obese South Asian women. *Nutrition* 2012;28:20–4.
30. Muthayya S, Kurpad AV, Duggan CP, *et al.* Low maternal vitamin B12 status is associated with intrauterine growth retardation in urban South Indians. *Eur J Clin Nutr* 2006;60:791–801.
31. West KP, Shamim AA, Mehra S, *et al.* Effect of maternal multiple micronutrient vs iron-folic acid supplementation on infant mortality and adverse birth outcomes in rural Bangladesh: the JiVitA-3 randomized trial. *JAMA* 2014;312:2649–58.
32. Potdar RD, Sahariah SA, Gandhi M, *et al.* Improving women's diet quality preconceptionally and during gestation: effects on birth weight and prevalence of low birth weight—a randomized controlled efficacy trial in India (Mumbai Maternal Nutrition Project). *Am J Clin Nutr* 2014;100:1257–68.
33. De Wals P, Tairou F, Van Allen MI, *et al.* Reduction in neural-tube defects after folic acid fortification in Canada. *N Engl J Med* 2007;357:135–42.
34. Lesser IA, Gasevic D, Lear SA. The association between acculturation and dietary patterns of South Asian immigrants. *PLoS One* 2014;9:e88495.
35. Barakat H. Effect of frying-cooking on nutritional and bioactive compounds of innovative ovo-vegetarian diets. *Food Nutr Sci* 2014;05:1577–90.
36. Rees GA, Doyle W, Srivastava A, *et al.* The nutrient intakes of mothers of low birth weight babies - a comparison of ethnic groups in East London, UK. *Matern Child Nutr* 2005;1:91–9.
37. Rao S, Yajnik CS, Kanade A, *et al.* Intake of micronutrient-rich foods in rural Indian mothers is associated with the size of their babies at birth: Pune Maternal Nutrition Study. *J Nutr* 2001;131:1217–24.
38. Wandel M, Råberg M, Kumar B, *et al.* Changes in food habits after migration among South Asians settled in Oslo: the effect of demographic, socio-economic and integration factors. *Appetite* 2008;50:376–85.
39. McDonald JT, Kennedy S. Is migration to Canada associated with unhealthy weight gain? overweight and obesity among Canada's immigrants. *Soc Sci Med* 2005;61:2469–81.
40. Sanou D, O'Reilly E, Ngnie-Teta I, *et al.* Acculturation and nutritional health of immigrants in Canada: a scoping review. *J Immigr Minor Health* 2014;16:24–34.
41. Aune D, Giovannucci E, Boffetta P, *et al.* Fruit and vegetable intake and the risk of cardiovascular disease, total cancer and all-cause mortality—a systematic review and dose-response meta-analysis of prospective studies. *Int J Epidemiol* 2017;46:1029–56.
42. Aune D, Keum N, Giovannucci E, *et al.* Whole grain consumption and risk of cardiovascular disease, cancer, and all cause and cause specific mortality: systematic review and dose-response meta-analysis of prospective studies. *BMJ* 2016;353:i2716.