

BMJ Open

BMJ Open is committed to open peer review. As part of this commitment we make the peer review history of every article we publish publicly available.

When an article is published we post the peer reviewers' comments and the authors' responses online. We also post the versions of the paper that were used during peer review. These are the versions that the peer review comments apply to.

The versions of the paper that follow are the versions that were submitted during the peer review process. They are not the versions of record or the final published versions. They should not be cited or distributed as the published version of this manuscript.

BMJ Open is an open access journal and the full, final, typeset and author-corrected version of record of the manuscript is available on our site with no access controls, subscription charges or pay-per-view fees (<http://bmjopen.bmj.com>).

If you have any questions on BMJ Open's open peer review process please email editorial.bmjopen@bmj.com

BMJ Open

The influencing factors of dysmenorrhea among hospital nurses: A questionnaire survey in Taiwan

Journal:	<i>BMJ Open</i>
Manuscript ID	bmjopen-2017-017615
Article Type:	Research
Date Submitted by the Author:	04-May-2017
Complete List of Authors:	Chiu, Min-Hui; Kaohsiung Medical University College of Nursing, Doctoral Student; Chi Mei Medical Center, Liouying, Department of Nursing Hsieh, Hsiu-Fen; Kaohsiung Medical University, College of Nursing Yang, Yi-Hsin; Kaohsiung Med Univ Chen, Huei-Mein; Chung Hwa University of Medical Technology Hsu, Su-Chen; Chi Mei Medical Center Wang, Hsiu-Hung; Kaohsiung Medical University, College of Nursing
Primary Subject Heading:	Communication
Secondary Subject Heading:	Intensive care
Keywords:	dysmenorrhea, prevalence rate, self-care behavior, hospital nurses

SCHOLARONE™
Manuscripts

The influencing factors of dysmenorrhea among hospital nurses: A questionnaire survey in Taiwan

ABSTRACT

Objectives: Nurses are at a high risk for dysmenorrhea while working intensely and focusing on their patients. The purposes of the current study were to (1) describe the basic demographic data, dysmenorrheal knowledge, and menstrual attitudes of dysmenorrheal and non-dysmenorrheal hospital nurses; (2) analyze these nurse populations to identify significant differences; and (3) examine their risk factors.

Methods: This cross-sectional survey employed a structured questionnaire. Two hospitals in southern Taiwan were recruited. The participants included (1) full-time hospital nurses (2) at least 18 years of age (3) who agreed to participate. All participants were recruited through random sampling. The questionnaire contained demographic data, the Dysmenorrheic Knowledge Scale, and the Menstrual Attitude Scale (MAS).

Results: A total of 420 nurses completed all questionnaires. Among them, 297 (70.7%) had experienced dysmenorrhea in the past six months, while 123 (29.3%) had not. Significant differences in age, marital status, childbearing status, age of menarche,

1
2
3
4 and rotating three-shift ratio were identified between the dysmenorrhea and
5
6 non-dysmenorrhea groups. Analysis of the MAS results revealed significant
7
8 differences between the groups regarding consideration of menstruation as a
9
10 debilitating or a bothersome event, anticipation and prediction of menstruation onset,
11
12 and denial of any effects from menstruation. The results of multiple logistic
13
14 regression showed that the predictive factors included age less than 40 years, working
15
16 three-shift rotations, marital status, acknowledging menstruation as a debilitating
17
18 event, anticipation and prediction of the onset of menstruation, and denial of any
19
20 effects from menstruation.
21
22
23
24
25
26
27
28
29

30 **Conclusions:** The risk factors of dysmenorrhea are age, working three-shift rotations,
31
32 marital status, acknowledging menstruation as a debilitating event, anticipation and
33
34 prediction of the onset of menstruation, and denial of any effects from menstruation
35
36 are predictive. These results will help managers design health-promoting menstrual
37
38 self-care programs for hospital nurses.
39
40
41
42
43

44 **Keywords:** dysmenorrhea, prevalence rate, self-care behavior, hospital nurses
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

STRENGTHS AND LIMITATIONS

- This study explored the risk factors for dysmenorrhea among nurses.
- The prevalence rate of dysmenorrhea in women is high and negatively affects their daily activity.
- The risk factors of dysmenorrhea are complex; thus, we used multiple logistic regression analysis to identify these factors.
- The study was limited in analyzing the effects of lifestyle choices such as smoking and drinking because few nurses reported those behaviors.
- The findings of the present study can improve health-promoting self-care and friendly environments for hospital nurses.

INTRODUCTION

More than 50% of women experience dysmenorrhea, defined as pain that accompanies menstruation. The majority (75.1%) of women believe dysmenorrhea to be a normal part of their lives and that the symptoms will continue to affect their daily lives until they near menopause¹ Dysmenorrhea occurs during the first one to three years after menarche and is commonly accompanied by sweating, lack of appetite, headache, distraction, nausea, vomiting, dizziness², and depression³. Dysmenorrhea arises during menstrual bleeding due to lower abdominal pain brought on by menses, with no relevant diagnoses from other gynecological examinations⁴. The prevalence rates of dysmenorrhea differ between age groups; however, over half of women in all age groups experience dysmenorrhea. The prevalence rate of dysmenorrhea is between 50.9% and 87.4% worldwide⁵⁻⁷ Parker et al found that 21% of Australian adolescents experienced severe dysmenorrhea, with 26% having school absences due to dysmenorrhea.⁸

In Taiwan, adolescent vocational nursing students 16–18 years of age had a dysmenorrhea prevalence rate of 73.3%, and another study reported a rate of 90.2% among the general population of women.¹⁰ A study on hospital nurses in Taiwan reported a prevalence rate of 90.7%.¹¹ Despite the high prevalence rate, nurses often neglect their own health while working and caring for patients.¹²

1
2
3
4 Dysmenorrhea can be cyclical or chronic. It can be possibly a result of pain
5
6 catastrophizing, and is sensitive to anxiety¹³. It not only causes physical pain but also
7
8 affects mental well-being and quality of life, leading to work or school absenteeism
9
10 and significant health burdens¹⁴. This problem not only affects performance but also
11
12 the daily lives of adolescent students in Taiwan⁹ and may cause women to take days
13
14 off, which may affect their work¹⁵. A study in Malaysia reported poor concentration,
15
16 absenteeism, and poor grades to be common factors of dysmenorrhea that affect daily
17
18 life. Despite its high prevalence rate and effect on daily life, 76.1% of women still
19
20 believe that dysmenorrhea is a natural part of a women's menstrual cycle, and only
21
22 14.8% believe that treatment is necessary. In terms of dysmenorrhea information, 62.3%
23
24 of adolescent girls claimed that they learned about the condition from their mothers,
25
26 while 52.9% said that they had obtained their knowledge from peers¹⁶. Besides
27
28 influencing daily learning, dysmenorrhea also causes tiredness, depression, difficulty
29
30 concentrating, etc. In addition, dysmenorrhea is negatively correlated with
31
32 self-awareness of one's own health; in other words, individuals with dysmenorrhea
33
34 have poorer self-awareness regarding their own health⁶. Another study of Malaysian
35
36 college students found that social activities, responsibilities, time spent resting, class
37
38 absenteeism, and daily life were affected in individuals with dysmenorrheal
39
40 individuals¹⁷.
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

1
2
3
4 In Taiwan, a study of 417 fifth- and sixth-grade students reported that
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

In Taiwan, a study of 417 fifth- and sixth-grade students reported that dysmenorrhea affected the emotions, daily lives, school activities, and social activities of 74.8%, 73.1%, 61.6%, and 50.1% of students, respectively⁸. Another study of 165 15- to 19-year-old students found that 82.4% experienced disruptions in daily activities because of dysmenorrhea and 12.7% reported school absences due to dysmenorrhea¹⁸. A study of 297 nurses (average age: 30.3 years) found that 8.4% had taken menstrual leave, 11.8% had booked a leave of absence for dysmenorrhea, and 28.6% saw their doctors due to the condition.¹⁵ Furthermore, 20.8% of adolescent girls from vocational schools in the same region had seen doctors for dysmenorrhea⁹. Menstrual leave is currently available in Japan, Indonesia, South Korea, and Taiwan. Focus groups studies in Taiwan showed that most interviewed women had menstrual discomfort and that they understood the implementation and regulation of menstrual leave in different companies. Yet in reality, the application rate is low due to factors including “the regulations are too rigid,” “no one else has taken menstrual leave,” “other kinds of leaves are available,” “no one will take my shift,” or “there has to be a doctor’s note”¹⁹.

Previous studies showed that the physical and mental health of nurses are significantly correlated with job satisfaction, tiredness, and comfort^{19,20}. The Gender Equality in Employment Act of Taiwan, approved in 2002, clearly defines regulations

1
2
3
4 regarding menstrual leave in which a female employee who has trouble working due
5
6
7 to discomfort during menstruation should take one day of menstrual leave every
8
9
10 month²¹. However, because of the particularity of nursing, temporary leave affects
11
12 the allocation of nurses. Cheng¹⁰ found that the long-term use of medicine to maintain
13
14 a ready-to-work status is a common experience among women; nurses, who have
15
16 frequent contact with drugs, use them extensively to overcome menstrual discomfort
17
18 at work²¹, including the use of sumatriptan–naproxen sodium to reduce the effect of
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
dysmenorrhea during work and leisure²².

Several risk factors affect dysmenorrhea, such as the following criteria. (a) Age: older women are generally less likely to have dysmenorrhea, although the relationship to marital status or birth condition remains unclear. (b) Age of menarche: age of menarche is significantly related to the occurrence of dysmenorrhea, with women with an earlier age of menarche more likely to experience more severe dysmenorrhea^{7 17} (c) Duration of menstrual cycle: a Nigerian study noted that longer menstrual flows are an important predictive factor of dysmenorrhea.⁷ (d) Regularity of menstrual cycle: women with irregular menstrual cycles are more likely to have dysmenorrhea⁹. In terms of lifestyle factors, the results of several studies indicated that smoking and alcohol consumption are risk factors for primary dysmenorrhea and that women with smoking or drinking habits are more likely to have dysmenorrhea²³

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

²⁴ With regard to work, several studies found that women who work rotations, especially nurses who have to work “sometimes day shifts, sometimes night shifts,” have relatively more serious menstrual discomfort ²¹. Nurses who work night shifts are more likely to have dysmenorrheal symptoms during their menstrual cycles compared to those who work regular shifts ¹⁵.

Furthermore, the attitudes of adolescent students toward menstruation correlate with the occurrence of dysmenorrheal ²⁵. Those who tend to agree that menstruation is a debilitating experience and those who anticipate and predict the onset of menstruation are more likely to have dysmenorrhea ²⁶. The effect of dysmenorrhea on women varies based on the degree and frequency of their pain.

Previous studies have revealed that women have inadequate knowledge about menstruation and that educational information about menstrual health care contains contradictory messages, including the idea that menstruation is a normal and natural event, but should be hidden ²⁷. Cheng ²⁸ observed that some female students knew about menstruation taboos; while they had doubts about those taboos, they still had negative attitudes about menstruation. Medical expenses due to dysmenorrhea increase with age, besides the topic of women menstrual health and their emotional problems, should also be taken seriously ²⁹. Nevertheless, since dysmenorrhea is so common, and women consider it a normal phenomenon and rarely seek medical

1
2
3
4 treatment due to their cultural and religious attitudes¹⁵ or due to the feeling that they
5
6
7 can tolerate the pain²³.
8

9
10 In Taiwan, 98.8% of hospital nurses are women. There is a high prevalence rate
11
12 of dysmenorrhea in women; due to the nature of nursing work, nurses are at a
13
14 high-risk for dysmenorrhea. However, previous studies on nurses did not discuss the
15
16 influencing factors. The purposes of this study were to (1) describe the basic
17
18 demographic data as well as menstruation characteristics, knowledge about
19
20 dysmenorrhea, and menstrual attitude of dysmenorrheal and non-dysmenorrheal
21
22 clinical nurses; (2) analyze the demographic data as well as menstruation
23
24 characteristics, knowledge about dysmenorrhea, and menstrual attitude of
25
26 dysmenorrheal and non-dysmenorrheal clinical nurses, and identify significant
27
28 differences between these groups; and (3) examine the risk factors of dysmenorrhea in
29
30 clinical nurses.
31
32
33
34
35
36
37
38
39
40
41
42
43

44 **METHODS**

45 **Study design and participants**

46
47
48 This study used a cross-sectional questionnaire survey based on structured
49
50 questionnaires for data collection. The participants were recruited from two medical
51
52 institutions in southern Taiwan. One institution was a medical center with 1,300 beds;
53
54
55
56
57
58
59
60

1
2
3
4 the other was a regional hospital with 650 beds. The participants included 2,000
5
6
7 clinical nurses. The inclusion criteria for this study included (1) full-time nurses at the
8
9
10 study hospitals (2) at least 18 years of age who (3) agreed to participate in this study
11
12 after reading an informed consent document. The sample size was estimated using
13
14 JMP 7.0 with an effect size of 0.3, α of 0.05, power of 0.80, resulting in a required
15
16 sample size of 350 individuals. With a predicted questionnaire recovery rate of 80%,
17
18 the sample size was estimated to be 438. We randomly sampled 450 nurses and
19
20
21 recovered 420 valid questionnaires (93.3%). Among the 420 nurses, 297 (70.7%) had
22
23
24 experienced dysmenorrhea in the past six months and were categorized in the
25
26
27 dysmenorrhea group, while the 123 (29.3%) nurses who had not experienced
28
29
30 dysmenorrhea were classified as the non-dysmenorrhea group.
31
32
33

34
35
36 Random sampling was conducted after this study received approval from the
37
38 Institutional Review Board (IRBD No. CLH0088) using lists of nurses from the two
39
40 hospitals. The questionnaires were delivered to the nurses by a trained research
41
42 assistant. The participants provided written consent and answered the questionnaires
43
44 independently. The written consent form indicated that participation in this study was
45
46 not compulsory and the questionnaire was anonymous. Anyone who was not willing
47
48 to participate in this study could simply choose not to return the questionnaire. No
49
50
51 individual rights or interests were compromised during this process. Those who
52
53
54
55
56
57
58
59
60

1
2
3
4 completed the questionnaire received a gift. The questionnaires were collected two
5
6
7 weeks after being issued, upon which the written consents and questionnaires were
8
9
10 filed separately and placed in locations chosen by the participants to protect their
11
12
13 personal privacy.

14 15 16 17 18 *Measurements*

19
20 This study was conducted in May 2010. The study instruments included the
21
22 Demographic Inventory (DI), Dysmenorrheal Knowledge Scale (DKS), and
23
24 Menstrual Attitude Scale (MAS). The DI was designed based on related studies^{9 30}
25
26 and included age, age of menarche, marital and childbirth status, discussion of
27
28 menstrual hygiene, dysmenorrhea experience, medical experience of dysmenorrhea,
29
30 working conditions, self-care health education experience regarding menstruation,
31
32 dysmenorrhea influences, and work coping. The DKS was designed using both the
33
34 researchers' clinical experiences and previous studies.[30] The scale included 20
35
36 questions with yes/no answers. Correct answers received a score of 1, while incorrect
37
38 answers received a score of 0. The total summed scores ranged from 0 to 20, with
39
40 higher scores representing a better knowledge of dysmenorrhea. The
41
42 Kuder-Richardson reliability coefficient was 0.63 in a previous study³⁰, and 0.72 in
43
44 this study. The MAS was developed by Brooks-Gunn and Ruble (1980) and modified
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

1
2
3
4 by Lee (1994). It includes 32 items with five dimensions, including menstruation as a
5
6 debilitating event, menstruation as a bothersome event, menstruation as a natural
7
8 event, anticipation and prediction of the onset of menstruation, and denial of the
9
10 effects of menstruation. Each item was scored on a 7-point Likert scale from 1
11
12 (disagree strongly) to 7 (agree strongly). The total summed scores ranged from 32 to
13
14 224, with higher scores representing a more positive attitude toward menstruation.
15
16
17
18
19
20
21 Cronbach's α was 0.58 in the original scale³¹, 0.61 and 0.75 in Morrison's study³²,
22
23
24 and 0.57 in the present study.
25
26
27
28
29

30 *Data Analysis*

31
32
33
34
35
36 The data in this study were stored using Microsoft Excel. JMP 7.0 (a business
37
38 unit of SAS) for statistical analysis. Based on the study's variables and purposes,
39
40 percentage, mean, and standard deviation were used to describe the demographics and
41
42 menstruation characteristics as well as the DKS and MAS distributions in the
43
44 dysmenorrhea and the non-dysmenorrhea groups. T- and chi-square tests were used to
45
46
47
48 detect differences in demographics, menstruation characteristics, DKS, and MAS
49
50
51 between the two groups. Finally, multiple logistic regression analysis was used to
52
53
54
55
56 examine the relationships between each variable and the occurrence of dysmenorrhea.
57
58
59
60

RESULTS

A total of 420 participants completed the questionnaire. Among them, 297 (70.7%) had experienced dysmenorrhea in the past six months, while 123 (29.3%) had not. Compared to the non-dysmenorrhea group, the dysmenorrhea group was significantly younger, had higher percentage of unmarried status, had no childbearing experience, had a higher percentage of age of menarche of less than 12 years, and higher percentage of those who worked a three-shift rotation (Table 1).

Table 1. Comparison of demographic and menstruation characteristics between the dysmenorrhea and non-dysmenorrhea groups

Variables	Dysmenorrheal (n = 297)		Non-dysmenorrheal (n = 123)		t or χ^2	p-value
	n	%	n	%		
Age (years) M \pm SD	30.33 \pm 3.61		32.04 \pm 4.47		-3.78	<0.001
Marital status					20.03	<0.001
Unmarried	231	77.78	69	56.10		
Married	66	22.22	54	43.90		
Childbearing					19.38	<0.001
No	247	83.16	78	63.41		
Yes	50	16.84	45	36.59		
Age of menarche					4.70	0.030
< 12	46	15.49	11	8.94		
\geq 12	251	84.51	112	91.06		
Regularity of menstrual cycle					0.10	0.755
Yes	193	64.98	85	69.11		
No	104	35.02	38	30.89		
Duration of menstrual cycle					1.99	0.575
\leq 4 days	110	37.04	43	34.96		
5-6 days	157	52.86	70	56.91		
\geq 7 days	30	10.10	10	8.13		
Prior health education on dysmenorrhea					0.99	0.321
No	163	54.88	74	60.16		
Yes	134	45.12	49	39.84		
Three-shift rotation					6.06	0.014
Yes	271	91.25	102	82.93		
No	26	8.75	21	17.07		

Table 1. Comparison of Demographic and Menstruation Characteristics Between Dysmenorrheal and Non-Dysmenorrheal Groups (continue)

Variables	Dysmenorrheal (n = 297)		Non-dysmenorrheal (n = 123)		t or χ^2	p-value
	n	%	n	%		
	Whether to know it is allowed to ask for menstrual leave					
Yes	242	81.48	107	86.99		
No	55	18.52	16	13.01		
Whether to consider menstrual leave necessary					1.68	.195
Yes	288	96.97	115	93.50		
No	9	3.03	8	6.50		
Have ever asked for menstrual leave					0.14	.707
No	272	91.58	114	92.68		
Yes	25	8.42	9	7.32		
Have ever asked for more leaves for dysmenorrhea?					3.59	.058
No	262	88.22	116	94.31		
Yes	35	11.78	7	5.69		

In terms of dysmenorrheal knowledge and menstrual attitudes, the dysmenorrhea group showed a significantly higher degree of acknowledgment than that of the non-dysmenorrhea group in considering menstruation a debilitating event, considering menstruation a bothersome event, and anticipation and prediction of menstruation onset. In terms of denial of the effects of menstruation, the dysmenorrhea group was significantly lower than that of the non-dysmenorrhea group. No significant differences were observed between groups in terms of considering menstruation a natural event (Table 2).

Table 2. Comparison of dysmenorrheic knowledge and menstrual attitude between the dysmenorrhea and non-dysmenorrhea groups

Variables	Dysmenorrheal (n = 297)		Non-dysmenorrheal (n = 123)		t-value	p-value
	M	SD	M	SD		
	Dysmenorrheic knowledge scale	14.12	2.19	13.40		
Menstrual attitude scale	148.64	12.74	146.85	12.69	1.31	0.192
Menstruation as a debilitating event	56.44	8.52	52.55	7.65	4.58	<.001
Menstruation as a bothersome event	29.05	5.01	27.54	6.64	2.27	0.024
Menstruation as a natural event	19.98	5.19	20.98	4.92	-1.87	0.062
Anticipation and prediction of the onset of menstruation	20.04	3.01	18.98	3.55	2.93	0.004
Denial of any effect of menstruation	23.12	6.75	26.80	5.70	5.70	<.001

Age less than 40 years, working three-shift rotations, and marital status were the predictive factors for the occurrence of dysmenorrhea. In terms of menstrual attitudes using mean as the contact point, the tendency to acknowledge menstruation as a debilitating event, and the tendency to deny the effects of menstruation were also predictive factors for dysmenorrhea occurrence. After controlling for working three-shift rotations, marital status, the tendency to acknowledge menstruation as a debilitating event, and the tendency to acknowledge the effects of menstruation, the rate of occurrence of dysmenorrhea in those younger than 40 years was 4.46 times

higher than that in those over 40 years of age. After controlling for age, marital status, the tendency to acknowledge menstruation as a debilitating event, and the tendency to acknowledge the effects of menstruation, the dysmenorrhea occurrence rate for those who worked three-shift rotations was 2.07 times higher than that in those who did not. After controlling for age, working three-shift rotations, marital status, and the tendency to acknowledge the effects of menstruation, those who acknowledged that menstruation was a debilitating event had a dysmenorrhea occurrence rate that was 2.72 times higher than that of those who did not. After controlling for age, working three-shift rotations, marital status, and the tendency to acknowledge menstruation as a debilitating event, those who acknowledged the effects of menstruation had a dysmenorrhea occurrence rate that was 2.59 times higher than that in those who did not (Table 3).

Table 3: Risk factors of dysmenorrhea

		Crude OR	95% CI	P-Value	Adjusted OR	95% CI	P-Value
Age	Age < 40	4.06	(1.33 , 13.7)	<0.05	4.46	(1.24 , 17.23)	<0.05
	Age ≥ 40	1.00	(- , -)		1.00	(- , -)	
Three-shift rotation	Rotating	2.15	(1.15 , 3.98)	<0.05	2.07	(1.01 , 4.21)	<0.05
	3-shift	1.00	(- , -)		1.00	(- , -)	

Married	No	2.74 (1.75 , 4.30)	<0.001	2.59 (1.57 , 4.28)	<0.001
	Yes	1.00 (- , -)		1.00 (- , -)	
Menstruation as a Higher debilitating event	Lower	2.53 (1.63 , 3.95)	<0.001	2.72 (1.67 , 4.52)	<0.001
	Higher	1.00 (- , -)		1.00 (- , -)	
Denial of any effect of menstruation	Lower	2.83 (1.83 , 4.46)	<0.001	2.59 (1.61 , 4.23)	<0.001
	Higher	1.00 (- , -)		1.00 (- , -)	

Reference group: Age \geq 40 years, no three-shift rotation, unmarried, reduced menstruation as a debilitating event, and increased denial of any effect on menstruation

CI, confidence interval; OR, odds ratio; SE, standard error.

DISCUSSION

Although the average age of the dysmenorrhea group was significantly lower than that of the non-dysmenorrhea group, when age was the only variable being compared, the dysmenorrhea occurrence rate of nurses younger than 40 years was 4.06 times higher than that of those over 40 years. After controlling for three-shift rotations, marital status, the tendency to acknowledge menstruation as a debilitating event, and the tendency to deny the effects of menstruation, the occurrence rate of dysmenorrhea in those under 40 years was 4.46 times higher than that in those over 40 years. Therefore, age was a risk factor for dysmenorrhea. To our knowledge, no other studies have directly discussed the correlation between age and dysmenorrhea. However, previous studies reported that dysmenorrhea occurrence rates were over 76%

1
2
3
4 in adolescent girls or students^{5 6 16}, compared to a rate of 55.9% in a study with 15- to
5
6
7 45-year-old women³³, with a decrease in the occurrence rate with increasing
8
9
10 participants' age. Since dysmenorrhea is part of being a woman, periodic
11
12 dysmenorrhea comes with the menstrual cycle. With age, women's dysmenorrhea and
13
14 coping experience also increase; therefore, dysmenorrheal women are "female
15
16 specialists with embodied knowledge," meaning that their experiences may affect
17
18 their subjective perceptions and tolerance of dysmenorrhea, which in turn affect the
19
20 occurrence of dysmenorrhea³⁴.
21
22
23
24
25
26
27
28
29

30 In terms of marital and childbirth status, both the marital status ratio and
31
32 childbirth ratio in the dysmenorrhea group were significantly lower than those in the
33
34 non-dysmenorrhea group, indicating a high dysmenorrhea occurrence rate among
35
36 unmarried, non-childbearing women. After adjusting for age, working three-shift
37
38 rotations, marital status, the tendency to acknowledge menstruation as a debilitating
39
40 event, and the tendency to deny the effects of menstruation, the dysmenorrhea
41
42 occurrence rate in unmarried women was 2.59 times higher than that in married
43
44 women. This result is consistent with the results of Chung³⁵ (average age = 27 years),
45
46
47 which reported a higher dysmenorrhea occurrence rate in single than in married and
48
49 divorced nurses³⁵. Many women are by their doctors that their dysmenorrhea will
50
51
52
53
54
55
56
57
58
59
60

1
2
3
4 improve after getting married and giving birth, which may improve menstrual blood
5
6 discharge. Furthermore, women may perform positive self-care behaviors for
7
8 menstruation after getting married due to childbearing needs, which may also be a
9
10 factor that indirectly relieves dysmenorrhea.¹¹ In terms of menstruation history, the
11
12 dysmenorrhea occurrence rate in women who had their menarche after 12 years of age
13
14 was significantly lower in the dysmenorrhea group than that in the non-dysmenorrhea
15
16 group, indicating that those who had menarche at a younger age were more likely to
17
18 have dysmenorrhea; however, after performing further logistic regression analysis,
19
20 this variable was not significant. Therefore, after adjusting, age of menarche was not
21
22 an important influencing factor for dysmenorrhea. Previous studies of dysmenorrhea
23
24 in adolescent girls or students found early menarche to be a risk factor for
25
26 dysmenorrhea.^{7 17 36} With age and dysmenorrhea experience, women develop coping
27
28 abilities; together with marital and childbearing status, these factors could weaken the
29
30 effect of age of menarche on dysmenorrhea occurrence.
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

In terms of work, the dysmenorrhea occurrence rate among those who worked three-shift rotations was significantly higher in the dysmenorrhea group than that in the non-dysmenorrhea group. After controlling for age, marital status, the tendency to acknowledge menstruation as a debilitating event, and the tendency to deny the

1
2
3
4 effects of menstruation, the dysmenorrhea occurrence rate in those who worked
5
6 three-shift rotations was 2.07 times higher than the rate in those who did not. This
7
8 indicates that work type (e.g., rotating nightshift) is a risk factor for dysmenorrhea
9
10 occurrence in nurses. However, Chung³⁵ found that rotating work shifts were not
11
12 significantly related to dysmenorrhea occurrence in nurses. Furthermore, one study on
13
14 Japanese junior high school students also found that nightshift work was not
15
16 significantly associated with dysmenorrhea³⁷. However, an interview of focus groups
17
18 conducted by Chang found that working rotating shifts was highly associated with
19
20 menstrual discomfort¹⁹. Due to the nature of nursing work, most nurses need to work
21
22 rotating night shifts, and the graveyard shift can easily cause uncomfortable menstrual
23
24 cycles or obvious irregularities²⁵. Coupling nurses' day/night activities and sleep
25
26 patterns with the increased pressure from working night shifts, menstrual discomfort
27
28 may be more common in those with frequent rotation changes.
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

In terms of menstrual attitude, the dysmenorrhea group was significantly higher than the non-dysmenorrhea group in considering menstruation to be a debilitating event, considering menstruation to be a bothersome event, and anticipating and predicting the onset of menstruation, while significantly lower than the non-dysmenorrhea group in denying the effects of menstruation. There was no

1
2
3
4 significant difference between the two groups with regard to considering menstruation
5
6
7 to be a natural event. In multiple logistic regressions, the tendency to acknowledge
8
9
10 menstruation as a debilitating event and the tendency to deny the effects of
11
12
13 menstruation were influencing factors for the occurrence of dysmenorrhea. In this
14
15
16 study, menstrual attitude was related to dysmenorrhea occurrence, particularly
17
18
19 considering menstruation to be a debilitating event and denying the effects of
20
21
22 menstruation. Firat found no significant correlation between menstrual discomfort and
23
24
25 menstrual attitude³⁸. In vocational nursing students, “menstruation is a debilitating
26
27
28 event” and “anticipation and prediction of the onset of menstruation” in the
29
30
31 dysmenorrhea group scored significantly higher than those in the non-dysmenorrhea
32
33
34 group. In terms of “menstruation is a bothersome event” and “menstruation is a
35
36
37 natural event,” the non-dysmenorrhea group was significantly higher than the
38
39
40 dysmenorrhea group; and no significant differences were observed between the two
41
42
43 groups in terms of “denial of any effect of menstruation”²⁶. These results are different
44
45
46 from those of the present study. The most obvious difference between the two studies
47
48
49 is the tendency to deny the effect of menstruation. The scores in the dysmenorrhea
50
51
52 and non-dysmenorrhea groups in the present study were 23.1 and 26.8, respectively,
53
54
55 compared to 26.6 and 28.0 in adolescent nursing students. In other words, compared
56
57
58 to adolescent nursing students, nurses in the present study did not deny the effects
59
60

1
2
3
4 caused by menstruation and recognized that they were affected by menstruation.
5
6
7 When a nurse experiences dysmenorrhea, her work may be affected. If she asks for
8
9
10 menstrual leave, temporary work arrangements and patient care may be affected.
11
12 Since dysmenorrhea is a periodic phenomenon, frequently asking for menstrual leave
13
14 may affect working relationships; therefore, nurses usually choose to work even when
15
16 they experience dysmenorrhea.
17
18
19

20
21 In this study, age under 40 years, working three-shift rotations, marital status,
22
23 using mean as the contact point, the tendency to acknowledge menstruation as a
24
25 debilitating event, and the tendency to deny the effects of menstruation were
26
27 predictive factors for dysmenorrhea occurrence. Chiou's ³⁰ study of adolescent
28
29 nursing students found that menstrual regularity and health education were predictive
30
31 factors for dysmenorrhea. Those who had irregular menstruations and had not
32
33 received dysmenorrhea-related health education had higher dysmenorrhea occurrence
34
35 rates ⁹; however, these two factors were not predictive factors in the present study.
36
37 Our participants were nurses from a wide range of age groups rather than adolescents;
38
39 therefore, they had more dysmenorrhea and self-care experience. In addition,
40
41 menstrual health is a part of nursing education; therefore, although nurses are not
42
43 directly involved in health education, education is a part of their professional training.
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

1
2
3
4 Thus, health education was not an important predictive factor in this study.
5
6
7
8

9
10 This study mainly evaluated the predictive factors for dysmenorrhea in hospital
11 nurses and aimed to build a friendly work environment for high-risk women. The
12 participants were randomly selected nurses from only two hospitals in southern
13 Taiwan; therefore, the results might be limited. In future studies, inclusion research
14 settings can be extended to examine any differences between different living areas. In
15 addition, rotating night shifts are required in many other occupations besides nursing.
16
17 Considering different professional backgrounds and life experiences, future studies
18 should also study women from different occupations with rotating shifts to evaluate
19 differences in the influence of their knowledge and attitudes on dysmenorrhea
20 occurrence. The variables assessed in this study included demographic characteristics,
21 menstruation history, dysmenorrheal knowledge, menstrual attitudes, etc.; however,
22 lifestyle choices such as smoking, drinking, and exercising were not included. Future
23 studies are necessary to analyze the effects of these lifestyle choices on dysmenorrhea.
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49

50 CONCLUSIONS

51
52 Nursing is a female-dominated occupation in Taiwan. The high prevalence rate
53 of dysmenorrhea severely affects nurses such that they may not be able to focus on
54
55
56
57
58
59
60

1
2
3
4 their work, thereby affecting the quality of patient care. Nurses may need to take
5
6 menstrual leave, which can cause a shortage of nursing resources. The predictive
7
8 factors for dysmenorrhea occurrence were assessed with the hope to produce
9
10 interventions for high-risk groups and improve workplace comfort for nurses. In this
11
12 study, age less than 40 years, working three-shift rotations, marital status, using mean
13
14 as the contact point, the tendency to acknowledge menstruation as a debilitating event,
15
16 and the tendency to deny the effects of menstruation were predictive factors for the
17
18 occurrence of dysmenorrhea. This finding could help nursing managers to offer
19
20 appropriate assistance for high-risk groups and build a health-promoting self-care and
21
22 friendly environment for hospital nurses.
23
24
25
26
27
28
29
30
31
32
33
34
35

36 In terms of clinical practices, if the majority of nurses are under 40 years of age
37
38 and unmarried, establishing a menstruation-supporting environment should be
39
40 considered. Since women with dysmenorrhea do not necessarily need to take a full
41
42 day off, if a supportive environment can be created that allows them to take a break
43
44 from work, then their comfort will be improved. Work environments, particularly
45
46 hospitals, can provide measures such as flexibility with menstrual leave or scheduling
47
48 menstrual leave, integrating menstrual self-care measures into annual occupational
49
50 safety and health training, and providing spaces at work for short breaks together with
51
52
53
54
55
56
57
58
59
60

thermotherapy, aromatherapy massage, etc. These measures will enable nurses to care for themselves at work, thus improving their workplace comfort and increasing their work satisfaction.

ACKNOWLEDGEMENTS

The authors thank the participants for their participation and the nurse managers of these two hospitals for their administrative assistance. We also thank Editage for their professional English review of this manuscript.

Competing interests

None.

Funding

This research received no specific grant from any funding agency in the public, commercial, or not-for-profit sectors.

Contributors

Min-Hui Chiu and Hsiu-Hung Wang designed the study, analyzed and interpreted the data, and drafted the manuscript. Min-Hui Chiu contributed to the study design, analysis, and interpretation of the data. All authors had full access to all data in the study and take responsibility for its integrity.

Data sharing statement

No additional data are available.

REFERENCES

1. Chen CX, Kwেকেboom KL, Ward SE. Beliefs About Dysmenorrhea and Their Relationship to Self-Management. *Res Nurs Health* 2016;39(4):263-76.
2. Eryilmaz G, Ozdemir F. Evaluation of menstrual pain management approaches by Northeastern Anatolian adolescents. *Pain management nursing : official journal of the American Society of Pain Management Nurses* 2009;10(1):40-7.
3. Kato T. Effects of Flexibility in Coping with Menstrual Pain on Depressive Symptoms. *Pain practice : the official journal of World Institute of Pain* 2017;17(1):70-77.
4. Zhu X, Wong F, Bensoussan A, Lo SK, Zhou C, Yu J. Are there any cross-ethnic differences in menstrual profiles? A pilot comparative study on Australian and Chinese women with primary dysmenorrhea. *The journal of obstetrics and gynaecology research* 2010;36(5):1093-101.
5. Abdul-Razzak KK, Ayoub NM, Abu-Taleb AA, Obeidat BA. Influence of dietary intake of dairy products on dysmenorrhea. *The journal of obstetrics and gynaecology research* 2010;36(2):377-83.
6. Agarwal AK, Agarwal A. A study of dysmenorrhea during menstruation in adolescent girls. *Indian journal of community medicine : official publication of Indian Association of Preventive & Social Medicine* 2010;35(1):159-64.
7. Loto OM, Adewumi TA, Adewuya AO. Prevalence and correlates of dysmenorrhea among Nigerian college women. *The Australian & New Zealand journal of obstetrics & gynaecology* 2008;48(4):442-4.
8. Chang YT, Chen YC. Menstrual Health Care Behavior and Associated Factors Among Female Elementary Students in the Hualien Region. *Journal of Nursing Research* 2008;16(1):9.
9. Chiou MH, Wang HH. Predictors of Dysmenorrhea and Self-Care Behavior Among Vocational Nursing School Female Students. *Journal of Nursing Research* 2008;16(1):17-25.
10. Cheng HF, Lin YH. Selection and efficacy of self-management strategies for dysmenorrhea in young Taiwanese women. *Journal of clinical nursing* 2011;20(7-8):1018-25.
11. Palmer D. "To Help a Million Sick, You Must Kill a Few Nurses": Nurses' Occupational Health, 1890–1914. *Nursing History Review* 2012;20(1):14-45.
12. Laschinger HK, Grau AL. The influence of personal dispositional factors and organizational resources on workplace violence, burnout, and health outcomes in new graduate nurses: a cross-sectional study. *International journal of nursing studies* 2012;49(3):282-91.

13. Payne LA, Rapkin AJ, Lung KC, Seidman LC, Zeltzer LK, Tsao JC. Pain Catastrophizing Predicts Menstrual Pain Ratings in Adolescent Girls with Chronic Pain. *Pain Med* 2016;17(1):16-24.
14. Berkley KJ, McAllister SL. Don't dismiss dysmenorrhea! *Pain* 2011;152(9):1940-1.
15. Chiu MH, Wang HH, Hsu SC, Liu IP. Dysmenorrhoea and self-care behaviours among hospital nurses: a questionnaire survey. *Journal of clinical nursing* 2013;22(21-22):3130-40.
16. Wong LP. Attitudes towards dysmenorrhoea, impact and treatment seeking among adolescent girls: a rural school-based survey. *The Australian journal of rural health* 2011;19(4):218-23.
17. Zukr SM, Naing L, Hamzah TNT, Hussain NHN. Primary Dysmenorrhea among Medical and Dental University Students in Kelantan: Prevalence and Associated Factors. *International Medical Journal* 2009;16(2):93-99.
18. Lu I-C. Dysmenorrhea and related factors in Taiwanese adolescent girls. *University of Texas at Austin* 2010.
19. Chang C, Chen FL, Chang CH, Hsu CH. A preliminary study on menstrual health and menstrual leave in the workplace in Taiwan. *Taiwan J Public Health* 2011;30(5):436-50.
20. Schluter PJ, Turner C, Huntington AD, Bain CJ, McClure RJ. Work/life balance and health: the Nurses and Midwives e-cohort study. *International Nursing Review* 2011(58):28-36.
21. Cady RK, Diamond ML, Diamond MP, Ballard JE, Lener ME, Dorner DP, et al. Sumatriptan-naproxen sodium for menstrual migraine and dysmenorrhea: satisfaction, productivity, and functional disability outcomes. *Headache* 2011;51(5):664-73.
22. Bettendorf B, Shay S, Tu F. Dysmenorrhea: contemporary perspectives. *Obstetrical and Gynecological Survey* 2008;63(9):597-603.
23. Perry M. Treatment options for dysmenorrhoea. *Practice Nursing* 2012;23(4):4.
24. Ju H, Jones M, Mishra GD. Smoking and trajectories of dysmenorrhoea among young Australian women. *Tobacco control* 2016;25(2):195-202.
25. Chiou MH, Wang HH. The Relationship Between Dysmenorrhea and Menstrual Attitudes Among Female Students in Vocational Nursing Schools. *The Journal of Nursing* 2004;51(4):45-52.
26. Parker MA, Sneddon AE, Arbon P. The menstrual disorder of teenagers (MDOT) study: determining typical menstrual patterns and menstrual disturbance in a large population-based study of Australian teenagers. *BJOG : an international journal of obstetrics and gynaecology* 2010;117(2):185-92.

- 1
 - 2
 - 3
 - 4
 - 5
 - 6
 - 7
 - 8
 - 9
 - 10
 - 11
 - 12
 - 13
 - 14
 - 15
 - 16
 - 17
 - 18
 - 19
 - 20
 - 21
 - 22
 - 23
 - 24
 - 25
 - 26
 - 27
 - 28
 - 29
 - 30
 - 31
 - 32
 - 33
 - 34
 - 35
 - 36
 - 37
 - 38
 - 39
 - 40
 - 41
 - 42
 - 43
 - 44
 - 45
 - 46
 - 47
 - 48
 - 49
 - 50
 - 51
 - 52
 - 53
 - 54
 - 55
 - 56
 - 57
 - 58
 - 59
 - 60
27. Stubbs ML. Cultural perceptions and practices around menarche and adolescent menstruation in the United States. *Annals of the New York Academy of Sciences* 2008;1135:58-66.
28. Cheng CY, Yang K, Liou SR. Taiwanese adolescents? gender differences in knowledge and attitudes towards menstruation. *Nursing & Health Sciences* 2007;9(2):127-34.
29. H LY. The Medical Utilization of Chinese and Western Medicine in Females with Dysmenorrhea. Kaohsiung Medical University, 2012.
30. Chiou M-H, Wang H-H, Yang Y-H. Effect of Systematic Menstrual Health Education on Dysmenorrheic Female Adolescents' Knowledge, Attitudes, and Self-care Behavior. *Kaohsiung J Med Sci* 2007;23(4):183-90.
31. S L. Prevalence and Associated Factors of Premenstrual Syndrome Among Nurses in Taiwan. In: Ministry of Science and Technology, R. O. C., editor, 1994.
32. Morrison LA, Larkspur L, Calibuso MJ, Brown S. Women's Attitudes About Menstruation and Associated Health and Behavioral Characteristics. *American Journal of Health Behavior* 2010;34(1):90-100.
33. Brooks-Gunn J, Ruble D. The menstrual attitude questionnaire. *Psychosomatic medicine* 1980;42:503-11.
34. Lin YX. *When "Experts" Meet the Medical Professionals: The Women with Menstrual Pain Encountering Western Medicine*. Kaohsiung Medical University, Taiwan, 2006.
35. Chung F, Yao C, Wan G. The associations between menstrual function and life style/working conditions among nurses in Taiwan. *Journal of occupational health* 2005;47(2):149-56.
36. Nagata C, Hirokawa K, Shimizu N, Shimizu H. Associations of menstrual pain with intakes of soy, fat and dietary fiber in Japanese women. *European journal of clinical nutrition* 2005;59(1):88-92.
37. Takeuchi H, Oishi T, Harada T. Morningness-eveningness preference, and mental and physical symptoms during the menstrual cycle of Japanese junior high school students. *Sleep and Biological Rhythms* 2003(1):245-47.
38. Firat MZ, Kulakac O, Oncel S, Akcan A. Menstrual Attitude Questionnaire: confirmatory and exploratory factor analysis with Turkish samples. *Journal of advanced nursing* 2009;65(3):652-62.

BMJ Open

Influencing factors of dysmenorrhea among hospital nurses: A questionnaire survey in Taiwan

Journal:	<i>BMJ Open</i>
Manuscript ID	bmjopen-2017-017615.R1
Article Type:	Research
Date Submitted by the Author:	13-Sep-2017
Complete List of Authors:	Chiu, Min-Hui; Kaohsiung Medical University College of Nursing, Doctoral Student; Chi Mei Medical Center, Liouying, Department of Nursing Hsieh, Hsiu-Fen; Kaohsiung Medical University, College of Nursing Yang, Yi-Hsin; Kaohsiung Med Univ Chen, Huei-Mein; Chung Hwa University of Medical Technology Hsu, Su-Chen; Chi Mei Medical Center Wang, Hsiu-Hung; Kaohsiung Medical University, College of Nursing
Primary Subject Heading:	Communication
Secondary Subject Heading:	Intensive care
Keywords:	dysmenorrhea, prevalence rate, self-care behavior, hospital nurses

SCHOLARONE™
Manuscripts

Title Page

Manuscript title

Influencing factors of dysmenorrhea among hospital nurses: A questionnaire survey in Taiwan

Running head

Dysmenorrhea and influencing factors

Authors

Min-Hui Chiu

1. Doctoral Student, College of Nursing, Kaohsiung Medical University, Taiwan
Address: No. 100 Shih-Chuan 1st Road, San-Ming District, Kaohsiung 80708, Taiwan
2. Director, Chi Mei Medical Center, Liouying, Tainan, Taiwan
Address: No.201, Taikang, Liouying Dist., Tainan City 736, Taiwan
Email: 300015@mail.chimei.org.tw

Hsiu-Fen Hsieh

Associate Professor, College of Nursing, Kaohsiung Medical University, Taiwan
Address: No. 100 Shih-Chuan 1st Road, San-Ming District, Kaohsiung 80708, Taiwan
Email: hsiufen96@gmail.com

Yi-Hsin Yang

Professor, College of Pharmacy, Kaohsiung Medical University, Kaohsiung, Taiwan
Address: No. 100 Shih-Chuan 1st Road, San-Ming District, Kaohsiung 80708, Taiwan
E-mail: yihhsya@kmu.edu.tw
Telephone: 886-7-3121101 ext. 2262

Huei-Mein Chen

Associate Professor, Chung Hwa University of Medical Technology, Tainan, Taiwan
Address: No. 89, Wen-hua1st St, Rende District, Tainan 71703, Taiwan
E-mail: hueimein@mail.hwai.edu.tw
Telephone: 886-6-2671214 ext. 512

Su-Chen Hsu

Chi Mei Medical Center, Tainan, Taiwan

1
2
3 Address: No.901, Zhonghua Rd., Yongkang Dist., Tainan City 710, Taiwan

4 E-mail: 300001@mail.chimei.org.tw

5
6 Telephone: 886-6-2812811 ext. 53000

7
8 Fax: 886-6-2827480

9
10
11 **Hsiu-Hung Wang***

12 Professor, College of Nursing, Kaohsiung Medical University, Taiwan

13 No. 100, Shih-Chuan 1st Road, Kaohsiung City 80708, Taiwan

14 Tel: 886-7-3121101 ext 2624

15 Fax: 886-7-3218364

16 E-mail: hhwang@kmu.edu.tw

17
18
19
20
21 **Corresponding author**

22 Hsiu-Hung Wang, RN, FAAN, PhD

23 Professor, College of Nursing, Kaohsiung Medical University

24 No. 100, Shih-Chuan 1st Road, Kaohsiung City 80708, Taiwan

25 Tel: 886-7-3121101 ext 2624

26 Fax: 886-7-3218364

27 E-mail: hhwang@kmu.edu.tw

28
29
30
31
32
33 **keywords**

34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
Dysmenorrhea, prevalence rate, self-care behavior, hospital nurses

Word count

4718

Influencing factors of dysmenorrhea among hospital nurses:

A questionnaire survey in Taiwan

ABSTRACT

Objectives: Nurses are at high risk for dysmenorrhea while working closely with patients. The study objectives were to: (1) describe basic demographic data and menstruation characteristics, knowledge about dysmenorrhea, and menstrual attitudes of dysmenorrheal and non-dysmenorrheal hospital nurses; (2) analyze the demographic data and menstruation characteristics, knowledge about dysmenorrhea, and menstrual attitudes of the both groups, and identify significant differences between the groups; and (3) examine influencing factors of dysmenorrhea among them.

Methods: This cross-sectional survey used a structured questionnaire, administered at two hospitals in southern Taiwan. Participants included full-time hospital nurses at least 18 years of age and agreed to participate. All participants were recruited through random sampling. The questionnaire included demographic data, the Dysmenorrhea Knowledge Scale, and Menstrual Attitude Scale (MAS).

Results: A total 420 nurses completed the questionnaire. Among them, 297 (70.7%)

1
2
3
4 had experienced dysmenorrhea in the past 6 months; 123 (29.3%) had not. Significant
5
6 differences in age ($p < 0.001$), marital status ($p < 0.001$), childbearing status ($p < 0.001$),
7
8 age of menarche ($p < 0.05$), and rotating three-shift ratio ($p < 0.05$) were identified
9
10 between the dysmenorrhea and non-dysmenorrhea groups. Analysis of MAS results
11
12 revealed significant differences between the groups regarding consideration of
13
14 menstruation as a debilitating ($p < 0.001$) or bothersome event ($p < 0.05$), anticipation
15
16 and prediction of menstruation onset ($p < 0.01$), and denial of any effects from
17
18 menstruation ($p < 0.001$). Results of multiple logistic regression showed that predictive
19
20 factors included age less than 40 years (4.46 vs. 1.00), working three-shift rotations
21
22 (2.07 vs. 1.00), marital status (2.59 vs. 1.00), acknowledging menstruation as a
23
24 debilitating event (2.72 vs. 1.00), and denial of effects from menstruation (2.59 vs.
25
26 1.00).
27
28
29
30
31
32
33
34
35
36
37
38

39 **Conclusions:** These findings could help nursing managers to create a caring climate
40
41 and friendly work environment for hospital nurses at risk of dysmenorrhea.
42
43

44 **Keywords:** Dysmenorrhea, prevalence rate, self-care behavior, hospital nurses
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

STRENGTHS AND LIMITATIONS

- The influencing factors of dysmenorrhea are complex; in this study, multiple logistic regression analysis was used to control for confounding variables and to identify these influencing factors.
- The findings of this study can improve health-promoting self-care and create friendly environments for hospital nurses.
- Participants in this study were aged 22–48 years; therefore, the results have limited application to hospital nurses aged under 22 years and more than 48 years.
- This study was limited in analyzing the effects of lifestyle choices, such as smoking and drinking, because few nurses reported those behaviors.

INTRODUCTION

More than 50% of women experience dysmenorrhea, defined as pain that accompanies menstruation. The majority (75.1%) of women believe dysmenorrhea to be a normal part of their lives and that the symptoms will continue to affect their daily life until they near menopause.¹ Dysmenorrhea occurs during the first 1 to 3 years after menarche and is commonly accompanied by sweating, lack of appetite, headache, distraction, nausea, vomiting, dizziness,² and depression.³ Dysmenorrhea arises during menstrual bleeding owing to lower abdominal pain brought on by menses, with no relevant diagnoses of other gynecological disorders.⁴ The prevalence rates of dysmenorrhea differ by age; however, over half of women in all age groups experience dysmenorrhea. The prevalence rate of dysmenorrhea worldwide is between 50.9% and 87.4%.⁵⁻⁷ Parker et al. found that 21% of Australian adolescents experienced severe dysmenorrhea, with 26% having school absences due to dysmenorrhea.⁸

In Taiwan, adolescent vocational nursing students 16–18 years of age had a reported dysmenorrhea prevalence rate of 73.3%;⁹ another study reported a rate of 90.2% among the general population of women in Taiwan.¹⁰ A study on hospital nurses in Taiwan reported a prevalence rate of 90.7%.¹¹ Despite the high prevalence of dysmenorrhea, nurses often neglect their own health while working and caring for

1
2
3
4 patients.¹²
5
6

7 Dysmenorrhea can be cyclical or chronic; in some cases, it can be a possible result
8
9 of pain catastrophizing and anxiety sensitivity.¹³ Dysmenorrhea not only causes
10
11 physical pain, it also affects mental well-being and quality of life, leading to work or
12
13 school absenteeism and significant health burdens.¹⁴⁻¹⁸ Despite its high prevalence
14
15 rate and effect on daily life, 76.1% of women still believe that dysmenorrhea is a
16
17 natural part of a women's menstrual cycle, and only 14.8% believe that treatment is
18
19 necessary.¹⁶ In terms of dysmenorrhea information, 62.3% of adolescent girls claim
20
21 that they learned about the condition from their mothers, and 52.9% state that they
22
23 obtained their knowledge from peers.¹⁶ In addition, dysmenorrhea is negatively
24
25 correlated with self-awareness of one's own health; in other words, individuals with
26
27 dysmenorrhea have poorer self-awareness regarding their own health.⁶
28
29
30
31
32
33
34
35
36
37
38

39 A study of 297 nurses (average age 30.3 years) found that 8.4% had taken
40
41 menstrual leave, 11.8% had taken a leave of absence for dysmenorrhea, and 28.6%
42
43 saw their doctors owing to the condition.¹⁵ Furthermore, 20.8% of adolescent girls
44
45 from vocational schools in the same region had seen doctors for dysmenorrhea.⁹
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
Menstrual leave is currently available in Japan, Indonesia, South Korea, and Taiwan.
Focus group studies in Taiwan have showed that most women had menstrual
discomfort and they understood the implementation and regulation of menstrual leave

1
2
3
4 in different companies. However, in reality, the application rate is low because of
5
6 factors expressed as follows: “the regulations are too rigid”, “no one else has taken
7
8 menstrual leave”, “other kinds of leave are available”, “no one will take my shift”, or
9
10 “there has to be a doctor’s note”.¹⁹
11
12

13
14
15 Previous studies have showed that the physical and mental health of nurses are
16
17 significantly correlated with job satisfaction, tiredness, and comfort.^{19, 20} The Gender
18
19 Equality in Employment Act of Taiwan, approved in 2002, clearly defines the
20
21 regulations regarding menstrual leave, in which a female employee who has trouble
22
23 working due to discomfort during menstruation should take 1 day of menstrual leave
24
25 every month.²¹ However, because of the particularities of nursing, temporary leave
26
27 affects the allocation of nurses. Cheng found that the long-term use of medication to
28
29 maintain a ready-to-work status is common among women. Nurses, who have
30
31 frequent contact with drugs, use them extensively to overcome menstrual discomfort
32
33 at work,¹⁰ including the use of sumatriptan–naproxen sodium to reduce the effects of
34
35 dysmenorrhea during work and leisure.^{21, 22}
36
37
38
39
40
41
42
43
44
45
46

47 Several risk factors affect dysmenorrhea, as follows. (a) Age: older women are
48
49 generally less likely to have dysmenorrhea, although the relationships with marital
50
51 status or childbirth history remain unclear. (b) Age of menarche: age of menarche is
52
53 significantly related to the occurrence of dysmenorrhea, with women who have an
54
55
56
57
58
59
60

1
2
3
4 earlier age of menarche more likely to experience more severe dysmenorrhea.^{7, 17} (c)
5
6
7 Duration of menstrual cycle: a Nigerian study noted that longer menstrual flow is an
8
9
10 important predictive factor of dysmenorrhea.⁷ (d) Regularity of menstrual cycle:
11
12 women with irregular menstrual cycles are more likely to have dysmenorrhea.⁹ In
13
14 terms of lifestyle factors, the results of several studies indicate that smoking and
15
16 alcohol consumption are risk factors for primary dysmenorrhea and that women who
17
18 habitually smoke or drink alcohol are more likely to have dysmenorrhea.^{23, 24} With
19
20 regard to work, several studies have found that women who work rotations, especially
21
22 nurses who have to alternate between day and night shifts, have relatively more
23
24 serious menstrual discomfort.²¹ Nurses who work night shifts are more likely to have
25
26 dysmenorrheal symptoms during their menstrual cycles compared with those who
27
28 work regular shifts.¹⁵
29
30
31
32
33
34
35
36
37

38
39 Furthermore, the attitudes of adolescent students toward menstruation correlate
40
41 with the occurrence of dysmenorrhea.²⁵ Adolescent girls who tend to feel that
42
43 menstruation is a debilitating experience and those who anticipate and predict the
44
45 onset of menstruation are more likely to experience dysmenorrhea.²⁶ The effect of
46
47 dysmenorrhea on women varies based on the degree and frequency of their pain.
48
49
50
51
52

53
54 Previous studies have revealed that women have inadequate knowledge about
55
56 menstruation and that educational information about menstrual health care contains
57
58
59
60

1
2
3
4 contradictory messages, including the idea that menstruation is a normal and natural
5
6 event but should be “invisible”.²⁷ Cheng observed that some female students knew
7
8 about menstruation taboos; while they had doubts about those taboos, they still had
9
10 negative attitudes about menstruation.²⁸ Medical expenses due to dysmenorrhea
11
12 increase with age; therefore, the topic of women’s menstrual health and
13
14 accompanying emotional challenges should be taken seriously.²⁹ Nevertheless,
15
16 because dysmenorrhea is so common, many women consider it a normal phenomenon
17
18 and rarely seek medical treatment owing to their cultural and religious attitudes¹⁵ or
19
20 because they feel that they can tolerate the pain.²³
21
22
23
24
25
26
27
28
29

30 In this study, we define dysmenorrhea as lower abdominal pain brought on by
31
32 menses at any time during the last 6 months.¹¹ In Taiwan, 98.8% of hospital nurses are
33
34 women. There is a high prevalence rate of dysmenorrhea among women; owing to the
35
36 nature of nursing work, nurses are at high risk for dysmenorrhea. However, previous
37
38 studies on nurses have not discussed the influencing factors. The objectives of this
39
40 study were to: (1) describe the basic demographic data as well as menstruation
41
42 characteristics, knowledge about dysmenorrhea, and menstrual attitudes of
43
44 dysmenorrheal and non-dysmenorrheal clinical nurses; (2) analyze the demographic
45
46 data as well as menstruation characteristics, knowledge about dysmenorrhea, and
47
48 menstrual attitudes of dysmenorrheal and non-dysmenorrheal clinical nurses, and
49
50
51
52
53
54
55
56
57
58
59
60

1
2
3
4 identify significant differences between these groups; and (3) examine the risk factors
5
6
7 of dysmenorrhea among clinical nurses.
8
9

10 11 12 **METHODS**

13 14 15 **Study design and participants**

16
17
18 This study used a cross-sectional questionnaire survey based on structured
19
20
21 questionnaires for data collection. Participants were recruited at two medical
22
23
24 institutions in southern Taiwan; one institution was a medical center with 1,300 beds,
25
26
27 the other was a regional hospital with 650 beds. The two institutions were at different
28
29
30 hospital levels within the same health care system. All 2,000 nurses at the two
31
32
33 institutions were potential participants. The inclusion criteria for this study were
34
35
36 full-time nurses at the study hospitals who were at least 18 years of age and agreed to
37
38
39 participate in the study after reading an informed consent document. The sample size
40
41
42 was estimated using JMP 7.0 (SAS Institute, Cary, NC, USA) with an effect size of
43
44
45 0.3, α of 0.05, and a power of 0.80, resulting in a required sample size of 350
46
47
48 individuals. With a predicted questionnaire recovery rate of 80%, the sample size was
49
50
51 estimated to be 438. We used simple random sampling to recruit the estimated number
52
53
54 of study participants. Each participant was chosen randomly and entirely by chance.
55
56
57 In total, 450 nurses were chosen for inclusion in the study. Among these 450 nurses,
58
59
60

1
2
3
4 finally 420 valid questionnaires were returned, with a response rate of 93.3%. Among
5
6 these 420 nurses, 297 (70.7%) had experienced dysmenorrhea in the past 6 months
7
8 and were categorized in the dysmenorrhea group; 123 (29.3%) nurses who had not
9
10 experienced dysmenorrhea were classified as the non-dysmenorrhea group.
11
12

13
14
15 Random sampling was conducted after the study received approval from the
16
17 Institutional Review Boards (IRBD No. CLH0088) of the two hospitals. The
18
19 questionnaires were administered to the nurses by a trained research assistant. All
20
21 participants provided their written consent and answered the questionnaires
22
23 independently. The written consent form indicated that participation in this study was
24
25 not compulsory and the questionnaire was anonymous. Anyone who was not willing
26
27 to participate in the study could simply choose not to return the questionnaire. No
28
29 individual rights or interests were compromised during this process. Those who
30
31 completed the questionnaire received a small gift. The questionnaires were collected 2
32
33 weeks after being issued, upon which the written consent forms and questionnaires
34
35 were filed separately and placed in locations chosen by participants, to protect their
36
37 personal privacy.
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52

53 **Measurements**

54
55
56 This study was conducted in May 2010. The study instruments included the
57
58
59
60

1
2
3
4 Demographic Inventory (DI), Dysmenorrheal Knowledge Scale (DKS), and
5
6 Menstrual Attitude Scale (MAS). The DI was designed based on related studies^{9, 30}
7
8 and included age, age of menarche, marital and childbirth status, discussion of
9
10 menstrual hygiene, dysmenorrhea experience, medical experience of dysmenorrhea,
11
12 menstrual hygiene, dysmenorrhea experience, medical experience of dysmenorrhea,
13
14 working conditions, self-care health education experience regarding menstruation,
15
16 dysmenorrhea influences, and work coping methods. The DKS was designed using
17
18 both the researchers' clinical experiences and previous studies.³⁰ The scale included
19
20 20 questions with yes/no answers, for example, "Women with dysmenorrhea are at
21
22 high risk for sterility" and "Applying hot compresses to the lower abdomen can ease
23
24 menstrual cramps". Correct answers received a score of 1, and incorrect answers
25
26 received a score of 0. The total summed scores ranged from 0 to 20, with higher
27
28 scores representing a better knowledge of dysmenorrhea. The Kuder-Richardson
29
30 reliability coefficient was 0.63 in a previous study³⁰ and 0.72 in this study. The MAS
31
32 was developed by Brooks-Gunn and Ruble (1980)³¹ and modified by Lee (1994). Lee
33
34 translated this scale into a Chinese version and deleted one items owing to cultural
35
36 differences.³² The Chinese version of MAS includes 32 items with five dimensions,
37
38 including consideration of menstruation as a debilitating event, menstruation as a
39
40 bothersome event, menstruation as a natural event, anticipation and prediction of the
41
42 onset of menstruation, and denial of the effects of menstruation. Each item is scored
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

1
2
3
4 on a 7-point Likert scale from 1 (strongly disagree) to 7 (strongly agree). The total
5
6
7 summed scores range from 32 to 224, with higher scores representing a more positive
8
9
10 attitude toward menstruation. Cronbach's α was 0.58 in the original scale³¹ and 0.61 in
11
12 Morrison's study.³³ Lo and Lin (1998) suggested that an acceptable Cronbach's α was
13
14 from 0.3 to 0.7.³⁴ As we were concerned about comparability with other studies, we
15
16
17 decided to use this scale with a Cronbach's α of 0.57 in the present study.
18
19

20 21 22 23 24 Data analysis

25
26
27 The data in this study were entered using Microsoft Excel. JMP 7.0 was used for
28
29
30 statistical analysis. Based on the study variables and objectives, percentage, mean,
31
32
33 and standard deviation were used to describe participant demographics and
34
35
36 menstruation characteristics as well as DKS and MAS distributions in the
37
38
39 dysmenorrhea and non-dysmenorrhea groups. We used *t*-tests and chi-square tests to
40
41
42 detect differences in demographics, menstruation characteristics, DKS, and MAS
43
44
45 between the two groups. Finally, multiple logistic regression analysis was used to
46
47
48 examine influencing factors (independent variables) of the occurrence of
49
50
51 dysmenorrhea (dependent variable). Independent variables with significant
52
53
54 differences were included in the multiple logistic regression analysis.
55
56
57
58
59
60

RESULTS

A total of 420 participants completed the questionnaire. Among them, 297 (70.7%) had experienced dysmenorrhea in the past 6 months, and 123 (29.3%) had not. Compared with the non-dysmenorrhea group, participants with dysmenorrhea were significantly younger ($t = -3.78$, $p < 0.001$), had a higher percentage of unmarried status (77.78%, $\chi^2 = 20.03$, $p < 0.001$), had no childbirth experience (83.16%, $\chi^2 = 19.38$, $p < 0.001$), had a higher percentage of age of menarche less than 12 years (15.49%, $\chi^2 = 4.70$, $p = 0.03$), and a higher percentage worked a three-shift rotation (91.25%, $\chi^2 = 6.06$, $p = 0.014$). The other variables showed no significant difference between the two groups (Table 1).

Table 1. Comparison of demographic and menstruation characteristics between the dysmenorrhea and non-dysmenorrhea groups

Variables	Dysmenorrheal (n = 297)		Non-dysmenorrheal (n = 123)		t or χ^2	p-value
	n	%	n	%		
Age (years) M \pm SD	30.33 \pm 3.61		32.04 \pm 4.47		-3.78	<0.001
Age < 40	292	98.32	115	93.50	6.74	<0.01
Age \geq 40	5	1.68	8	6.50		
Marital status					20.03	<0.001
Unmarried	231	77.78	69	56.10		
Married	66	22.22	54	43.90		
Childbearing					19.38	<0.001
No	247	83.16	78	63.41		
Yes	50	16.84	45	36.59		
Age of menarche					4.70	0.030
< 12	46	15.49	11	8.94		
\geq 12	251	84.51	112	91.06		
Regularity of menstrual cycle					0.10	0.755
Yes	193	64.98	85	69.11		
No	104	35.02	38	30.89		
Duration of menstrual cycle					1.99	0.575
\leq 4 days	110	37.04	43	34.96		
5-6 days	157	52.86	70	56.91		
\geq 7 days	30	10.10	10	8.13		
Prior health education on dysmenorrhea					0.99	0.321
No	163	54.88	74	60.16		
Yes	134	45.12	49	39.84		
Three-shift rotation					6.06	0.014
Yes	271	91.25	102	82.93		
No	26	8.75	21	17.07		

Table 1. Comparison of Demographic and Menstruation Characteristics Between Dysmenorrheal and Non-Dysmenorrheal Groups (continue)

Variables	Dysmenorrheal (n = 297)		Non-dysmenorrheal (n = 123)		t or χ^2	p-value
	n	%	n	%		
	Whether to know it is allowed to ask for menstrual leave					
Yes	242	81.48	107	86.99		
No	55	18.52	16	13.01		
Whether to consider menstrual leave necessary					1.68	0.195
Yes	288	96.97	115	93.50		
No	9	3.03	8	6.50		
Have ever asked for menstrual leave					0.14	0.707
No	272	91.58	114	92.68		
Yes	25	8.42	9	7.32		
Have ever asked for more leaves for dysmenorrhea?					3.59	0.058
No	262	88.22	116	94.31		
Yes	35	11.78	7	5.69		
Secondary dysmenorrhea					0.02	0.886
No	267	89.90	110	89.43		
Yes	30	10.10	13	10.57		

Note: Regularity of periods means that the duration between consecutive first-day menses is similar.

With respect to knowledge about dysmenorrhea, there was a non-significant difference between the two groups ($p = 0.238$). The highest scoring items among both groups were “applying hot compresses”, “adopting a knee-to-chest position”, and “drinking hot beverages including warm water or milk or ginger soup to ease cramps.” The lowest scoring items were “normal bleeding amount every cycle”, “dysmenorrhea

1
2
3
4 is a disease of the reproductive system”, and “women with dysmenorrhea are at high
5
6 risk for sterility.” Regarding menstrual attitudes, after standardizing, the highest
7
8 scoring dimension among the dysmenorrhea group was “considering menstruation a
9
10 debilitating event” (73.30) and “considering menstruation a natural event” (74.93) in
11
12 the non-dysmenorrhea group. The lowest scoring dimension in both groups was
13
14 “denial of any effect of menstruation” (47.18 in the dysmenorrhea group and 54.69 in
15
16 the dysmenorrhea group). Women with dysmenorrhea had significantly higher scores
17
18 than women in the non-dysmenorrhea group with respect to the following:
19
20 considering menstruation a debilitating event ($p < 0.001$), considering menstruation a
21
22 bothersome event ($p = 0.024$), and anticipation and prediction of menstruation onset
23
24 ($p = 0.004$). In terms of denying the effects of menstruation, scores for the
25
26 dysmenorrhea group were significantly lower than those of the non-dysmenorrhea
27
28 group ($p < 0.001$). No significant differences were observed between the groups in
29
30 terms of considering menstruation a natural event (Table 2).
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51

52 Table 2. Comparison of dysmenorrheic knowledge and menstrual attitude between the
53
54 dysmenorrhea and non-dysmenorrhea groups
55
56

Variables	Dysmenorrheal		Non-dysmenorrheal		t-value	p-value
	(n = 297)		(n = 123)			
	M	SD	M	SD		
Dysmenorrheic knowledge	14.12	2.19	13.40	2.40	1.18	0.238

1	scale						
2							
3							
4	Menstrual attitude scale	148.64	12.74	146.85	12.69	1.31	0.192
5							
6	Menstruation as a						
7	debilitating event	56.44	8.52	52.55	7.65	4.58	<0.001
8							
9	Menstruation as a						
10	bothersome event	29.05	5.01	27.54	6.64	2.27	0.024
11							
12	Menstruation as a natural						
13	event	19.98	5.19	20.98	4.92	-1.87	0.062
14							
15	Anticipation and prediction						
16	of the onset of menstruation	20.04	3.01	18.98	3.55	2.93	0.004
17							
18	Denial of any effect of						
19	menstruation	23.12	6.75	26.80	5.70	5.70	<0.001

20

21

22

23

24 We conducted a multiple logistic regression analysis using dysmenorrhea and

25 non-dysmenorrhea as dependent variables and significant differences between the

26 variables of demographics, menstruation characteristics, DKS, and MAS as predictors.

27

28

29

30 The results of the analysis showed that crucial predictive factors for the occurrence of

31 dysmenorrhea among hospital nurses include age less than 40 years, working

32 three-shift rotations, and marital status. In terms of menstrual attitudes, using the

33 mean as the contact point, the tendency to consider menstruation to be a debilitating

34 event and the tendency to deny the effects of menstruation were also predictive factors

35 for dysmenorrhea occurrence. The adjusted odd ratios are shown in Table 3.

36

37

38

39

40

41

42

43

44

45

46

47

48

49

50 After controlling for working three-shift rotations, marital status, the tendency to

51 consider menstruation a debilitating event, and the tendency to deny the effects of

52 menstruation, the rate of occurrence of dysmenorrhea in women younger than 40

53

54

55

56

57

58

59

60

years was 4.46 times higher than that in women over 40 years of age. After controlling for age, marital status, the tendency to consider menstruation a debilitating event, and tendency to deny the effects of menstruation, the dysmenorrhea occurrence rate for nurses who worked three-shift rotations was 2.07 times higher than that in nurses who did not. After controlling for age, working three-shift rotations, marital status, and the tendency to deny the effects of menstruation, participants who felt that menstruation was a debilitating event had a dysmenorrhea occurrence rate that was 2.72 times higher than that of participants who did not. After controlling for age, working three-shift rotations, marital status, and the tendency to think of menstruation as a debilitating event, women who did not acknowledge the effects of menstruation had a dysmenorrhea occurrence rate that was 2.59 times higher than that in women who did not deny these effects.

Table 3: Risk factors of dysmenorrhea

		Crude OR	95% CI	P-Value	Adjusted OR	95% CI	P-Value
Age	Age<40	4.06	(1.33 , 13.7)	<0.05	4.46	(1.24 , 17.23)	<0.05
	Age≥40	1.00	(- , -)		1.00	(- , -)	
Three-shift rotation	Yes	2.15	(1.15 , 3.98)	<0.05	2.07	(1.01 , 4.21)	<0.05
	No	1.00	(- , -)		1.00	(- , -)	
Married	No	2.74	(1.75 , 4.30)	<0.001	2.59	(1.57 , 4.28)	<0.001

	Yes	1.00 (- , -)		1.00 (- , -)
Menstruation as a Higher debilitating event	Lower	2.53 (1.63 , 3.95)	<0.001	2.72 (1.67 , 4.52)
	Higher	1.00 (- , -)		1.00 (- , -)
Denial of any effect of menstruation	Lower	2.83 (1.83 , 4.46)	<0.001	2.59 (1.61 , 4.23)
	Higher	1.00 (- , -)		1.00 (- , -)

Reference group: Age ≥ 40 years, no three-shift rotation, unmarried, reduced menstruation as a debilitating event, and increased denial of any effect on menstruation

CI, confidence interval; OR, odds ratio; SE, standard error.

DISCUSSION

Although the average age of participants in the dysmenorrhea group was significantly lower than that of women in the non-dysmenorrhea group, when age was the only variable being compared, the dysmenorrhea occurrence rate of nurses younger than 40 years was 4.06 times higher than that of nurses over age 40 years; after controlling for confounding variables, the rate of dysmenorrhea in younger women was 4.46 times higher than that in women over 40 years old. Therefore, age was a risk factor for dysmenorrhea. To our knowledge, no other studies have directly discussed the correlation between age and dysmenorrhea. However, previous studies have reported dysmenorrhea occurrence rates over 76% in adolescent girls or students,^{5, 6, 16} compared with a rate of 55.9% in a study among women aged 15–45-years,³¹ with a

1
2
3
4 decrease in the occurrence with increasing age. As dysmenorrhea is very common,
5
6
7 periodic dysmenorrhea comes naturally with the menstrual cycle. With age,
8
9
10 experiences of dysmenorrhea and coping among women also increase; therefore,
11
12
13 women with dysmenorrhea can be thought of as “female specialists with embodied
14
15
16 knowledge,” meaning that their experiences may affect their subjective perception and
17
18
19 tolerance of dysmenorrhea, which can in turn affect the occurrence of
20
21
22 dysmenorrhea.³⁵ Nursing is a female-dominated occupation in Taiwan. In this study,
23
24
25 97% of participants were younger than 40 years old. The high prevalence rate of
26
27
28 dysmenorrhea severely affects nurses such that they may not be able to focus on their
29
30
31 work, thereby affecting the quality of patient care. Nurses may need to take menstrual
32
33
34 leave, which can cause a shortage of nursing resources.

35
36 In terms of marital status, the unmarried ratio in the dysmenorrhea group was
37
38
39 2.74 times that of the non-dysmenorrhea group, indicating a high dysmenorrhea
40
41
42 occurrence rate among unmarried women; after adjusting for confounding variables, it
43
44
45 was 2.59 times higher than that among married women. This result is consistent with
46
47
48 those of Chung (average study participant age 27 years), which reported a higher
49
50
51 dysmenorrhea occurrence rate in single nurses than in married and divorced ones.³⁶
52
53
54 Many women are told by their doctors that their dysmenorrhea will improve after
55
56
57 getting married and giving birth, which may improve menstrual blood discharge.
58
59
60

1
2
3
4 Furthermore, women may engage in positive self-care for menstruation after getting
5
6 married for childbearing reasons, which may also be a factor that indirectly relieves
7
8 dysmenorrhea.¹¹ In terms of work, the dysmenorrhea occurrence rate among nurses in
9
10 the dysmenorrhea group who worked three-shift rotations was 2.15 times that of
11
12 nurses in the non-dysmenorrhea group; after adjusting for confounding variables, it
13
14 was 2.07 times higher than among nurses who did not work three-shift rotations. This
15
16 indicates that work type (e.g., rotating nightshifts) is a risk factor for dysmenorrhea in
17
18 nurses. However, Chung found that rotating work shifts were not significantly related
19
20 to dysmenorrhea occurrence in nurses.³⁶ Furthermore, one study among Japanese
21
22 junior high school students also found that nightshift work was not significantly
23
24 associated with dysmenorrhea.³⁷ However, an interview of focus groups conducted by
25
26 Chang found that working rotating shifts was highly associated with menstrual
27
28 discomfort.¹⁹ Owing to the nature of nursing work, most nurses need to work rotating
29
30 night shifts, and the graveyard shift can easily cause uncomfortable menstrual cycles
31
32 or obvious irregularities.²⁵ Coupling nurses' day/night activities and sleep patterns
33
34 with the increased pressure of working night shifts, menstrual discomfort may be
35
36 more common in those with frequent rotation changes.
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51

52
53 In terms of menstrual attitude, the dysmenorrhea group had significantly higher
54
55 scores than the non-dysmenorrhea group in considering menstruation to be a
56
57
58
59
60

1
2
3
4 debilitating event and significantly lower scores than the non-dysmenorrhea group in
5
6
7 denying the effects of menstruation. In multiple logistic regression for menstrual
8
9
10 attitude, the tendency to consider menstruation as a debilitating event in the
11
12
13 dysmenorrhea group was 2.53 times higher than that of the non-dysmenorrhea group;
14
15
16 after adjusting for confounding variables, women with dysmenorrhea had a 2.72 times
17
18
19 higher tendency to think of menstruation as a debilitating event than women who did
20
21
22 not consider menstruation to be debilitating. The tendency to deny the effects of
23
24
25 menstruation in the dysmenorrhea group was 2.83 times higher than the
26
27
28 non-dysmenorrhea group; after adjusting for confounding variables, women with
29
30
31 dysmenorrhea had a 2.59 times higher tendency to deny the effects of menstruation.
32
33
34 In this study, menstrual attitude was related to dysmenorrhea occurrence. Firat found
35
36
37 no significant correlation between menstrual discomfort and menstrual attitude.³⁸ In
38
39
40 that study among vocational nursing students with dysmenorrhea, “menstruation is a
41
42
43 debilitating event” and “anticipation and prediction of the onset of menstruation”
44
45
46 scored significantly higher than in those who did not have dysmenorrhea. In terms of
47
48
49 “menstruation is a bothersome event” and “menstruation is a natural event,” the
50
51
52 non-dysmenorrhea group scored significantly higher than the dysmenorrhea group; no
53
54
55 significant differences were observed between the two groups in terms of “denial of
56
57
58 any effect of menstruation”.²⁶ These results are different from those of the present
59
60

1
2
3
4 study. The most obvious difference between the two studies is the tendency to deny
5
6 the effects of menstruation. Scores in the dysmenorrhea and non-dysmenorrhea
7
8 groups in the present study were 23.1 and 26.8, respectively, compared with 26.6 and
9
10 28.0 among adolescent nursing students in Firat's study. In other words, compared
11
12 with adolescent nursing students, nurses in the present study did not deny the effects
13
14 caused by menstruation and recognized that they were affected by menstruation.
15
16 When a nurse experiences dysmenorrhea, her work may be affected. If she requests
17
18 menstrual leave, temporary work arrangements must be made and patient care may be
19
20 affected. Because dysmenorrhea is a periodic phenomenon, taking frequent menstrual
21
22 leave may affect working relationships; therefore, nurses usually choose to work even
23
24 when they experience dysmenorrhea.
25
26
27
28
29
30
31
32
33
34
35

36 In summary, influencing factors for dysmenorrhea occurrence included age under
37
38 40 years, working three-shift rotations, marital status, and using mean as the contact
39
40 point, the tendency to consider menstruation to be a debilitating event and tendency to
41
42 deny the effects of menstruation. Chiou's study of adolescent nursing students found
43
44 that menstrual regularity and health education were predictive factors for
45
46 dysmenorrhea.³⁰ Women who had irregular menstrual cycles and had not received
47
48 dysmenorrhea-related health education had higher dysmenorrhea occurrence rates;⁹
49
50 however, these two factors were not predictive factors in the present study. Our
51
52
53
54
55
56
57
58
59
60

1
2
3
4 participants were nurses from a wide range of age groups rather than only adolescents;
5
6
7 therefore, they had more experience with dysmenorrhea and self-care. In addition,
8
9
10 menstrual health is a part of nursing education; therefore, although nurses are not
11
12 directly involved in health education, education is a part of their professional training.
13
14 Thus, health education was not an important predictive factor in this study. This study
15
16 mainly evaluated the influencing factors for dysmenorrhea in hospital nurses with the
17
18 aim of helping to build a caring climate and friendly work environment for high-risk
19
20 nurses. The participants were randomly selected nurses from only two hospitals in
21
22 southern Taiwan; therefore, the results might have limited generalizability to nurses in
23
24 other parts of the country. In future studies, inclusion criteria should be extended to
25
26 examine any differences between different areas of Taiwan. In addition, rotating night
27
28 shifts are required in many other occupations besides nursing. Considering different
29
30 professional backgrounds and life experiences, future studies should also include
31
32 women from different occupations who work rotating shifts, to evaluate differences in
33
34 the influence of their knowledge and attitudes on dysmenorrhea occurrence. The
35
36 variables assessed in this study included demographic characteristics, menstruation
37
38 history, dysmenorrhea knowledge, and menstrual attitudes, among others; however,
39
40 lifestyle choices such as smoking, drinking, and exercising were not included. Future
41
42 studies are necessary to analyze the effects of these lifestyle choices on dysmenorrhea.
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

CONCLUSIONS

We assessed various influencing factors for dysmenorrhea occurrence with the aim of helping nursing managers to offer appropriate assistance for high-risk groups and build a caring climate and friendly environment for at-risk hospital nurses. One example of this is hospital nursing managers can assist nurses who are at risk for dysmenorrhea by setting up a more accommodating system of working hours that is based on nurses' menstrual cycle. Because women with dysmenorrhea do not necessarily need to take an entire day off of work, if a supportive environment can be created that allows them to take a break from work to apply hot compresses or drink hot beverages, their comfort will be improved. Work environments, particularly hospitals, can provide measures such as flexible menstrual leave or scheduling menstrual leave, thereby integrating menstrual self-care into annual occupational safety and health training. In addition, spaces can be provided at work for women to take short breaks and supportive measures can be offered, such as thermotherapy, aromatherapy massage, and others. Such measures will enable nurses to care for themselves at work, thus improving their workplace comfort and increasing their work satisfaction and performance.

ACKNOWLEDGEMENTS

The authors thank the participants for their participation and the nurse managers of the two hospitals for their administrative assistance. We also thank Editage for their professional English review of this manuscript.

Competing interests

None.

Funding

This research received no specific grants from any funding agency in the public, commercial, or not-for-profit sectors.

Contributors

Min-Hui Chiu and Hsiu-Hung Wang designed the study, analyzed and interpreted the data, and drafted the manuscript. Yi-Hsin Yang assisted in defining the statistical analysis. Min-Hui Chiu and Su-Chen Hsu contributed to the data collection and analysis. Hsiu-Fen Hsieh and Huei-Mein Chen assisted to draft the manuscript. All authors had full access to all data in the study and take responsibility for its integrity.

Data sharing statement

No additional data are available.

REFERENCES

1. Chen CX, Kwekkeboom KL, Ward SE. Beliefs about dysmenorrhea and their relationship to self-management. *Res Nurs Health* 2016;39(4):263-76.
2. Eryilmaz G, Ozdemir F. Evaluation of menstrual pain management approaches by northeastern Anatolian adolescents. *Pain management nursing: official journal of the American Society of Pain Management Nurses* 2009;10(1):40-7.
3. Kato T. Effects of flexibility in coping with menstrual pain on depressive symptoms. *Pain practice: the official journal of World Institute of Pain* 2017;17(1):70-77.
4. Zhu X, Wong F, Bensoussan A, Lo SK, Zhou C, Yu J. Are there any cross-ethnic differences in menstrual profiles? A pilot comparative study on Australian and Chinese women with primary dysmenorrhea. *The journal of obstetrics and gynaecology research* 2010;36(5):1093-101.
5. Abdul-Razzak KK, Ayoub NM, Abu-Taleb AA, Obeidat BA. Influence of dietary intake of dairy products on dysmenorrhea. *The journal of obstetrics and gynaecology research* 2010;36(2):377-83.
6. Agarwal AK, Agarwal A. A study of dysmenorrhea during menstruation in adolescent girls. *Indian journal of community medicine : official publication of Indian Association of Preventive & Social Medicine* 2010;35(1):159-64.
7. Loto OM, Adewumi TA, Adewuya AO. Prevalence and correlates of dysmenorrhea among Nigerian college women. *The Australian & New Zealand journal of obstetrics & gynaecology* 2008;48(4):442-4.
8. Chang YT, Chen YC. Menstrual health care behavior and associated factors among female elementary students in the Hualien region. *Journal of Nursing Research* 2008;16(1):11-6.
9. Chiou MH, Wang HH. Predictors of dysmenorrhea and self-care behavior among vocational nursing school female students. *Journal of Nursing Research* 2008;16(1):17-25.
10. Cheng HF, Lin YH. Selection and efficacy of self-management strategies for dysmenorrhea in young Taiwanese women. *Journal of clinical nursing* 2011;20:1018-25.
11. Palmer D. "To help a million sick, you must kill a few nurses": Nurses' occupational health, 1890–1914. *Nursing History Review* 2012;20(1):14-45.
12. Laschinger HK, Grau AL. The influence of personal dispositional factors and organizational resources on workplace violence, burnout, and health outcomes in new graduate nurses: a cross-sectional study. *International journal of nursing studies* 2012;49(3):282-91.
13. Payne LA, Rapkin AJ, Lung KC, Seidman LC, Zeltzer LK, Tsao JC. Pain catastrophizing predicts menstrual pain ratings in adolescent girls with chronic pain. *Pain Med* 2016;17(1):16-24.

14. Berkley KJ, McAllister SL. Don't dismiss dysmenorrhea! *Pain* 2011;152(9):1940-1.
15. Chiu MH, Wang HH, Hsu SC, Liu IP. Dysmenorrhoea and self-care behaviours among hospital nurses: a questionnaire survey. *Journal of clinical nursing* 2013;22(22):3130-40.
16. Wong LP. Attitudes towards dysmenorrhoea, impact and treatment seeking among adolescent girls: a rural school-based survey. *The Australian journal of rural health* 2011;19(4):218-23.
17. Zukr SM, Naing L, Hamzah TNT, Hussain NHN. Primary dysmenorrhea among medical and dental university students in Kelantan: Prevalence and associated factors. *International Medical Journal* 2009;16(2):93-99.
18. Lu I-C. Dysmenorrhea and related factors in Taiwanese adolescent girls. *University of Texas at Austin* 2010.
19. Chang C, Chen FL, Chang CH, Hsu CH. A preliminary study on menstrual health and menstrual leave in the workplace in Taiwan. *Taiwan J Public Health* 2011;30(5):436-50.
20. Schluter PJ, Turner C, Huntington AD, Bain CJ, McClure RJ. Work/life balance and health: the nurses and midwives e-cohort study. *International Nursing Review* 2011(58):28-36.
21. Cady RK, Diamond ML, Diamond MP, Ballard JE, Lener ME, Dorner DP, et al. Sumatriptan-naproxen sodium for menstrual migraine and dysmenorrhea: satisfaction, productivity, and functional disability outcomes. *Headache* 2011;51(5):664-73.
22. Bettendorf B, Shay S, Tu F. Dysmenorrhea: contemporary perspectives. *Obstetrical and Gynecological Survey* 2008;63(9):597-603.
23. Perry M. Treatment options for dysmenorrhoea. *Practice Nursing* 2012;23(4):4.
24. Ju H, Jones M, Mishra GD. Smoking and trajectories of dysmenorrhoea among young Australian women. *Tobacco control* 2016;25(2):195-202.
25. Chiou MH, Wang HH. The relationship between dysmenorrhea and menstrual attitudes among female students in vocational nursing schools. *The Journal of Nursing* 2004;51(4):45-52.
26. Parker MA, Sneddon AE, Arbon P. The menstrual disorder of teenagers (MDOT) study: determining typical menstrual patterns and menstrual disturbance in a large population-based study of Australian teenagers. *BJOG : an international journal of obstetrics and gynaecology* 2010;117(2):185-92.
27. Stubbs ML. Cultural perceptions and practices around menarche and adolescent menstruation in the United States. *Annals of the New York Academy of Sciences* 2008;1135:58-66.

- 1
 - 2
 - 3
 - 4
 - 5
 - 6
 - 7
 - 8
 - 9
 - 10
 - 11
 - 12
 - 13
 - 14
 - 15
 - 16
 - 17
 - 18
 - 19
 - 20
 - 21
 - 22
 - 23
 - 24
 - 25
 - 26
 - 27
 - 28
 - 29
 - 30
 - 31
 - 32
 - 33
 - 34
 - 35
 - 36
 - 37
 - 38
 - 39
 - 40
 - 41
 - 42
 - 43
 - 44
 - 45
 - 46
 - 47
 - 48
 - 49
 - 50
 - 51
 - 52
 - 53
 - 54
 - 55
 - 56
 - 57
 - 58
 - 59
 - 60
28. Cheng CY, Yang K, Liou SR. Taiwanese adolescents? Gender differences in knowledge and attitudes towards menstruation. *Nursing & Health Sciences* 2007;9(2):127-34.
29. H LY. The medical utilization of chinese and western medicine in females with dysmenorrhea. Kaohsiung Medical University, 2012.
30. Chiou M-H, Wang H-H, Yang Y-H. Effect of systematic menstrual health education on dysmenorrheic female adolescents' knowledge, attitudes, and self-care behavior. *Kaohsiung J Med Sci* 2007;23(4):183-90.
31. Brooks-Gunn J, Ruble D. The menstrual attitude questionnaire. *Psychosomatic medicine* 1980;42:503-11.
32. S L. Prevalence and associated factors of premenstrual syndrome among nurses in Taiwan. In: Ministry of Science and Technology, R. O. C., editor, 1994.
33. Morrison LA, Larkspur L, Calibuso MJ, Brown S. Women's attitudes about menstruation and associated health and behavioral characteristics. *American Journal of Health Behavior* 2010;34(1):90-100.
34. Lo Sing-Kai, Lin Ching-Fen. On the use and interpretation of Cronbach's alpha. *The Journal of Nursing Research* 1998;6:82-9.
35. Lin YX. *When "Experts" meet the medical professionals: The women with menstrual pain encountering western medicine*. Kaohsiung Medical University, Taiwan, 2006.
36. Chung F, Yao C, Wan G. The associations between menstrual function and life style/working conditions among nurses in Taiwan. *Journal of occupational health* 2005;47(2):149-56.
37. Takeuchi H, Oishi T, Harada T. Morningness-eveningness preference, and mental and physical symptoms during the menstrual cycle of Japanese junior high school students. *Sleep and Biological Rhythms* 2003(1):245-47.
38. Firat MZ, Kulakac O, Oncel S, Akcan A. Menstrual attitude questionnaire: confirmatory and exploratory factor analysis with Turkish samples. *Journal of advanced nursing* 2009;65(3):652-62.

STROBE Statement—checklist of items that should be included in reports of observational studies

	Item No	Page No	Recommendation
Title and abstract	1	1	(a) Indicate the study's design with a commonly used term in the title or the abstract
		2	(b) Provide in the abstract an informative and balanced summary of what was done and what was found
Introduction			
Background/rationale	2	4-8	Explain the scientific background and rationale for the investigation being reported
Objectives	3	4	State specific objectives, including any prespecified hypotheses
Methods			
Study design	4	9	Present key elements of study design early in the paper
Setting	5	9	Describe the setting, locations, and relevant dates, including periods of recruitment, exposure, follow-up, and data collection
Participants	6	9	<i>Cross-sectional study</i> —Give the eligibility criteria, and the sources and methods of selection of participants
Variables	7	10-12	Clearly define all outcomes, exposures, predictors, potential confounders, and effect modifiers. Give diagnostic criteria, if applicable
Data sources/ measurement	8*	10-12	For each variable of interest, give sources of data and details of methods of assessment (measurement). Describe comparability of assessment methods if there is more than one group
Bias	9	3	Describe any efforts to address potential sources of bias
Study size	10	9	Explain how the study size was arrived at
Quantitative variables	11	12	Explain how quantitative variables were handled in the analyses. If applicable, describe which groupings were chosen and why
Statistical methods	12	12	(a) Describe all statistical methods, including those used to control for confounding
			(b) Describe any methods used to examine subgroups and interactions
			(c) Explain how missing data were addressed
			(d) <i>Cohort study</i> —If applicable, explain how loss to follow-up was addressed
			<i>Case-control study</i> —If applicable, explain how matching of cases and controls was addressed
			<i>Cross-sectional study</i> —If applicable, describe analytical methods taking account of sampling strategy
			(e) Describe any sensitivity analyses

Continued on next page

Results			
Participants	13*	9	(a) Report numbers of individuals at each stage of study—eg numbers potentially eligible, examined for eligibility, confirmed eligible, included in the study, completing follow-up, and analysed (b) Give reasons for non-participation at each stage
Descriptive data	14*	13	(a) Give characteristics of study participants (eg demographic, clinical, social) and information on exposures and potential confounders (b) Indicate number of participants with missing data for each variable of interest
Outcome data	15*	13	<i>Cross-sectional study</i> —Report numbers of outcome events or summary measures
Main results	16	13	(a) Give unadjusted estimates and, if applicable, confounder-adjusted estimates and their precision (eg, 95% confidence interval). Make clear which confounders were adjusted for and why they were included (b) Report category boundaries when continuous variables were categorized (c) If relevant, consider translating estimates of relative risk into absolute risk for a meaningful time period
Other analyses	17	-	Report other analyses done—eg analyses of subgroups and interactions, and sensitivity analyses
Discussion			
Key results	18	19-24	Summarise key results with reference to study objectives
Limitations	19	3	Discuss limitations of the study, taking into account sources of potential bias or imprecision. Discuss both direction and magnitude of any potential bias
Interpretation	20	19-24	Give a cautious overall interpretation of results considering objectives, limitations, multiplicity of analyses, results from similar studies, and other relevant evidence
Generalisability	21	19-24	Discuss the generalisability (external validity) of the study results
Other information			
Funding	22	26	Give the source of funding and the role of the funders for the present study and, if applicable, for the original study on which the present article is based

*Give information separately for cases and controls in case-control studies and, if applicable, for exposed and unexposed groups in cohort and cross-sectional studies.

Note: An Explanation and Elaboration article discusses each checklist item and gives methodological background and published examples of transparent reporting. The STROBE checklist is best used in conjunction with this article (freely available on the Web sites of PLoS Medicine at <http://www.plosmedicine.org/>, Annals of Internal Medicine at <http://www.annals.org/>, and Epidemiology at <http://www.epidem.com/>). Information on the STROBE Initiative is available at www.strobe-statement.org.

BMJ Open

Influencing factors of dysmenorrhea among hospital nurses: A questionnaire survey in Taiwan

Journal:	<i>BMJ Open</i>
Manuscript ID	bmjopen-2017-017615.R2
Article Type:	Research
Date Submitted by the Author:	15-Nov-2017
Complete List of Authors:	Chiu, Min-Hui; Kaohsiung Medical University College of Nursing, Doctoral Student; Chi Mei Medical Center, Liouying, Department of Nursing Hsieh, Hsiu-Fen; Kaohsiung Medical University, College of Nursing Yang, Yi-Hsin; Kaohsiung Med Univ Chen, Huei-Mein; Chung Hwa University of Medical Technology Hsu, Su-Chen; Chi Mei Medical Center Wang, Hsiu-Hung; Kaohsiung Medical University, College of Nursing
Primary Subject Heading:	Communication
Secondary Subject Heading:	Intensive care
Keywords:	dysmenorrhea, prevalence rate, self-care behavior, hospital nurses

SCHOLARONE™
Manuscripts

Title Page

Manuscript title

Influencing factors of dysmenorrhea among hospital nurses: A questionnaire survey in Taiwan

Running head

Dysmenorrhea and influencing factors

Authors

Min-Hui Chiu

1. Doctoral Student, College of Nursing, Kaohsiung Medical University, Taiwan
Address: No. 100 Shih-Chuan 1st Road, San-Ming District, Kaohsiung 80708, Taiwan
2. Director, Chi Mei Medical Center, Liouying, Tainan, Taiwan
Address: No.201, Taikang, Liouying Dist., Tainan City 736, Taiwan
Email: 300015@mail.chimei.org.tw

Hsiu-Fen Hsieh

Associate Professor, College of Nursing, Kaohsiung Medical University, Taiwan
Address: No. 100 Shih-Chuan 1st Road, San-Ming District, Kaohsiung 80708, Taiwan
Email: hsiufen96@gmail.com

Yi-Hsin Yang

Professor, College of Pharmacy, Kaohsiung Medical University, Kaohsiung, Taiwan
Address: No. 100 Shih-Chuan 1st Road, San-Ming District, Kaohsiung 80708, Taiwan
E-mail: yihnya@kmu.edu.tw
Telephone: 886-7-3121101 ext. 2262

Huei-Mein Chen

Associate Professor, Chung Hwa University of Medical Technology, Tainan, Taiwan
Address: No. 89, Wen-hua1st St, Rende District, Tainan 71703, Taiwan
E-mail: hueimein@mail.hwai.edu.tw
Telephone: 886-6-2671214 ext. 512

Su-Chen Hsu

Chi Mei Medical Center, Tainan, Taiwan

1
2
3 Address: No.901, Zhonghua Rd., Yongkang Dist., Tainan City 710, Taiwan

4 E-mail: 300001@mail.chimei.org.tw

5 Telephone: 886-6-2812811 ext. 53000

6 Fax: 886-6-2827480

7
8
9
10 **Hsiu-Hung Wang***

11 Professor, College of Nursing, Kaohsiung Medical University, Taiwan

12
13
14 **Corresponding author**

15 Hsiu-Hung Wang, RN, FAAN, PhD

16 Dean and Professor, College of Nursing, Kaohsiung Medical University

17 No. 100, Shih-Chuan 1st Road, Kaohsiung City 80708, Taiwan

18 Tel: 886-7-3121101 ext 2624

19 Fax: 886-7-3218364

20 E-mail: hhwang@kmu.edu.tw

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

keywords

Dysmenorrhea, prevalence rate, self-care behavior, hospital nurses

Word count

4005

For peer review only

Influencing factors of dysmenorrhea among hospital nurses: A questionnaire survey in Taiwan

ABSTRACT

Objectives: Nurses are at high risk for dysmenorrhea while working with patients.

The study objectives were to: (1) describe demographic and menstruation characteristics, knowledge about dysmenorrhea, and menstrual attitudes towards menstruation among dysmenorrheal and non-dysmenorrheal hospital nurses; (2) identify significant differences between the groups; and (3) examine the factors influencing dysmenorrhea.

Methods: This cross-sectional survey used a structured questionnaire, administered at two hospitals in southern Taiwan. Participants included hospital nurses at least 18 years of age who agreed to participate. All participants were recruited through random sampling. The questionnaire included demographic data, the Dysmenorrhea Knowledge Scale, and Menstrual Attitude Scale (MAS).

Results: A total of 420 nurses completed the questionnaire. Among them, 297 (70.7%) had experienced dysmenorrhea in the past 6 months and 123 (29.3%) had not. Significant differences in age ($p < 0.001$), marital status ($p < 0.001$), childbearing status ($p < 0.001$), age at menarche ($p < 0.05$), and rotating three-shift ratio ($p < 0.05$) were identified between the dysmenorrhea and non-dysmenorrhea groups. Analysis of the MAS results revealed significant differences between the groups regarding consideration of menstruation as a debilitating ($p < 0.001$) or bothersome event ($p < 0.05$), anticipation and prediction of menstruation onset ($p < 0.01$), and denial of any effects from menstruation ($p < 0.001$). Results of the multiple logistic regression

1
2
3 showed that predictive factors included age less than 40 years (4.46 vs. 1.00), working
4 three-shift rotations (2.07 vs. 1.00), marital status (2.59 vs. 1.00), acknowledging
5 menstruation as a debilitating event (2.72 vs. 1.00), and denial of effects from
6 menstruation (2.59 vs. 1.00).
7
8
9
10

11 **Conclusions:** These findings could help nursing managers to create a caring climate
12 and friendly work environment for hospital nurses at risk for dysmenorrhea.
13
14

15 **Keywords:** Dysmenorrhea, prevalence rate, self-care behavior, hospital nurses
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

Strengths and limitations of this study

- The factors influencing dysmenorrhea are complex; in this study, multiple logistic regression analysis was used to control for confounding variables and to identify these factors.
- Participants in this study were aged 22–48 years; therefore, the results have limited application to hospital nurses aged under 22 years and above 48 years.
- This study was limited in terms of analyzing the effects of lifestyle choices, such as smoking and alcohol consumption, because only a few nurses reported those behaviors.

INTRODUCTION

More than 50% of women experience dysmenorrhea, defined as pain that accompanies menstruation. The majority (75.1%) of women believe dysmenorrhea to be a normal part of their lives and that the symptoms will continue to affect their daily life until they near menopause.¹ Dysmenorrhea occurs during the first 1 to 3 years after menarche and is commonly accompanied by sweating, lack of appetite, headache, distraction, nausea, vomiting, dizziness,² and depression.³ Dysmenorrhea arises during menstrual bleeding owing to lower abdominal pain brought on by menses, and is not any other gynecological disorders.⁴ The prevalence rates of dysmenorrhea differ by age; however, over 50% of women in all age groups experience dysmenorrhea. The prevalence rate of dysmenorrhea worldwide is between 50.9% and 87.4%.⁵⁻⁷ Parker et al. found that 21% of Australian adolescents experienced severe dysmenorrhea, with 26% requiring absence from school due to dysmenorrhea.⁸

In Taiwan, the reported prevalence rate of dysmenorrhea among adolescent vocational nursing students aged 16–18 years was 73.3%;⁹ another study reported a prevalence rate of 90.2% among the general population of women in Taiwan.¹⁰ A study on hospital nurses in Taiwan reported a prevalence rate of 90.7%.¹¹ Despite the high prevalence of dysmenorrhea, nurses often neglect their own health while working and caring for patients.¹²

Dysmenorrhea can be cyclical or chronic; in some cases, it can be a possible result of pain catastrophizing and anxiety sensitivity.¹³ Dysmenorrhea not only causes physical pain, it also affects mental well-being and quality of life, leading to work or school absenteeism and significant health burdens.¹⁴⁻¹⁸ Despite its high prevalence rate and effect on daily life, 76.1% of women still believe that dysmenorrhea is a natural part of a women's menstrual cycle, and only 14.8% believe that treatment is

1
2
3 necessary.¹⁶ In terms of dysmenorrhea information, 62.3% of adolescent girls claim
4 that they learned about the condition from their mothers, and 52.9% state that they
5 obtained their knowledge from peers.¹⁶ In addition, dysmenorrhea is negatively
6 correlated with self-awareness of one's own health; in other words, individuals with
7 dysmenorrhea have poorer self-awareness regarding their own health.⁶

8
9
10
11
12
13
14 A study of 297 nurses (average age 30.3 years) revealed that 8.4% had taken
15 menstrual leave, 11.8% had taken a leave of absence for dysmenorrhea, and 28.6%
16 saw their doctors owing to the condition.¹⁵ Furthermore, 20.8% of adolescent girls
17 from vocational schools in the same region had visited doctors for consultation
18 regarding dysmenorrhea.⁹ Menstrual leave is currently available in Japan, Indonesia,
19 South Korea, and Taiwan. Focus group studies in Taiwan have shown that most
20 women have experienced menstrual discomfort and that they understand the
21 implementation and regulation of menstrual leave in different companies. However, in
22 reality, menstrual leave is rarely availed of owing to the following factors: "the
23 regulations are too rigid," "no one else has taken menstrual leave," "other kinds of
24 leave are available," "no one will take my shift," or "there has to be a doctor's note."¹⁹

25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
Previous studies have shown that the physical and mental health of nurses is
significantly correlated with job satisfaction, tiredness, and comfort.^{19, 20} The Gender
Equality in Employment Act of Taiwan, approved in 2002, clearly defines the
regulations regarding menstrual leave, in which a female employee who has trouble
working due to discomfort during menstruation should take 1 day of menstrual leave
every month.²¹ However, because of the particularities of nursing, temporary leave
affects the allocation of nurses. Cheng found that the long-term use of medication to
maintain a ready-to-work status is common among women. Nurses, who have
frequent contact with drugs, use them extensively to overcome menstrual discomfort

1
2
3 at work,¹⁰ including the use of sumatriptan–naproxen sodium to reduce the effects of
4
5 dysmenorrhea during work and leisure.^{21, 22}
6

7
8 Several risk factors affect dysmenorrhea. (a) Age: older women are generally less
9
10 likely to experience dysmenorrhea, although the relationships with marital status or
11
12 childbirth history remain unclear. (b) Age at menarche: age at menarche is
13
14 significantly related to the occurrence of dysmenorrhea, and women who demonstrate
15
16 an earlier age at menarche are more likely to experience more severe dysmenorrhea.^{7,}
17
18 ¹⁷ (c) Duration of menstrual cycle: a Nigerian study noted that longer menstrual flow
19
20 duration is an important predictive factor of dysmenorrhea.⁷ (d) Regularity of
21
22 menstrual cycle: women with irregular menstrual cycles are more likely to have
23
24 dysmenorrhea.⁹ In terms of lifestyle factors, the results of several studies indicate that
25
26 smoking and alcohol consumption are risk factors for primary dysmenorrhea and that
27
28 women who habitually smoke or drink alcohol are more likely to have
29
30 dysmenorrhea.^{23, 24} With regards to work, several studies have found that women who
31
32 work rotations, especially nurses who have to alternate between day and night shifts,
33
34 experience relatively more serious menstrual discomfort.²¹ Nurses who work night
35
36 shifts are more likely to experience dysmenorrheal symptoms during their menstrual
37
38 cycles than those who work regular shifts.¹⁵
39
40

41
42 Furthermore, the attitudes of adolescent students toward menstruation correlate
43
44 with the occurrence of dysmenorrhea.²⁵ Adolescent girls who tend to feel that
45
46 menstruation is a debilitating experience and those who anticipate and predict the
47
48 onset of menstruation are more likely to experience dysmenorrhea.²⁶ The effect of
49
50 dysmenorrhea on women varies based on the degree and frequency of their pain.
51

52
53 Previous studies have revealed that women have inadequate knowledge about
54
55 menstruation and that educational information about menstrual health care contains
56

1
2
3 contradictory messages, including the idea that menstruation is a normal and natural
4 event but should be “invisible.”²⁷ Cheng observed that some female students knew
5 about menstruation taboos; although they did have doubts about those taboos, they
6 still had negative attitudes towards menstruation.²⁸ Medical expenses due to
7 dysmenorrhea increase with age; therefore, the topic of women’s menstrual health and
8 accompanying emotional challenges should be taken seriously.²⁹ Nevertheless,
9 because dysmenorrhea is so common, many women consider it a natural phenomenon
10 and rarely seek medical treatment owing to their cultural and religious attitudes¹⁵ or
11 because they feel that they can tolerate the pain.²³

12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

Herein, we defined dysmenorrhea as lower abdominal pain brought on by menses at any time during the past 6 months.¹¹ In Taiwan, 98.8% of hospital nurses are women. There is a high prevalence rate of dysmenorrhea among women; owing to the nature of nursing work, nurses are at high risk for dysmenorrhea. However, previous relevant studies on nurses have not discussed the factors influencing dysmenorrhea. The objectives of this study were to: (1) describe the basic demographic data as well as menstruation characteristics, knowledge about dysmenorrhea, and attitudes towards menstruation among dysmenorrheal and non-dysmenorrheal clinical nurses; (2) analyze the demographic data as well as menstruation characteristics, knowledge about dysmenorrhea, and attitudes towards menstruation among dysmenorrheal and non-dysmenorrheal clinical nurses, and identify significant differences between these groups; and (3) examine the risk factors for dysmenorrhea among clinical nurses.

METHODS

Study design and participants

1
2
3 This study used a cross-sectional survey based on structured questionnaires for
4 data collection. Participants were recruited from two medical institutions in southern
5 Taiwan; one institution was a medical center with 1,300 beds, the other was a regional
6 hospital with 650 beds. The two institutions were at different hospital levels within
7 the same health care system. All 2,000 nurses at the two institutions were potential
8 participants. The inclusion criteria for this study were full-time nurses employed at
9 the study hospitals who were at least 18 years of age and agreed to participate in the
10 study after providing informed consent. The sample size was estimated using JMP 7.0
11 (SAS Institute, Cary, NC, USA) with an effect size of 0.3, α of 0.05, and a power of
12 0.80, resulting in a required sample size of 350 individuals. With a predicted
13 questionnaire recovery rate of 80%, the sample size was estimated to be 438. After
14 obtaining a list of the names of the potential participants, serial numbers were
15 assigned to each participant. The estimated number of study participants were
16 randomly selected using the sampling function of Microsoft Excel 2007. In total, 450
17 nurses were chosen for inclusion in the study, and finally, 420 valid questionnaires
18 were returned, yielding a response rate of 93.3%. Among these 420 nurses, 297
19 (70.7%) had experienced dysmenorrhea in the past 6 months and were classified as
20 the dysmenorrhea group; 123 (29.3%) nurses who had not experienced dysmenorrhea
21 were classified as the non-dysmenorrhea group.

22
23
24 The random sampling was conducted after the study received approval from the
25 Institutional Review Boards (IRBD No. CLH0088) of the two hospitals. The
26 questionnaires were administered to the nurses by a trained research assistant. All
27 participants provided their written consent and answered the questionnaires
28 independently. The written consent form indicated that participation in this study was
29 not compulsory and the questionnaire was anonymous. Anyone who was not willing
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

1
2
3 to participate in the study could simply choose not to return the questionnaire. No
4 individual rights or interests were compromised during this process. Those who
5 completed the questionnaire received a small gift. The questionnaires were collected 2
6 weeks after being issued, upon which the written consent forms and questionnaires
7 were filed separately and placed in locations chosen by participants, to protect their
8 personal privacy.
9
10
11
12
13
14

15 16 17 18 **Measurements**

19
20 This study was conducted in May 2010. The study instruments included the
21 Demographic Inventory (DI), Dysmenorrheal Knowledge Scale (DKS), and
22 Menstrual Attitude Scale (MAS). The DI was designed based on related studies^{9, 30}
23 and included age, age at menarche, marital and childbirth status, discussion of
24 menstrual hygiene, dysmenorrhea experience, medical experience of dysmenorrhea,
25 working conditions, self-care health education regarding menstruation, factors
26 affecting dysmenorrhea, and work coping methods. The DKS was designed using
27 both the researchers' clinical experiences and previous studies.³⁰ The scale included
28 20 questions with yes/no answers, for example, "Women with dysmenorrhea are at
29 high risk for sterility" and "Applying hot compresses to the lower abdomen can ease
30 menstrual cramps". Correct answers received a score of 1, and incorrect answers
31 received a score of 0. The total summed scores ranged from 0 to 20, with higher
32 scores representing better knowledge of dysmenorrhea. The Kuder-Richardson
33 reliability coefficient was 0.63, according to a previous study³⁰ and 0.72 in this study.
34 The MAS was developed by Brooks-Gunn and Ruble (1980)³¹ and modified by Lee
35 (1994). Lee translated this scale into a Chinese version and deleted one item owing to
36 cultural differences.³² The Chinese version of the MAS includes 32 items with five
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

1
2
3 dimensions, including consideration of menstruation as a debilitating event,
4 menstruation as a bothersome event, menstruation as a natural event, anticipation and
5 prediction of the onset of menstruation, and denial of the effects of menstruation.
6
7 Each item is scored on a 7-point Likert scale from 1 (strongly disagree) to 7 (strongly
8 agree). The total summed scores range from 32 to 224, with higher scores
9 representing a more positive attitude toward menstruation. Cronbach's α was 0.58 in
10 the original scale³¹ and 0.61 in Morrison's study.³³ Lo and Lin (1998) suggested that
11 an acceptable Cronbach's α ranged from 0.3 to 0.7.³⁴ Because we were concerned
12 about the comparability of our study with other studies, we decided to use this scale
13 with a Cronbach's α of 0.57 in the present study.
14
15
16
17
18
19
20
21
22
23
24
25

26 Data analysis

27
28 The data in this study were entered using Microsoft Excel. JMP 7.0 was used for
29 the statistical analysis. Based on the study variables and objectives, percentage, mean,
30 and standard deviation were used to describe participant demographics and
31 menstruation characteristics as well as DKS and MAS score distributions in the
32 dysmenorrhea and non-dysmenorrhea groups. We used *t*-tests and chi-square tests to
33 detect differences in demographics, menstruation characteristics, DKS score, and
34 MAS score between the two groups. Finally, multiple logistic regression analysis was
35 used to examine factors (independent variables) influencing the occurrence of
36 dysmenorrhea (dependent variable). Independent variables with significant
37 differences were included in the multiple logistic regression analysis.
38
39
40
41
42
43
44
45
46
47
48
49
50
51

52 RESULTS

53
54 A total of 420 participants completed the questionnaire. Among them, 297
55
56
57

1
2
3 (70.7%) had experienced dysmenorrhea in the past 6 months, and 123 (29.3%) had
4
5 not. Compared with the non-dysmenorrhea group, participants with dysmenorrhea
6
7 were significantly younger ($t = -3.78$, $p < 0.001$), were more frequently unmarried
8
9 (77.78%, $\chi^2 = 20.03$, $p < 0.001$) and without a history of childbirth (83.16%, $\chi^2 =$
10
11 19.38, $p < 0.001$), and more often demonstrated an age at menarche of less than 12
12
13 years (15.49%, $\chi^2 = 4.70$, $p = 0.03$); moreover, a higher percentage of participants
14
15 with dysmenorrhea worked a three-shift rotation (91.25%, $\chi^2 = 6.06$, $p = 0.014$). The
16
17 other variables showed no significant differences between the two groups (Table 1).
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

Table 1. Comparison of demographic and menstruation characteristics between the dysmenorrhea and non-dysmenorrhea groups

Variables	Dysmenorrheal (n = 297)		Non-dysmenorrheal (n = 123)		t or χ^2	p-value
	n	%	n	%		
Age (years) M± SD	30.33± 3.61		32.04±4.47		-3.78	<0.001
Age<40	292	98.32	115	93.50	6.74	<0.01
Age≥40	5	1.68	8	6.50		
Marital status					20.03	<0.001
Unmarried	231	77.78	69	56.10		
Married	66	22.22	54	43.90		
Childbearing					19.38	<0.001
No	247	83.16	78	63.41		
Yes	50	16.84	45	36.59		
Age of menarche					4.70	0.030
<12	46	15.49	11	8.94		
≥12	251	84.51	112	91.06		
Regularity of menstrual cycle					0.10	0.755
Yes	193	64.98	85	69.11		
No	104	35.02	38	30.89		
Duration of menstrual cycle					1.99	0.575
≤4 days	110	37.04	43	34.96		
5-6 days	157	52.86	70	56.91		
≥7 days	30	10.10	10	8.13		
Prior health education on dysmenorrhea					0.99	0.321
No	163	54.88	74	60.16		
Yes	134	45.12	49	39.84		
Three-shift rotation					6.06	0.014
Yes	271	91.25	102	82.93		
No	26	8.75	21	17.07		

Table 1. Comparison of demographic and menstruation characteristics between dysmenorrhoeal and non-dysmenorrhoeal groups (continued)

Variables	Dysmenorrhoeal (n = 297)		Non-dysmenorrhoeal (n = 123)		t or χ^2	p-value
	n	%	n	%		
Whether to know it is allowed to ask for menstrual leave					2.03	0.154
Yes	242	81.48	107	86.99		
No	55	18.52	16	13.01		
Whether to consider menstrual leave necessary					1.68	0.195
Yes	288	96.97	115	93.50		
No	9	3.03	8	6.50		
Have ever asked for menstrual leave					0.14	0.707
No	272	91.58	114	92.68		
Yes	25	8.42	9	7.32		
Have ever asked for more leaves for dysmenorrhoea?					3.59	0.058
No	262	88.22	116	94.31		
Yes	35	11.78	7	5.69		
Secondary dysmenorrhoea					0.02	0.886
No	267	89.90	110	89.43		
Yes	30	10.10	13	10.57		

Note: Regularity of periods indicates that the duration between consecutive first-day menses is similar.

With respect to knowledge about dysmenorrhoea, there was a non-significant difference between the two groups ($p = 0.238$). The highest scoring items among both

groups were “applying hot compresses,” “adopting a knee-to-chest position,” and “drinking hot beverages including warm water or milk or ginger soup to ease cramps.” The lowest scoring items were “normal bleeding amount every cycle,” “dysmenorrhea is a disease of the reproductive system,” and “women with dysmenorrhea are at high risk for sterility.” Regarding attitudes towards menstruation, after standardizing, the highest scoring dimension among the dysmenorrhea group was “considering menstruation a debilitating event” (73.30) and “considering menstruation a natural event” (74.93) in the non-dysmenorrhea group. The lowest scoring dimension in both groups was “denial of any effect of menstruation” (47.18 in the dysmenorrhea group and 54.69 in the non-dysmenorrhea group). Women who experienced dysmenorrhea had significantly higher scores than women in the non-dysmenorrhea group with respect to the following: considering menstruation a debilitating event ($p < 0.001$), considering menstruation a bothersome event ($p = 0.024$), and anticipation and prediction of menstruation onset ($p = 0.004$). In terms of denying the effects of menstruation, scores for the dysmenorrhea group were significantly lower than those of the non-dysmenorrhea group ($p < 0.001$). No significant differences were observed between the groups in terms of considering menstruation a natural event (Table 2).

Table 2. Comparison of dysmenorrheic knowledge and attitudes towards menstruation between the dysmenorrhea and non-dysmenorrhea groups

Variables	Dysmenorrheal (n = 297)		Non-dysmenorrheal (n = 123)		t-value	p-value
	M	SD	M	SD		
Dysmenorrheic knowledge scale	14.12	2.19	13.40	2.40	1.18	0.238
Menstrual attitude scale	148.6	12.74	146.85	12.69	1.31	0.192

	4						
Menstruation as a debilitating event	56.44	8.52	52.55	7.65	4.58	<0.001	
Menstruation as a bothersome event	29.05	5.01	27.54	6.64	2.27	0.024	
Menstruation as a natural event	19.98	5.19	20.98	4.92	-1.87	0.062	
Anticipation and prediction of the onset of menstruation	20.04	3.01	18.98	3.55	2.93	0.004	
Denial of any effect of menstruation	23.12	6.75	26.80	5.70	5.70	<0.001	

We conducted a multiple logistic regression analysis using dysmenorrhea and non-dysmenorrhea as dependent variables, which yielded significant differences in the variables of demographics, menstruation characteristics, DKS score, and MAS score between the two groups. The results of the analysis showed that crucial predictive factors for the occurrence of dysmenorrhea among hospital nurses included age less than 40 years, working three-shift rotations, and marital status. Moreover, using the mean as the contact point, we found that the tendency to consider menstruation to be a debilitating event and the tendency to deny the effects of menstruation were also predictive factors for dysmenorrhea occurrence. The adjusted odd ratios are shown in Table 3.

After controlling for working three-shift rotations, marital status, the tendency to consider menstruation a debilitating event, and the tendency to deny the effects of menstruation, the rate of occurrence of dysmenorrhea in women younger than 40 years was 4.46 times higher than that in women over 40 years of age. After controlling for age, marital status, the tendency to consider menstruation a debilitating

event, and tendency to deny the effects of menstruation, the rate of occurrence of dysmenorrhea among nurses who worked three-shift rotations was 2.07 times higher than that in nurses who did not. After controlling for age, working three-shift rotations, marital status, and the tendency to deny the effects of menstruation, participants who felt that menstruation was a debilitating event had a dysmenorrhea occurrence rate that was 2.72 times higher than that of participants who did not. After controlling for age, working three-shift rotations, marital status, and the tendency to think of menstruation as a debilitating event, women who did not acknowledge the effects of menstruation had a dysmenorrhea occurrence rate that was 2.59 times higher than that in women who did not deny these effects.

Table 3: Risk factors for dysmenorrhea

		Crude OR	95% CI	P-Value	Adjusted OR	95% CI	P-Value
Age	Age < 40	4.06	(1.33 , 13.7)	<0.05	4.46	(1.24 , 17.23)	<0.05
	Age ≥ 40	1.00	(- , -)		1.00	(- , -)	
Three-shift rotation	Yes	2.15	(1.15 , 3.98)	<0.05	2.07	(1.01 , 4.21)	<0.05
	No	1.00	(- , -)		1.00	(- , -)	
Married	No	2.74	(1.75 , 4.30)	<0.001	2.59	(1.57 , 4.28)	<0.001
	Yes	1.00	(- , -)		1.00	(- , -)	
Menstruation as a debilitating event	Higher	2.53	(1.63 , 3.95)	<0.001	2.72	(1.67 , 4.52)	<0.001
	Lower	1.00	(- , -)		1.00	(- , -)	

Denial of any effect of menstruation	Lower	2.83 (1.83 , 4.46)	<0.001	2.59 (1.61 , 4.23)	<0.001
	Higher	1.00 (- , -)		1.00 (- , -)	

Reference group: Age ≥ 40 years, no three-shift rotation, unmarried, reduced menstruation as a debilitating event, and increased denial of any effect on menstruation

CI, confidence interval; OR, odds ratio; SE, standard error.

DISCUSSION

Although the average age of participants in the dysmenorrhea group was significantly lower than that of women in the non-dysmenorrhea group, when age was the only variable being compared, the dysmenorrhea occurrence rate among nurses younger than 40 years was 4.06 times higher than that of nurses over age 40 years; after controlling for confounding variables, the rate of dysmenorrhea in younger women was 4.46 times higher than that in women over 40 years old. Therefore, age was a risk factor for dysmenorrhea. To our knowledge, no other studies have directly discussed the correlation between age and dysmenorrhea. However, previous studies have reported dysmenorrhea occurrence rates over 76% in adolescent girls or students,^{5,6,16} and another reported a rate of 55.9% among women aged 15–45 years,³¹ wherein a decrease in the occurrence of dysmenorrhea with increasing age was observed. As dysmenorrhea is very common, periodic dysmenorrhea is a natural phenomenon observed during the menstrual cycle. With age, the number of episodes of dysmenorrhea and the number of times coping strategies were used among women also increase; therefore, women with dysmenorrhea can be thought of as “female specialists with embodied knowledge,” meaning that their experiences may affect their subjective perception and tolerance of dysmenorrhea, which can in turn affect

1
2
3 the occurrence of dysmenorrhea.³⁵ Nursing is a female-dominated occupation in
4 Taiwan. In this study, 97% of participants were younger than 40 years old. The high
5 prevalence rate of dysmenorrhea severely affects nurses such that they may not be
6 able to focus on their work, thereby affecting the quality of patient care. Nurses may
7 need to take menstrual leave, which can cause a shortage of nursing resources.
8
9

10
11
12
13
14 In terms of marital status, the ratio of unmarried to married women in the
15 dysmenorrhea group was 2.74 times that of the non-dysmenorrhea group, indicating a
16 high dysmenorrhea occurrence rate among unmarried women; after adjusting for
17 confounding variables, it was 2.59 times higher than that among married women. This
18 result is consistent with those of Chung (average study participant age 27 years),
19 which reported a higher dysmenorrhea occurrence rate in single nurses than in
20 married and divorced ones.³⁶ Many women are told by their doctors that their
21 dysmenorrhea will improve after getting married and giving birth, which may
22 improve menstrual blood discharge. Furthermore, women may engage in positive
23 self-care for menstruation after getting married for childbearing reasons, which may
24 also be a factor that indirectly relieves dysmenorrhea.¹¹ In terms of work, the
25 dysmenorrhea occurrence rate among nurses in the dysmenorrhea group who worked
26 three-shift rotations was 2.15 times that of nurses in the non-dysmenorrhea group;
27 after adjusting for confounding variables, it was 2.07 times higher than among nurses
28 who did not work three-shift rotations. This indicates that work type (e.g., rotating
29 nightshifts) is a risk factor for dysmenorrhea in nurses. However, Chung found that
30 rotating work shifts were not significantly related to dysmenorrhea occurrence in
31 nurses.³⁶ Furthermore, one study among Japanese junior high school students also
32 found that nightshift work was not significantly associated with dysmenorrhea.³⁷
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
However, focus group interviews conducted by Chang revealed that working rotating

1
2
3 shifts was highly associated with menstrual discomfort.¹⁹ Owing to the nature of
4 nursing work, most nurses need to work rotating night shifts, and the graveyard shift
5 can easily cause uncomfortable menstrual cycles or obvious irregularities.²⁵ Owing to
6 nurses' day/night activities and sleep patterns, along with the increased pressure of
7 working night shifts, menstrual discomfort may be more common in those with
8 frequent rotation changes.
9

10
11
12
13
14
15
16 In terms of attitudes towards menstruation, the dysmenorrhea group had
17 significantly higher scores than the non-dysmenorrhea group in terms of considering
18 menstruation to be a debilitating event, and significantly lower scores than the
19 non-dysmenorrhea group in terms of denying the effects of menstruation. In the
20 multiple logistic regression analysis of attitudes towards menstruation, the tendency to
21 consider menstruation as a debilitating event in the dysmenorrhea group was 2.53
22 times higher than that in the non-dysmenorrhea group; after adjusting for confounding
23 variables, women with dysmenorrhea had a 2.72 times higher tendency to think of
24 menstruation as a debilitating event than women who did not consider menstruation to
25 be debilitating. The tendency to deny the effects of menstruation in the dysmenorrhea
26 group was 2.83 times higher than in the non-dysmenorrhea group; after adjusting for
27 confounding variables, women with dysmenorrhea had a 2.59 times higher tendency
28 to deny the effects of menstruation. In this study, attitude towards menstruation was
29 related to dysmenorrhea occurrence. Firat found no significant correlation between
30 menstrual discomfort and attitude towards menstruation.³⁸ In that study,
31 "menstruation is a debilitating event" and "anticipation and prediction of the onset of
32 menstruation" obtained significantly higher scores from the vocational nursing
33 students with dysmenorrhea than those who did not have dysmenorrhea. In terms of
34 "menstruation is a bothersome event" and "menstruation is a natural event," the
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

1
2
3 non-dysmenorrhea group demonstrated significantly higher scores than the
4
5 dysmenorrhea group; no significant differences were observed between the two
6
7 groups in terms of “denial of any effect of menstruation”.²⁶ These results are different
8
9 from those of the present study. The most obvious difference between the two studies
10
11 is the tendency to deny the effects of menstruation. Scores in the dysmenorrhea and
12
13 non-dysmenorrhea groups in the present study were 23.1 and 26.8, respectively,
14
15 compared with 26.6 and 28.0 among adolescent nursing students in Firat’s study. In
16
17 other words, compared with adolescent nursing students, the nurses in the present
18
19 study did not deny the effects caused by menstruation and recognized that they were
20
21 affected by menstruation. When a nurse experiences dysmenorrhea, her work may be
22
23 affected. If she requests for menstrual leave, temporary work arrangements must be
24
25 made and patient care may be affected. Because dysmenorrhea is a periodic
26
27 phenomenon, taking frequent menstrual leave may affect working relationships;
28
29 therefore, nurses usually choose to work even when they experience dysmenorrhea.
30
31
32

33 In summary, the factors influencing dysmenorrhea occurrence included age
34
35 under 40 years, working three-shift rotations, and marital status, and when mean was
36
37 used as the contact point, the tendency to consider menstruation to be a debilitating
38
39 event and tendency to deny the effects of menstruation. Chiou’s study of adolescent
40
41 nursing students revealed that menstrual regularity and health education were
42
43 predictive factors for dysmenorrhea.³⁰ Women who had irregular menstrual cycles and
44
45 had not received dysmenorrhea-related health education demonstrated higher
46
47 dysmenorrhea occurrence rates;⁹ however, these two factors were not predictive
48
49 factors in the present study. Our participants were nurses from a wide range of age
50
51 groups rather than only adolescents; therefore, they had more experience with
52
53 dysmenorrhea and self-care. In addition, menstrual health is a part of nursing
54
55
56
57
58
59
60

1
2
3 education; therefore, although nurses are not directly involved in health education,
4 education is a part of their professional training. Thus, health education was not an
5 important predictive factor in this study. This study mainly evaluated the factors
6 influencing dysmenorrhea in hospital nurses in order to help institutions build a caring
7 and friendly work environment for high-risk nurses. The participants were randomly
8 selected from only two hospitals in southern Taiwan; therefore, the results might have
9 limited generalizability to nurses in other parts of the country. In future studies, the
10 inclusion criteria should be extended to examine any differences across regions in
11 Taiwan. In addition, rotating night shifts are required in many other occupations
12 besides nursing. Considering different professional backgrounds and life experiences,
13 future studies should also include women from different occupations who work
14 rotating shifts, to evaluate the differences in the influence of their knowledge and
15 attitudes on the prevalence of dysmenorrhea. The variables assessed in this study
16 included demographic characteristics, menstruation history, dysmenorrhea-related
17 knowledge, and attitudes towards menstruation; however, lifestyle choices such as
18 smoking, alcohol consumption, and exercising were not included. Further studies that
19 analyze the effects of these lifestyle choices on dysmenorrhea are warranted.
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41

42 CONCLUSIONS

43
44 We assessed the various factors influencing the occurrence of dysmenorrhea with
45 the aim of helping nursing managers to offer appropriate assistance for high-risk
46 groups and build a caring and friendly work environment for at-risk hospital nurses,
47 which were as follows: (a) setting up a more accommodating working hours system
48 that is based on the nurses' menstrual cycle; (b) creating a supportive workplace
49 environment that allows nurses to take a break from work to apply hot compresses or
50
51
52
53
54
55
56
57
58
59
60

1
2
3 drink hot beverages; (c) allowing flexible menstrual leave or scheduling menstrual
4 leave, thereby integrating menstrual self-care into the annual occupational safety and
5 health training; (d) providing spaces at work for nurses to take short breaks and
6 offering supportive measures, such as thermotherapy, aromatherapy massage, and
7 others. Such measures will enable nurses to care for themselves at work, thus
8 improving their workplace comfort levels and increasing their work satisfaction and
9 performance.
10
11
12
13
14
15
16
17
18
19

20 **ACKNOWLEDGEMENTS**

21
22 The authors thank the participants for their participation and the nurse managers of
23 the two hospitals for their administrative assistance. We also thank Editage for their
24 professional English review of this manuscript.
25
26
27

28 **Competing interests**

29
30 None.
31
32

33 **Funding**

34
35 This research received no specific grants from any funding agency in the public,
36 commercial, or not-for-profit sectors.
37
38

39 **Contributors**

40
41 Min-Hui Chiu and Hsiu-Hung Wang designed the study, analyzed and interpreted the
42 data, and drafted the manuscript. Yi-Hsin Yang assisted in defining the statistical
43 analysis. Min-Hui Chiu and Su-Chen Hsu contributed to the data collection and
44 analysis. Hsiu-Fen Hsieh and Huei-Mein Chen assisted with drafting the manuscript.
45
46 All authors had full access to all data in the study and take responsibility for its
47 integrity.
48
49
50
51
52
53

54 **Data sharing statement**

No additional data are available.

REFERENCES

1. Chen CX, Kwekkeboom KL, Ward SE. Beliefs about dysmenorrhea and their relationship to self-management. *Res Nurs Health* 2016;39(4):263-76.
2. Eryilmaz G, Ozdemir F. Evaluation of menstrual pain management approaches by northeastern Anatolian adolescents. *Pain management nursing: official journal of the American Society of Pain Management Nurses* 2009;10(1):40-7.
3. Kato T. Effects of flexibility in coping with menstrual pain on depressive symptoms. *Pain practice: the official journal of World Institute of Pain* 2017;17(1):70-77.
4. Zhu X, Wong F, Bensoussan A, Lo SK, Zhou C, Yu J. Are there any cross-ethnic differences in menstrual profiles? A pilot comparative study on Australian and Chinese women with primary dysmenorrhea. *The journal of obstetrics and gynaecology research* 2010;36(5):1093-101.
5. Abdul-Razzak KK, Ayoub NM, Abu-Taleb AA, Obeidat BA. Influence of dietary intake of dairy products on dysmenorrhea. *The journal of obstetrics and gynaecology research* 2010;36(2):377-83.
6. Agarwal AK, Agarwal A. A study of dysmenorrhea during menstruation in adolescent girls. *Indian journal of community medicine : official publication of Indian Association of Preventive & Social Medicine* 2010;35(1):159-64.
7. Loto OM, Adewumi TA, Adewuya AO. Prevalence and correlates of dysmenorrhea among Nigerian college women. *The Australian & New Zealand journal of obstetrics & gynaecology* 2008;48(4):442-4.
8. Chang YT, Chen YC. Menstrual health care behavior and associated factors among female elementary students in the Hualien region. *Journal of Nursing Research*

- 2008;16(1):11-6.
9. Chiou MH, Wang HH. Predictors of dysmenorrhea and self-care behavior among vocational nursing school female students. *Journal of Nursing Research* 2008;16(1):17-25.
10. Cheng HF, Lin YH. Selection and efficacy of self-management strategies for dysmenorrhea in young Taiwanese women. *Journal of clinical nursing* 2011;20:1018-25.
11. Palmer D. "To help a million sick, you must kill a few nurses": Nurses' occupational health, 1890–1914. *Nursing History Review* 2012;20(1):14-45.
12. Laschinger HK, Grau AL. The influence of personal dispositional factors and organizational resources on workplace violence, burnout, and health outcomes in new graduate nurses: a cross-sectional study. *International journal of nursing studies* 2012;49(3):282-91.
13. Payne LA, Rapkin AJ, Lung KC, Seidman LC, Zeltzer LK, Tsao JC. Pain catastrophizing predicts menstrual pain ratings in adolescent girls with chronic pain. *Pain Med* 2016;17(1):16-24.
14. Berkley KJ, McAllister SL. Don't dismiss dysmenorrhea! *Pain* 2011;152(9):1940-1.
15. Chiu MH, Wang HH, Hsu SC, Liu IP. Dysmenorrhoea and self-care behaviours among hospital nurses: a questionnaire survey. *Journal of clinical nursing* 2013;22(22):3130-40.
16. Wong LP. Attitudes towards dysmenorrhoea, impact and treatment seeking among adolescent girls: a rural school-based survey. *The Australian journal of rural health* 2011;19(4):218-23.
17. Zukr SM, Naing L, Hamzah TNT, Hussain NHN. Primary dysmenorrhea among

- 1
2
3 medical and dental university students in Kelantan: Prevalence and associated
4 factors. *International Medical Journal* 2009;16(2):93-99.
5
6
7 18. Lu I-C. Dysmenorrhea and related factors in Taiwanese adolescent girls.
8
9 *University of Texas at Austin* 2010.
10
11 19. Chang C, Chen FL, Chang CH, Hsu CH. A preliminary study on menstrual health
12 and menstrual leave in the workplace in Taiwan. *Taiwan J Public Health*
13 2011;30(5):436-50.
14
15 20. Schluter PJ, Turner C, Huntington AD, Bain CJ, McClure RJ. Work/life balance
16 and health: the nurses and midwives e-cohort study. *International Nursing*
17 *Review* 2011(58):28-36.
18
19 21. Cady RK, Diamond ML, Diamond MP, Ballard JE, Lener ME, Dorner DP, et al.
20 Sumatriptan-naproxen sodium for menstrual migraine and dysmenorrhea:
21 satisfaction, productivity, and functional disability outcomes. *Headache*
22 2011;51(5):664-73.
23
24 22. Bettendorf B, Shay S, Tu F. Dysmenorrhea: contemporary perspectives.
25 *Obstetrical and Gynecological Survey* 2008;63(9):597-603.
26
27 23. Perry M. Treatment options for dysmenorrhoea. *Practice Nursing* 2012;23(4):4.
28
29 24. Ju H, Jones M, Mishra GD. Smoking and trajectories of dysmenorrhoea among
30 young Australian women. *Tobacco control* 2016;25(2):195-202.
31
32 25. Chiou MH, Wang HH. The relationship between dysmenorrhea and menstrual
33 attitudes among female students in vocational nursing schools. *The Journal of*
34 *Nursing* 2004;51(4):45-52.
35
36 26. Parker MA, Sneddon AE, Arbon P. The menstrual disorder of teenagers (MDOT)
37 study: determining typical menstrual patterns and menstrual disturbance in a large
38 population-based study of Australian teenagers. *BJOG : an international journal*
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56

- 1
2
3 *of obstetrics and gynaecology* 2010;117(2):185-92.
- 4
5 27. Stubbs ML. Cultural perceptions and practices around menarche and adolescent
6 menstruation in the United States. *Annals of the New York Academy of Sciences*
7 2008;1135:58-66.
- 8
9
10
11 28. Cheng CY, Yang K, Liou SR. Taiwanese adolescents? Gender differences in
12 knowledge and attitudes towards menstruation. *Nursing & Health Sciences*
13 2007;9(2):127-34.
- 14
15
16
17 29. H LY. The medical utilization of chinese and western medicine in females with
18 dysmenorrhea. Kaohsiung Medical University, 2012.
- 19
20
21
22 30. Chiou M-H, Wang H-H, Yang Y-H. Effect of systematic menstrual health
23 education on dysmenorrheic female adolescents' knowledge, attitudes, and
24 self-care behavior. *Kaohsiung J Med Sci* 2007;23(4):183-90.
- 25
26
27
28 31. Brooks-Gunn J, Ruble D. The menstrual attitude questionnaire. *Psychosomatic*
29 *medicine* 1980;42:503-11.
- 30
31
32
33 32. S L. Prevalence and associated factors of premenstrual syndrome among nurses in
34 Taiwan. In: Ministry of Science and Technology, R. O. C., editor, 1994.
- 35
36
37
38 33. Morrison LA, Larkspur L, Calibuso MJ, Brown S. Women's attitudes about
39 menstruation and associated health and behavioral characteristics. *American*
40 *Journal of Health Behavior* 2010;34(1):90-100.
- 41
42
43
44 34. Lo Sing-Kai, Lin Ching-Fen. On the use and interpretation of Cronbach's alpha.
45 *The Journal of Nursing Research* 1998;6:82-9.
- 46
47
48
49 35. Lin YX. *When "Experts" meet the medical professionals: The women with*
50 *menstrual pain encountering western medicine*. Kaohsiung Medical University,
51 Taiwan, 2006.
- 52
53
54
55 36. Chung F, Yao C, Wan G. The associations between menstrual function and life

1
2
3 style/working conditions among nurses in Taiwan. *Journal of occupational health*
4
5 2005;47(2):149-56.
6

7 37. Takeuchi H, Oishi T, Harada T. Morningness-eveningness preference, and mental
8
9 and physical symptoms during the menstrual cycle of Japanese junior high school
10
11 students. *Sleep and Biological Rhythms* 2003(1):245-47.
12

13
14 38. Firat MZ, Kulakac O, Oncel S, Akcan A. Menstrual attitude questionnaire:
15
16 confirmatory and exploratory factor analysis with Turkish samples. *Journal of*
17
18 *advanced nursing* 2009;65(3):652-62.
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

STROBE Statement—checklist of items that should be included in reports of observational studies

	Item No	Page No	Recommendation
Title and abstract	1	1	(a) Indicate the study's design with a commonly used term in the title or the abstract
		2	(b) Provide in the abstract an informative and balanced summary of what was done and what was found
Introduction			
Background/rationale	2	4-8	Explain the scientific background and rationale for the investigation being reported
Objectives	3	4	State specific objectives, including any prespecified hypotheses
Methods			
Study design	4	9	Present key elements of study design early in the paper
Setting	5	9	Describe the setting, locations, and relevant dates, including periods of recruitment, exposure, follow-up, and data collection
Participants	6	9	<i>Cross-sectional study</i> —Give the eligibility criteria, and the sources and methods of selection of participants
Variables	7	10-12	Clearly define all outcomes, exposures, predictors, potential confounders, and effect modifiers. Give diagnostic criteria, if applicable
Data sources/ measurement	8*	10-12	For each variable of interest, give sources of data and details of methods of assessment (measurement). Describe comparability of assessment methods if there is more than one group
Bias	9	3	Describe any efforts to address potential sources of bias
Study size	10	9	Explain how the study size was arrived at
Quantitative variables	11	12	Explain how quantitative variables were handled in the analyses. If applicable, describe which groupings were chosen and why
Statistical methods	12	12	(a) Describe all statistical methods, including those used to control for confounding
			(b) Describe any methods used to examine subgroups and interactions
			(c) Explain how missing data were addressed
			(d) <i>Cohort study</i> —If applicable, explain how loss to follow-up was addressed
			<i>Case-control study</i> —If applicable, explain how matching of cases and controls was addressed
			<i>Cross-sectional study</i> —If applicable, describe analytical methods taking account of sampling strategy
			(e) Describe any sensitivity analyses

Continued on next page

Results			
Participants	13*	9	(a) Report numbers of individuals at each stage of study—eg numbers potentially eligible, examined for eligibility, confirmed eligible, included in the study, completing follow-up, and analysed (b) Give reasons for non-participation at each stage
Descriptive data	14*	13	(a) Give characteristics of study participants (eg demographic, clinical, social) and information on exposures and potential confounders (b) Indicate number of participants with missing data for each variable of interest
Outcome data	15*	13	<i>Cross-sectional study</i> —Report numbers of outcome events or summary measures
Main results	16	13	(a) Give unadjusted estimates and, if applicable, confounder-adjusted estimates and their precision (eg, 95% confidence interval). Make clear which confounders were adjusted for and why they were included (b) Report category boundaries when continuous variables were categorized (c) If relevant, consider translating estimates of relative risk into absolute risk for a meaningful time period
Other analyses	17	-	Report other analyses done—eg analyses of subgroups and interactions, and sensitivity analyses
Discussion			
Key results	18	19-24	Summarise key results with reference to study objectives
Limitations	19	3	Discuss limitations of the study, taking into account sources of potential bias or imprecision. Discuss both direction and magnitude of any potential bias
Interpretation	20	19-24	Give a cautious overall interpretation of results considering objectives, limitations, multiplicity of analyses, results from similar studies, and other relevant evidence
Generalisability	21	19-24	Discuss the generalisability (external validity) of the study results
Other information			
Funding	22	26	Give the source of funding and the role of the funders for the present study and, if applicable, for the original study on which the present article is based

*Give information separately for cases and controls in case-control studies and, if applicable, for exposed and unexposed groups in cohort and cross-sectional studies.

Note: An Explanation and Elaboration article discusses each checklist item and gives methodological background and published examples of transparent reporting. The STROBE checklist is best used in conjunction with this article (freely available on the Web sites of PLoS Medicine at <http://www.plosmedicine.org/>, Annals of Internal Medicine at <http://www.annals.org/>, and Epidemiology at <http://www.epidem.com/>). Information on the STROBE Initiative is available at www.strobe-statement.org.