

Supplemental table 1 - Proportion of missing data among singleton live births from 37 to 40 weeks' gestation.

	n (%)
Study population	18,652 (100.0)
Capacity of hospital	0 (0.0)
Baby's friendly initiative hospital	0 (0.0)
Type of payment for birth	0 (0.0)
Age	3 (0.1)
Years of schooling	94 (0.5)
Previous births	0 (0.0)
Previous caesarean section	64 (0.3)
Previous stillbirth or neonatal death	64 (0.3)
Maternal or foetal condition	64 (0.3)
Precursors of birth	0 (0.0)
Mode of Birth	0 (0.0)
Gestational age	0 (0.0)
Resuscitation at labour ward	61 (0.3)
Oxygen therapy during hospital stay	60 (0.3)
Neonatal ICU admission	61 (0.3)
Neonatal death	0 (0.0)
Transient tachypnoea of the newborn	189 (1.0)
Antibiotic use during hospital stay	188 (1.0)
Hypoglycemia in the first 48h of life	189 (1.0)
Phototherapy in the first 72h of life	190 (1.0)
Breastfeeding up to one hour after birth	1,114 (5.9)
Exclusive breastfeeding during hospital stay	20 (0.1)
Readmission for Phototherapy*	51 (0.4)

* Follow-up study population as the denominator (n= 12,775).