BMJ Open Association of endotoxaemia with serum free fatty acids in metabolically healthy and unhealthy abdominally obese individuals: a case-control study in northwest of Iran

Maryam Saghafi-Asl,¹ Parichehr Amiri,² Mahsa Naghizadeh,²
Seyed Mostafa Ghavami,³ Nahid Karamzad⁴

To cite: Saghafi-Asl M, Amiri P, Naghizadeh M, *et al.* Association of endotoxaemia with serum free fatty acids in metabolically healthy and unhealthy abdominally obese individuals: a case-control study in northwest of Iran. *BMJ Open* 2017;**7**:e015910. doi:10.1136/bmjopen-2017-015910

► Prepublication history and additional material are available. To view these files please visit the journal online (<http://dx.doi.org/10.1136/bmjopen-2017-015910>).

Received 9 January 2017

Revised 24 March 2017

Accepted 27 March 2017

CrossMark

For numbered affiliations see end of article.

Correspondence to

Mrs. Nahid Karamzad;
nahidkaramzad@gmail.com

ABSTRACT

Objectives This study aimed to compare serum free fatty acids (FFAs) and lipopolysaccharide-binding protein (LBP) between metabolically healthy abdominally obese (MHAO) and metabolically unhealthy abdominally obese (MUAO) individuals. We also examined the association between serum FFAs and LBP in the participants.

Methods In this age-matched and gender-matched case-control study, 164 abdominally obese subjects were recruited from June to November 2015 in the northwest of Iran. Demographic data, dietary intake, body composition, anthropometric indices and physical activity (PA) were assessed. Basal blood samples were collected to determine serum metabolic parameters, FFAs and LBP. Abdominal obesity was defined as having waist circumference ≥ 95 cm. Those with three or more metabolic alterations were defined as MUAO and those having two or less were classified as MHAO. Data were analysed using SPSS V.17.0.

Results There were no significant differences in dietary intake, anthropometric indices, body composition and PA between the two groups. The odds of MUAO significantly increased by increments in serum fasting blood sugar (OR 3.79, 95% CI 2.25 to 6.40), triglycerides (OR 1.10, 95% CI 1.05 to 1.15), systolic blood pressure (OR 1.02, 95% CI 1.00 to 1.04) and diastolic blood pressure (OR 1.03, 95% CI 1.01 to 1.06) and decreased by increase in serum high-density lipoprotein cholesterol (OR 0.32, 95% CI 0.20 to 0.52). The levels of LBP and FFAs showed no significant differences between the two groups. However, significant correlations were found between LBP and FFAs in pooled population ($r=0.712$; $p<0.001$) as well as in cases ($r=0.717$; $p<0.001$) and controls ($r=0.704$; $p<0.001$). Neither FFAs nor LBP were significantly correlated with dietary intake or metabolic parameters ($p>0.05$).

Conclusion The results indicated that serum LBP and FFAs are highly correlated both in MHAO and MUAO states. In addition, the levels of LBP and FFAs seem to be more related to abdominal obesity than to the presence or absence of metabolic health.

INTRODUCTION

Obesity is increasingly prevalent worldwide.¹ There are well-established health

Strengths and limitations of this study

- The association of lipopolysaccharide-binding protein and free fatty acids levels in metabolically healthy abdominally obese and metabolically unhealthy abdominally obese individuals for the first time and found significant differences between the two parameters.
- This was a case-control study in which causality could not be assessed.
- Insulin levels were not measured in our study population; therefore, insulin resistance was not studied.
- The present work was carried out on volunteer participants. Though all volunteers were randomly recruited from general population after public announcement and based on the eligible criteria.

consequences of obesity such as type 2 diabetes, metabolic syndrome (MetS) and cardiovascular disease.² However, not all obese people are at higher risk of metabolic diseases. In a subtype of obese persons, described as 'metabolically healthy obese (MHO)', the obese phenotype may exist devoid of metabolic dysfunction.³ Despite there still being no uniform definition for MHO, it is thought to account for approximately one-fifth of the obese population.⁴

Evidence increasingly identifies inflammation as a potential mechanism linking adiposity, especially abdominal fat and metabolic dysfunction.⁵ However, published results are rare and conflicting regarding the role of inflammation in the metabolic differences observed between metabolically healthy and unhealthy individuals.^{6,7} Studies on postmenopausal obese women suggest that the MHO may have more favourable inflammatory profiles⁸ and less visceral fat⁹ than

their counterparts with insulin resistance (IR) and other metabolic abnormalities.¹⁰ In contrast, another study¹¹ reported that MHO women displayed abnormal levels of inflammatory profile, despite not having increased 10 year risk of cardiovascular disease.

The basic mechanism accounted for inflammation in adipose tissue is still unknown, but some factors including plasma free fatty acids (FFAs) are suggested.¹² It was clarified that plasma FFAs are increased among the obese as they are released from inflamed adipose tissue¹³ and through the lipolysis of adipocytes.¹⁴ However, little is known about the contribution of FFAs to the development of inflammation in obesity. Therefore, examining the association of FFAs with inflammatory markers seems to be warranted.

Lipopolysaccharide (LPS) molecules, also known as bacterial endotoxins, may trigger inflammation, leading to activation of immunity and cytokine release. LPS infusion and consequent subclinical endotoxaemia results in elevated levels of proinflammatory markers and metabolic aberrations.^{15 16} LPS has a short half-life¹⁷ and there is no agreement on the measurement of its plasma level.¹⁸ Hence, lipopolysaccharide-binding protein (LBP) is introduced with longer half-life and more reliable measurement.^{19 20} Also, serum LBP level is a proxy of serum LPS level.²¹ A population-based study²² found that LBP was significantly associated with MetS in normal weight participants. Another study²³ reported that among MetS components LBP concentration was independently associated with abdominal obesity.

In prior studies, inflammatory parameters were compared between obese and lean subjects.^{19 24 25} Therefore, it remains unclear whether the observed alterations in serum FFAs and/or inflammatory parameters in metabolically unhealthy obese patients are due to adiposity and/or metabolic state. Therefore, regarding the significant confounding effect of abdominal obesity, we used waist circumference (WC), as a reflection of visceral adipose tissue,²⁶ to define abdominal obesity and examined differences in characteristics and inflammatory markers (serum LBP and FFAs) between 'metabolically healthy' and 'unhealthy' abdominally obese individuals. We also examined the association between serum FFAs and LBP in pooled population as well as in each group.

METHODS

Study design and participants

A total of 81 metabolically healthy abdominally obese (MHAO) with 83 age-matched and gender-matched metabolically unhealthy abdominally obese (MUHAO) were recruited in this case-control study, carried out from 15 June to 6 November 2015 in the northwest of Iran. Frequency matching was carried out for the present study. Apparently healthy individuals aged 18–60 years with abdominal obesity were included in the study. We excluded pregnant or lactating women, those with diarrhoea for 3 consecutive days within the previous 3

months, diagnosed diabetes, coronary heart disease, stroke, myocardial infarction, cardiovascular and kidney, liver or infectious diseases including tuberculosis, AIDS and hepatitis; thyroid problems, severe mental disorders or physical disabilities and malignancies; taking oral anti-diabetic agents or insulin or other drugs for the past 2 months or antibiotics used for 3 consecutive days within the previous 3 months; smokers or alcohol consumers; misreported dietary intakes (<800 kcal/day or >4200 kcal/day) or being on specific diets in the past 6 months and having gastrointestinal surgery within the past 1 year.

Abdominal obesity was defined as having WC ≥ 95 cm according to the Iranian National Committee of Obesity.²⁷ According to Meigs *et al*,²⁸ metabolic health was defined as the presence of <3 of the following metabolic abnormalities including abdominal obesity (WC ≥ 95 cm for both genders)²⁷; high serum triglyceride (TG) concentration (≥ 150 mg/dL); low serum high-density lipoprotein cholesterol (HDL-C) (<40 mg/dL for men and <50 mg/dL for women); elevated blood pressure (BP) ($\geq 130/85/85$ mm Hg) and fasting blood sugar (FBS) (≥ 100 mg/dL). Eighty-one individuals with ≥ 3 criteria entered the case group (MUAO) and 83 with ≤ 2 criteria formed the control group (MHAO).

Sampling procedures

After public announcement for the study, 500 volunteers were recruited from general population. Of these, 178 people could enter the study based on the defined eligibility criteria for the present study. Informed consent was taken from each participant before the study. After taking blood samples and anthropometric measurements, 14 of them were excluded due to diabetes (FBS ≥ 126 mg/dL²⁹ in two occasions), leaving 164 people (82 men, 82 women) to conduct the research.

Biochemical assays

After a 12 hour overnight fast, 5 cm³ blood was obtained for serum analyses. After centrifugation at 3000 rpm for 5 min, metabolic parameters were analysed immediately, but serum FFAs and LBP were analysed after supplying in -80°C .

FBS was measured by the enzymatic colorimetric method using glucose oxidase. Serum TG concentration was measured by commercially available enzymatic reagents with glycerol phosphate oxidase. Serum HDL-C was measured after precipitation of the apolipoprotein B-containing lipoproteins with phosphotungstic acid. Assays were performed using Pars Azmoon kits (Pars Azmoon, Tehran, Iran) and a Selectra 2 auto-analyzer (Vital Scientific, Spankeren, the Netherlands). Interassay and intra-assay coefficient of variation (CV) was <5% for all assays. Serum samples for both LBP and FFAs assays were stored at -80°C until analysis. Both serum LBP and FFAs levels were determined by a sandwich ELISA (Bioassay Technology Laboratory, Shanghai Korean Biotech, Shanghai City, China) according to the manufacturer's instructions. The intra-assay and interassay CVs were <8% and <10%, respectively.

Measurements

All anthropometric indices were measured by a trained researcher. Height (without shoes in standard situation with precision of 0.1 cm and with an inelastic measuring tape) and weight (with Seca scale, light clothes and precision of 0.1 kg) were measured and body mass index (BMI) was calculated as weight in kilogram divided by the square of height in metres.³⁰ WC was measured using a non-stretchable fibre measuring tape. The subjects were asked to stand erect in relaxed position with both feet together on flat surface. WC was measured as the smallest horizontal girth between the costal and iliac crests at minimal respiration. Hip circumference was taken as the greatest circumference at the level of greater trochanters (the widest portion of the hip) on both sides. Waist to hip ratio was calculated by dividing WC (cm) by hip circumference (cm).³¹ BP was recorded in a comfortable sitting position in the left arm

after at least a 5 min rest using the mercury sphygmomanometer. Two measurements were taken and the mean of the two measurements was considered as the BP.³² Bioelectrical Impedance Analysis (BC-418MA, Tanita, Japan) was used to describe fat per cent, fat mass and fat-free mass. Dietary intake was assessed using a 3-day food record (1 weekend day and 2 workdays). *Nutritionist IV* software (Axxya Systems, Stafford, Texas, USA), modified for Iranian foods, was used for dietary data analysis. Physical activity (PA) was measured via long-form International Physical Activity Questionnaire (IPAQ).³³

Statistical analysis and sample size

To examine the normal distribution of variables, Kolmogorov-Smirnov tests and histograms were applied. The independent samples t-test was used to compare the means (SD) of normally distributed variables between the two

Table 1 Demographic, anthropometric and dietary intake parameters in metabolically unhealthy abdominally obese (MUAO) and metabolically healthy abdominally obese (MHAO) patients

Variables	MUAO (n=81)	MHAO (n=83)	OR (95% CI)	p Value
Age (years)*	38.23 (8.52)	37.13 (8.64)	1.01 (0.97 to 1.05)	0.412†
Men (%)	50.6	49.4	1.05 (0.57 to 1.93)	0.876‡
Physical activity score§	3144 (1416–5166)	2412 (1260–5211)	1.00 (0.99 to 1.00)	0.451¶
Weight (kg)*	87.21 (13.90)	84.78 (13.98)	1.01 (0.99 to 1.03)	0.266†
Height (cm)*	165.09 (11.56)	164.56 (10.60)	1.00 (0.97 to 1.03)	0.762†
Waist circumference (cm)*	106.02 (8.30)	105.06 (8.63)	1.01 (0.97 to 1.05)	0.470†
Hip circumference (cm)*	110.90 (6.92)	111.31 (8.26)	0.99 (0.95 to 1.03)	0.730†
Waist-to-hip ratio*	0.95 (0.05)	0.94 (0.06)	2.73 (0.15 to 48.00)	0.209†
BMI (kg/m ²)*	32.16 (4.25)	31.35 (4.12)	1.04 (0.97 to 1.12)	0.214†
Body fat percentage (%)*				
Males	26.86 (5.15)	25.07 (4.86)	1.07 (0.98 to 1.17)	0.093†
Females	38.59 (4.38)	39.88 (4.79)	0.93 (0.85 to 1.03)	0.227†
Body fat mass (kg)*				
Males	25.91 (7.27)	22.79 (6.76)	1.06 (1.00 to 1.13)	0.062†
Females	31.11 (7.33)	32.4 (8.63)	0.98 (0.92 to 1.03)	0.441†
Body fat-free mass (kg)*				
Males	69.09 (6.69)	66.98 (7.85)	1.04 (0.98 to 1.10)	0.130†
Females	48.72 (5.08)	47.61 (4.94)	1.04 (0.95 to 1.14)	0.425†
Total energy intake (kcal/day)*	2152.9 (765.1)	2206.8 (862.9)	1.00 (0.97 to 1.04)	0.700 †
Carbohydrate intake (% energy)*	60.20 (10.09)	59.29 (9.11)	1.01 (0.97 to 1.04)	0.499†
Protein intake (% energy)*	14.22 (2.95)	14.17 (4.10)	1.00 (0.91 to 1.09)	0.937†
Total fat intake (% energy)*	25.58 (10.54)	26.54 (12.14)	0.98 (0.95 to 1.02)	0.380†
Total SFA intake (% energy)§	14.06 (10.6–21.87)	14.21 (10.18–21.49)	0.99 (0.96 to 1.02)	0.780¶
Total MUFA intake (% energy)§	16.39 (11.01–24.68)	18.26 (11.76–26.27)	0.99 (0.97 to 1.01)	0.183 ¶
Total PUFA intake (% energy)§	12.7 (9.59–22.19)	14.41 (8.94–19.01)	1.01 (0.98 to 1.03)	0.943¶

*Variables with normal numeric scales are reported as mean (SD).

†Independent samples t-test.

‡ χ^2 test.

§Variables with non-normal numeric scales are reported as median (25th, 75th).

¶Mann-Whitney U test.

BMI, body mass index; MUFA, monounsaturated fatty acids; PUFA, polyunsaturated fatty acids; SFA, saturated fatty acids.

Figure 1 Flow chart of the study. BMI, body mass index; FBS, fasting blood sugar; TG, triglyceride.

groups. The Mann-Whitney U test was used for values with skewed distribution and in such conditions median (25th, 75th) was reported. In order to assess the association of two categorical variables, χ^2 test was applied. The correlation between serum FFAs and LBP was assessed using Spearman's correlation coefficient analysis. ORs and their 95% CIs were reported using logistic regression test.

The larger sample size was calculated for serum FFAs compared with LBP using literature-derived data³⁴ for patients with non-alcoholic fatty liver disease (NAFLD);

the effect size for serum FFAs was 0.20 nmol/L ($SD_1=0.34$ nmol/L for controls and $SD_2=0.53$ nmol/L for patients with NAFLD). Therefore, sample size estimation was based upon this parameter with 80% power and α -error of 5% and a case-to-control ratio of 1:1. It was predicted that 79 persons in each group would detect changes in serum FFAs as well as serum LBP level using the two-means formula. Data were analysed using SPSS V.17.0 for Windows (PASW Statistics). p value <0.05 was considered significant. Finally we reported the study based on the STROBE statement for Case-Control studies (Supplementary file 1).

Figure 2 Metabolic characteristics of metabolically unhealthy abdominally obese and metabolically unhealthy abdominally obese subjects. $p<0.001$ for all except waist circumference (WC), using χ^2 . FBS, fasting blood sugar; HDL-C, high-density lipoprotein cholesterol; Htn, hypertension; TG, triglycerides.

RESULTS

Males comprised about 50% of the study participants in the two groups ($p=0.87$). The age range of the subjects was 20–59 years. Participants of the two study groups similarly had WC ≥ 95 cm, that is, the cut-off point of WC for Iranian population. Overall, there were no significant differences in age, gender, anthropometric indices and body composition between the two groups. Dietary parameters, especially total fat, saturated fatty acids (SFA), monounsaturated fatty acids (MUFA) and polyunsaturated fatty acids (PUFA) intakes, were more or less the same between the two study groups (table 1). Marital status, education level and job of the cases and controls were also similar. However, mean number of

Table 2 Biochemical characteristics in metabolically unhealthy abdominally obese (MUAO) and metabolically healthy abdominally obese (MHAO) patients

Variables	MUAO (n=81)	MHAO (n=83)	OR (95% CI)	p Value
FBS (mg/dL)†	87.72 (5.82)	95.50 (9.76)	3.79 (2.25 to 6.40)*	<0.001‡
TG (mg/dL)†	193 (151–241)	112 (88 to 146)	1.10 (1.05 to 1.15)*	<0.001§
HDL-C (mg/dL)¶	39.53 (6.65)	46.44 (9.20)	0.32 (0.20 to 0.52)*	<0.001‡
SBP (mg/dL)†	115 (16.45)	108.13 (16.60)	1.02 (1.00 to 1.04)*	0.009‡
DBP (mg/dL)†	77.31 (13.86)	70.84 (12.94)	1.03 (1.01 to 1.06)*	0.002‡
Cholesterol (mg/dL)¶	193.60 (41.37)	187.37 (32.91)	1.00 (0.99 to 1.01)	0.286‡

*Statistically significant ($p < 0.05$).

†Variables with non-normal numeric scales are reported as median (25th, 75th).

‡Independent samples t-test.

§Mann-Whitney U test.

¶Variables with normal numeric scales are reported as mean (SD).

DBP, diastolic blood pressure; FBS, fasting blood sugar; HDL-C, high-density lipoprotein cholesterol; SBP, systolic blood pressure; TG, triglycerides.

metabolic aberrations were significantly higher in cases than controls (3.25 ± 0.72 vs 1.67 ± 0.50 ; $p < 0.001$) (data not shown). The flow chart of the study is shown in [figure 1](#).

Except for WC which was matched between the two groups, metabolic aberrations including low HDL-C (82% vs 34%), high TG (78% vs 24%), high FBS (33% vs 0%) and hypertension (34% vs 10%) were significantly higher in the cases than controls, respectively ([figure 2](#)). The current study indicated that each 10-unit increment in serum FBS level increased the risk of MUAO about 3.8 times (OR 3.79, 95% CI 2.25 to 6.40). Additionally, the odds of MUAO was significantly increased per one increment in serum TG level (OR 1.10, 95% CI 1.05 to 1.15), the systolic blood pressure (SBP) (OR 1.02, 95% CI 1.00 to 1.04) and diastolic blood pressure (DBP) (OR 1.03, 95% CI 1.01 to 1.06). However, the odds of having MUAO was significantly decreased by 68% per 10-unit increment in serum HDL-C level (OR 0.32, 95% CI 0.20 to 0.52) ([table 2](#)).

The median of LBP ($12.32 \mu\text{g/mL}$ in cases vs $12.76 \mu\text{g/mL}$ in controls, $p = 0.483$) and FFAs (1294 nmol/L in cases vs 1333 nmol/L in controls; $p = 0.686$) showed no significant difference between the two groups ([figure 3](#)). However, a significant correlation was found between LBP and FFAs in pooled population ($r = 0.712$; $p < 0.001$) as well as in cases ($r = 0.717$; $p < 0.001$) and controls ($r = 0.704$; $p < 0.001$) ([figure 4](#)). The results of partial correlation indicated much stronger correlation between LBP and FFAs, when controlling for WC ($r = 0.961$; $p < 0.001$). Moreover, number of metabolic aberrations was significantly correlated with HDL-C ($r = -0.537$; $p < 0.001$), TG ($r = 0.468$; $p < 0.001$), FBS ($r = 0.534$; $p < 0.001$), SBP ($r = 0.247$; $p = 0.001$) and DBP ($r = 0.315$; $p < 0.001$). Neither FFAs nor LBP were significantly correlated with dietary intake of total fat, SFA, MUFA and PUFA (data not shown). There were also no significant correlations of LBP and FFAs with metabolic parameters ([table 3](#)).

Figure 3 Lipopolysaccharide-binding protein (LBP) ($\mu\text{g/mL}$) (A) and free fatty acids (FFAs) concentrations (nmol/L) (B) in subjects with metabolically healthy abdominally obese (controls) and metabolically unhealthy abdominally obese (cases). $p = \text{not significant}$. Data are presented as box plot where boxes represent the IQR, the line within boxes represents the median and the lines outside the boxes represent the lower quartile minus 1.5 times the IQR or the upper quartile plus 1.5 times the IQR.

Figure 4 Spearman's correlation between lipopolysaccharide-binding protein (LBP) and free fatty acids (FFAs) in pooled population (A) as well as in each study group (B).

DISCUSSION

The present study, to the best of our knowledge, examined the association of LBP and FFAs levels in MHAO and MUAO individuals for the first time and found significant differences between the two parameters. Anthropometric indices as well as body composition profile were similar between the two groups. Moreover, there were no significant differences in LBP and FFAs between MHAO and MUAO.

In the present study, we matched metabolically healthy with metabolically unhealthy individuals on abdominal fat which might explain why we did not find any differences in levels of FFAs and LBP, as inflammatory markers, and body composition between the two groups. Therefore, the levels of LBP and FFAs seem to be more related to abdominal obesity than to the presence or absence of metabolic health. This is further supported by the observation that serum FFAs or LBP levels were not correlated with metabolic parameters. Therefore, our findings suggest that increased levels of these two markers are not necessarily related to the presence of metabolic aberrations.

A few smaller studies have examined differences in body composition and/or inflammatory profile between metabolically healthy and unhealthy obese postmenopausal women.^{35–37} In line with our result, Engström *et al.*³⁶ in a research on 58 obese postmenopausal women, found no

significant differences in levels of inflammatory markers between those with MetS compared with those without MetS. Additionally, in the population-based study of Phillips *et al.*³⁸, no significant difference was noted in C reactive protein (CRP) level between MHAO and MUAO, based on metabolic health criteria of the Meigs *et al.*²⁸ study. It is noteworthy that in the present study we used Meigs's metabolic health definition in which WC has also been considered.

A recent study revealed that the association between inflammatory biomarkers and metabolically healthy obesity depends on the criteria used. Since in that research, a significant difference was noted in the levels of CRP and interleukin-6 (IL-6) with some but not all MHAO definitions, which disappeared after adjustment for abdominal obesity or per cent body fat.³⁹ This study confirms our results. However, Phillips *et al.*⁴⁰ showed that obese women and men with MetS had significantly higher levels of inflammatory cytokines than obese persons without MetS. Beasley *et al.*⁴¹ showed that visceral adiposity, and not abdominal subcutaneous fat, was most consistently associated with significantly higher levels of IL-6 and CRP levels in black and white men and women in the Health ABC study. We could not measure visceral fat in our study, though, abdominal obesity measured through WC, can reflect visceral adiposity.²⁶ On the other

Table 3 Correlation of free fatty acids (FFAs) and lipopolysaccharide-binding protein(LBP) with metabolic parameters in metabolically unhealthy abdominally obese (MUAO) and metabolically healthy abdominally obese (MHAO) individuals

Variables	MUAO		MHAO	
	FFAs	LBP	FFAs	LBP
WC	0.07 (0.51)	0.02 (0.85)	0.06 (0.58)	0.03 (0.74)
TG	−0.02 (0.79)	0.008 (0.94)	−0.07 (0.48)	0.07 (0.49)
FBs	−0.005 (0.96)	−0.18 (0.09)	0.07 (0.50)	0.09 (0.40)
HDL-C	0.01 (0.91)	0.08 (0.45)	0.10 (0.33)	0.41 (0.09)
SBP	−0.03 (0.78)	−0.08 (0.46)	−0.05 (0.64)	−0.11 (0.29)
DBP	−0.08 (0.43)	−0.18 (1.00)	0.06 (0.58)	−0.09 (0.39)

p = not significant, using Spearman's correlation coefficient test.

DBP, diastolic blood pressure; HDL-C, high-density lipoprotein cholesterol; SBP, systolic blood pressure; TG, triglycerides; WC, waist circumference.

hand, a recent work observed no significant differences in visceral fat between the obese-insulin-resistant and obese-insulin-sensitive persons.⁴²

Several studies have demonstrated a strong association of IR with obesity, low HDL-C, hypertriglyceridaemia and hypertension^{10 43} as well as inflammatory factors.⁴⁴ However, in our study, we could not assess IR due to some financial deficits.

In the present work, dietary intake was compared between the two obese groups; therefore, no significant difference was found in terms of energy or macronutrients, especially fat intake. Moreover, habitual PA was controlled between the study groups. And, in our previous report, there were no significant differences between the two groups in terms of PA (unpublished data). Since different levels of habitual PA might affect levels of serum inflammatory markers.⁴⁵

Obesity, as a well-known metabolic risk factor, is usually associated with mild chronic inflammation.⁴⁶ The relationship between obesity and increased inflammation may be justified, in part, by FFAs⁴⁷ which are released from adipocytes through lipolysis and are elevated in obesity due to increased adipose tissue.⁴⁸ Inflammatory cytokines such as IL-6 can stimulate lipolysis and increase levels of FFAs.⁴⁹ On the other hand, in healthy persons an acute increase in FFAs can induce inflammatory changes.⁵⁰ Therefore, FFAs are not only increased by inflammation, but also promote inflammation. The results of the present research showed that FFAs are positively correlated with LBP levels either in the pooled population or in each group. It shows that any increase in the level of FFAs, observed in the abdominally obese, regardless of their metabolic aberrations can lead to a significant elevation in the level of LBP.

LBP has been considered a key inflammatory marker which mediates LPS-triggered innate immunity.⁵¹ Although LBP concentration was previously reported to be associated with various anthropometric and metabolic factors such as BMI, WC and so on,^{23–25} in our study the relationship only existed between the two biomarkers, FFAs and LBP, but not with the metabolic or anthropometric parameters. It is notable that the positive relationship between LBP and BMI was not observed in either normal weight or obese groups in the Yang *et al*⁵² study after multivariate analyses. In their research, the level of LBP significantly reduced after bariatric surgery and consequent reduction in WC (from 121.6 cm to 90.6 cm; $p < 0.001$) which indicates the strong association of LBP with WC. Liu *et al*,²² in a population-based follow-up study on 2529 Chinese, also found that the association of LBP with MetS was significant only in normal weight participants, but not in their overweight/obese counterparts after multivariate adjustments including BMI, which supports our study's findings. It is assumed that the association between serum LBP level and MetS observed in previous studies^{19 24 25} is mediated by BMI or WC, and finding no association between serum LBP level and incidence of MetS in our study, in which the WC-matched controls were included, is not unexpected.

Overall, what makes our research different from most of the previous ones is that in our study we matched the two groups based on WC, rarely observed in prior reports. Most of the previous studies have examined either MetS patients versus those without the syndrome or metabolically healthy versus metabolically unhealthy, regardless of their BMI or WC status and based on different metabolic health criteria.^{3 8 28} A few have examined inflammatory markers between metabolically healthy and unhealthy persons, considering WC or abdominal obesity.^{36 38 39}

CONCLUSION

Our study indicated that WC could be a strong mediator of the association between serum LBP, FFAs and metabolic alterations. In fact, the levels of LBP and FFAs seem to be more related to abdominal obesity than to the presence or absence of metabolic health. The results also suggested a significant correlation between serum FFAs and LBP in abdominally obese population, which seems to be independent of metabolic aberrations.

Author affiliations

¹Department of Biochemistry and Diet Therapy, Nutrition Research Center, School of Nutrition and Food Sciences, Tabriz University of Medical Sciences, Tabriz, The Islamic Republic of Iran

²Student Research Committee, School of Nutrition and Food Sciences, Tabriz University of Medical Sciences, Tabriz, The Islamic Republic of Iran

³Department of Radiology, School of Paramedicine, Tabriz University of Medical Sciences, Tabriz, The Islamic Republic of Iran

⁴Department of Public Health, School of Nursing and Midwifery, Maragheh University of Medical Sciences, Maragheh, The Islamic Republic of Iran

Acknowledgements The authors thank those who participated in this study.

Contributors MS-A and NK conceived the study design and wrote the study protocol. MS-A and NK analysed and interpreted the data. MS-A, PA, MN, SMG and NK have been involved in drafting the manuscript or revising it critically for content. All authors have given final approval of the version to be published.

Funding This work was financially supported by MRGUMS and TBZMED, Iran.

Disclaimer None.

Competing interests None declared.

Patient consent Obtained.

Ethics approval All protocols were approved by ethics committees of Maragheh University of Medical Sciences (MRGUMS) and Tabriz University of Medical Sciences (TBZMED). Research was carried out in compliance with the Declaration of Helsinki.

Provenance and peer review Not commissioned; externally peer reviewed.

Data sharing statement No additional data available.

Open Access This is an Open Access article distributed in accordance with the Creative Commons Attribution Non Commercial (CC BY-NC 4.0) license, which permits others to distribute, remix, adapt, build upon this work non-commercially, and license their derivative works on different terms, provided the original work is properly cited and the use is non-commercial. See: <http://creativecommons.org/licenses/by-nc/4.0/>

© Article author(s) (or their employer(s) unless otherwise stated in the text of the article) 2017. All rights reserved. No commercial use is permitted unless otherwise expressly granted.

REFERENCES

1. Kelly T, Yang W, Chen CS, *et al*. Global burden of obesity in 2005 and projections to 2030. *Int J Obes* 2008;32:1431–7.
2. Malik VS, Willett WC, Hu FB. Global obesity: trends, risk factors and policy implications. *Nat Rev Endocrinol* 2013;9:13–27.

3. Karelis AD, Rabasa-Lhoret R. Inclusion of C-reactive protein in the identification of metabolically healthy but obese (MHO) individuals. *Diabetes Metab* 2008;34:183–4.
4. Primeau V, Coderre L, Karelis AD, et al. Characterizing the profile of obese patients who are metabolically healthy. *Int J Obes* 2011;35:971–81.
5. Hotamisligil GS. Inflammation and metabolic disorders. *Nature* 2006;444:860–7.
6. Desai MY, Dalal D, Santos RD, et al. Association of body mass index, metabolic syndrome, and leukocyte count. *Am J Cardiol* 2006;97:835–8.
7. Van Guilder GP, Hoetzer GL, Greiner JJ, et al. Influence of metabolic syndrome on biomarkers of oxidative stress and inflammation in obese adults. *Obesity* 2006;14:2127–31.
8. Messier V, Karelis AD, Prud'homme D, et al. Identifying metabolically healthy but obese individuals in sedentary postmenopausal women. *Obesity* 2010;18:911–7.
9. Messier V, Karelis AD, Robillard ME, et al. Metabolically healthy but obese individuals: relationship with hepatic enzymes. *Metabolism* 2010;59:20–4.
10. Karelis AD, Faraj M, Bastard JP, et al. The metabolically healthy but obese individual presents a favorable inflammation profile. *J Clin Endocrinol Metab* 2005;90:4145–50.
11. Wildman RP, Kaplan R, Manson JE, et al. Body size phenotypes and inflammation in the Women's Health Initiative Observational Study. *Obesity* 2011;19:1482–91.
12. Ye J. Emerging role of adipose tissue hypoxia in obesity and insulin resistance. *Int J Obes* 2009;33:54–66.
13. Karpe F, Dickmann JR, Frayn KN. Fatty acids, obesity, and insulin resistance: time for a reevaluation. *Diabetes* 2011;60:2441–9.
14. Boden G. Role of fatty acids in the pathogenesis of insulin resistance and NIDDM. *Diabetes* 1997;46:3–10.
15. Laugerette F, Vors C, Peretti N, et al. Complex links between dietary lipids, endogenous endotoxins and metabolic inflammation. *Biochimie* 2011;93:39–45.
16. Mehta NN, McGillicuddy FC, Anderson PD, et al. Experimental endotoxemia induces adipose inflammation and insulin resistance in humans. *Diabetes* 2010;59:172–81.
17. Munford RS. Invited review: detoxifying endotoxin: time, place and person. *J Endotoxin Res* 2005;11:69–84.
18. Amar J, Burcelin R, Ruidavets JB, et al. Energy intake is associated with endotoxemia in apparently healthy men. *Am J Clin Nutr* 2008;87:1219–23.
19. Albillos A, de la Hera A, González M, et al. Increased lipopolysaccharide binding protein in cirrhotic patients with marked immune and hemodynamic derangement. *Hepatology* 2003;37:208–17.
20. Ruiz AG, Casafont F, Crespo J, et al. Lipopolysaccharide-binding protein plasma levels and liver TNF-alpha gene expression in obese patients: evidence for the potential role of endotoxin in the pathogenesis of non-alcoholic steatohepatitis. *Obes Surg* 2007;17:1374–80.
21. Román J, Caixàs A, Escoté X, et al. Lipopolysaccharide-binding protein is increased in patients with psoriasis with metabolic syndrome, and correlates with C-reactive protein. *Clin Exp Dermatol* 2013;38:81–4.
22. Liu X, Lu L, Yao P, et al. Lipopolysaccharide binding protein, obesity status and incidence of metabolic syndrome: a prospective study among middle-aged and older Chinese. *Diabetologia* 2014;57:1834–41.
23. Gonzalez-Quintela A, Alonso M, Campos J, et al. Determinants of serum concentrations of lipopolysaccharide-binding protein (LBP) in the adult population: the role of obesity. *PLoS One* 2013;8:e54600.
24. Moreno-Navarrete JM, Ortega F, Serino M, et al. Circulating lipopolysaccharide-binding protein (LBP) as a marker of obesity-related insulin resistance. *Int J Obes* 2012;36:1442–9.
25. Sun L, Yu Z, Ye X, et al. A marker of endotoxemia is associated with obesity and related metabolic disorders in apparently healthy Chinese. *Diabetes Care* 2010;33:1925–32.
26. Kuk JL, Lee S, Heymsfield SB, et al. Waist circumference and abdominal adipose tissue distribution: influence of age and sex. *Am J Clin Nutr* 2005;81:1330–4.
27. Azizi F, Khalili D, Aghajani H, et al. Appropriate waist circumference cut-off points among Iranian adults: the first report of the Iranian National Committee of Obesity. *Arch Iran Med* 2010;13:243.
28. Meigs JB, Wilson PW, Fox CS, et al. Body mass index, metabolic syndrome, and risk of type 2 diabetes or cardiovascular disease. *J Clin Endocrinol Metab* 2006;91:2906–12.
29. Moradi S, Keshavarzi A, Tabatabaee SM. Is stress hyperglycemia a predicting factor of developing Diabetes in Future? *Exp Clin Endocrinol Diabetes* 2015;123:614–6.
30. Mirmiran P, Esmailzadeh A, Azizi F. Detection of cardiovascular risk factors by anthropometric measures in Tehranian adults: receiver operating characteristic (ROC) curve analysis. *Eur J Clin Nutr* 2004;58:1110–8.
31. Taylor RW, Jones IE, Williams SM, et al. Evaluation of waist circumference, waist-to-hip ratio, and the conicity index as screening tools for high trunk fat mass, as measured by dual-energy X-ray absorptiometry, in children aged 3–19 y. *Am J Clin Nutr* 2000;72:490–5.
32. Azizi F, Ghanbarian A, Madjid M, et al. Distribution of blood pressure and prevalence of hypertension in Tehran adult population: tehran lipid and glucose study (TLGS), 1999–2000. *J Hum Hypertens* 2002;16:305–12.
33. Vasheghani-Farahani A, Tahmasbi M, Asheri H, et al. The Persian, last 7-day, long form of the International Physical Activity Questionnaire: translation and validation study. *Asian J Sports Med* 2011;2:106.
34. Zhang J, Zhao Y, Xu C, et al. Association between serum free fatty acid levels and nonalcoholic fatty liver disease: a cross-sectional study. *Sci Rep* 2014;4:5832.
35. Halkes CJ, van Dijk H, de Jaegere PP, et al. Postprandial increase of complement component 3 in normolipidemic patients with coronary artery disease: effects of expanded-dose simvastatin. *Arterioscler Thromb Vasc Biol* 2001;21:1526–30.
36. Engström G, Hedblad B, Eriksson KF, et al. Complement C3 is a risk factor for the development of diabetes: a population-based cohort study. *Diabetes* 2005;54:570–5.
37. Expert Panel on Detection, Evaluation, and Treatment of High Blood Cholesterol in Adults. Executive summary of the third report of the national cholesterol education program (NCEP) Expert panel on detection, evaluation, and treatment of high blood cholesterol in adults (Adult treatment panel III). *JAMA* 2001;285:2486–97.
38. Phillips CM, Kesse-Guyot E, Ahluwalia N, et al. Dietary fat, abdominal obesity and smoking modulate the relationship between plasma complement component 3 concentrations and metabolic syndrome risk. *Atherosclerosis* 2012;220:513–9.
39. Marques-Vidal P, Velho S, Waterworth D, et al. The association between inflammatory biomarkers and metabolically healthy obesity depends on the definition used. *Eur J Clin Nutr* 2012;66:426–35.
40. Phillips CM, Dillon C, Harrington JM, et al. Defining metabolically healthy obesity: role of dietary and lifestyle factors. *PLoS One* 2013;8:e76188.
41. Beasley LE, Koster A, Newman AB, et al. Inflammation and race and gender differences in computerized tomography-measured adipose depots. *Obesity* 2009;17:1062–9.
42. Muscari A, Massarelli G, Bastagli L, et al. Relationship of serum C3 to fasting insulin, risk factors and previous ischaemic events in middle-aged men. *Eur Heart J* 2000;21:1081–90.
43. Moro E, Gallina P, Pais M, et al. Hypertriglyceridemia is associated with increased insulin resistance in subjects with normal glucose tolerance: evaluation in a large cohort of subjects assessed with the 1999 World Health Organization criteria for the classification of diabetes. *Metabolism* 2003;52:616–9.
44. Schmidt MI, Duncan BB. Diabetes: an inflammatory metabolic condition. *Clin Chem Lab Med* 2003;41:1120–30.
45. You T, Nicklas BJ. Effects of exercise on adipokines and the metabolic syndrome. *Curr Diab Rep* 2008;8:7–11.
46. Fantuzzi G. Adipose tissue, adipokines, and inflammation. *J Allergy Clin Immunol* 2005;115:911–9.
47. Björntorp P, Bergman H, Varnauskas E. Plasma free fatty acid turnover rate in obesity. *Acta Med Scand* 1969;185:351–6.
48. Jensen MD, Haymond MW, Rizza RA, et al. Influence of body fat distribution on free fatty acid metabolism in obesity. *J Clin Invest* 1989;83:1168–73.
49. Boden G, Cheung P, Stein TP, et al. FFA cause hepatic insulin resistance by inhibiting insulin suppression of glycogenolysis. *Am J Physiol Endocrinol Metab* 2002;283:E12–E19.
50. Santomauro AT, Boden G, Silva ME, et al. Overnight lowering of free fatty acids with acipimox improves insulin resistance and glucose tolerance in obese diabetic and nondiabetic subjects. *Diabetes* 1999;48:1836–41.
51. Schumann RR. Old and new findings on lipopolysaccharide-binding protein: a soluble pattern-recognition molecule. *Biochem Soc Trans* 2011;39:989–93.
52. Yang PJ, Lee WJ, Tseng PH, et al. Bariatric surgery decreased the serum level of an endotoxin-associated marker: lipopolysaccharide-binding protein. *Surg Obes Relat Dis* 2014;10:1182–7.