BMJ Open Double-blind randomised controlled trial of the independent and synergistic effect of *Spirulina maxima* with exercise (ISESE) on general fitness, lipid profile and redox status in overweight and obese subjects: study protocol

Marco Antonio Hernández-Lepe,¹ José Alberto López-Díaz,² Laura Alejandra de la Rosa,² Rosa Patricia Hernández-Torres,³ Abraham Wall-Medrano,² Marco Antonio Juárez-Oropeza,⁴ José Pedraza-Chaverri,⁵ Rene Urquidez-Romero,² Arnulfo Ramos-Jiménez²

To cite: Hernández-Lepe MA, López-Díaz JA, Rosa LA, et al. Double-blind randomised controlled trial of the independent and synergistic effect of *Spirulina maxima* with exercise (ISESE) on general fitness, lipid profile and redox status in overweight and obese subjects: study protocol. *BMJ Open* 2017;7:e013744. doi:10.1136/bmjopen-2016-013744

► Prepublication history and additional material is available. To view please visit the journal (<http://dx.doi.org/10.1136/bmjopen-2016-013744>).

Received 4 August 2016

Revised 9 April 2017

Accepted 27 April 2017

CrossMark

For numbered affiliations see end of article.

Correspondence to

Dr Arnulfo Ramos-Jiménez; aramos@uacj.mx

ABSTRACT

Introduction In order to reduce cardiovascular disease risk factors, a healthy diet must include dietary antioxidants from different sources (eg, *Spirulina maxima*) and regular practice of exercise should be promoted. There is some evidence from animal studies that *S. maxima* and exercise decrease cardiovascular disease risks factors. However, very few studies have proved the independent or synergistic effect of *S. maxima* plus exercise in humans. This study attempts to address the independent and synergistic effects in overweight and obese subjects participating in a systematic physical exercise programme at moderate intensity on general fitness, plasma lipid profile and antioxidant capacity.

Methods and analysis Using a randomised, double-blind, placebo-controlled, counterbalanced crossover study design, 80 healthy overweight and obese subjects will be evaluated during a 12-week isoenergetic diet accompanied by 4.5 g/day *S. maxima* intake and/or a physical systematic exercise programme at moderate intensity. Body composition, oxygen uptake, heart rate, capillary blood lactate, plasma concentrations of triacylglycerols, total, low-density and high-density lipoprotein cholesterol, antioxidant status, lipid oxidation, protein carbonyls, superoxide dismutase, catalase, glutathione, glutathione peroxidase, glutathione reductase and paraoxonase will be assessed.

Ethics and dissemination This study and all the procedures have been approved by the Universidad Autonoma de Ciudad Juárez Bioethics Committee. Findings will be disseminated through peer-reviewed journals, national and international conferences.

Trial registration number ClinicalTrials.gov: NCT02837666.

BACKGROUND

Cardiovascular diseases (CVD) are the leading causes of death worldwide. Each

Strengths and limitations of this study

- Working with obese people, one of the main risk factors for cardiovascular diseases.
- This study can be extrapolated to different populations.
- Double-blind randomised controlled trial.
- There are no possible limitations to perform all the procedures described in the present paper.

year, 17.5 million people die because of these diseases.¹ Dyslipidaemias are a predisposing factor for CVD and are characterised by high concentrations of triacylglycerols (TAG), total cholesterol (TC), low-density lipoprotein cholesterol (LDL-c) and low concentrations of high-density lipoprotein cholesterol (HDL-c).² Besides dyslipidaemias, oxidative stress (OxS) is also observed to rise in obesity,^{3,4} and it is also a predisposing factor for CVD.^{5–8}

In order to reduce OxS, dyslipidaemias and the incidence of CVD, the intake of antioxidants from a fruit and vegetable-rich diet or nutritional supplements, mainly by unprocessed foods, has been proposed.^{9,10} In this sense, the cyanobacterium *Spirulina* is an important source of antioxidants, currently associated with cardiovascular protection properties.^{9,11}

For centuries *Spirulina maxima* (*S. maxima*) has been cultivated and used as a nutritional supplement due to its content of amino acids and essential fatty acids, vitamin C, vitamin E, tocopherols and phycocyanins.¹² Recently, studies on *S. maxima* have focused on verifying

the biological activity of its components, including hypolipidaemic and antioxidant effects.^{13–15} However, most studies have been conducted in animal models, with only a few studies focused on the biological effect in humans.

Furthermore, it is known that the practice of systematic physical exercise ameliorates the risk of CVD,^{10 16} and it has been observed that it also plays a role in OxS, especially physical exercise at high intensity (PEHI).¹⁷ PEHI considerably increases general metabolism, oxygen uptake (VO_2) and mitochondrial activity, thus increasing the production of reactive oxygen species (ROS).¹⁸ However, physical exercise of moderate intensity (PEMI) has the best protective effect against CVD, mainly due to physiological adaptations including the expression of antioxidant enzymes,¹⁹ which stop the formation and propagation of ROS and improving the redox status of the organism.^{20 21}

Algae and cyanobacteria have shown promising pharmacological properties providing health benefits and physical improvements.²² These effects are attributed to their specific profile in bioactive compounds such as uncommon carotenoids, phenolic compounds and tocopherols in addition to their well-known content in vitamins and good quality proteins associated with their specific chemical structure and interaction with biological membranes.²³

There is evidence that the cyanobacteria *Spirulina*, in addition to exercise, lowers the risks of CVD. This has mainly been observed in animal models.^{24–26} No studies with *Spirulina* and exercise experimental designs proving these benefits have yet been reported in humans. Therefore, the ISESE study will analyse the independent and synergistic effects of the intake of *S. maxima* with PEMI on general fitness, plasma lipid profile and redox status in overweight and obese subjects.

METHODS/DESIGN

Hypothesis

Spirulina maxima intake and a dosed physical activity programme will improve the lipid profile, general fitness and antioxidant status in overweight and obese subjects.

Objectives

The ISESE project's main objective is to demonstrate that ingestion of *S. maxima* and a dosed physical activity programme will decrease, both independently and synergistically, some CVD risks in overweight and obese subjects.

Outcome measures

The primary outcome measure will be changes in the plasma lipid profile after 6 weeks of treatment (plasma TAG, TC, HDL-c and LDL-c using standardised enzymatic methods).

The secondary outcome measure will be changes in general fitness (maximal oxygen uptake (VO_2 max), heart rate (HR), onset blood lactate accumulation and body fat mass) as well as changes in redox status (malondialdehyde

(MDA), protein carbonyls, paraoxonase (PON1), superoxide dismutase (SOD), catalase glutathione (GSH), glutathione reductase (GR) and glutathione peroxidase (GPx)).

Participant eligibility

The inclusion criteria will be subjects aged 18–45 years of both sexes: 40 overweight (body mass index (BMI) 25–29.9 kg/m²) and 40 obese (BMI >30 kg/m²). The exclusion criteria will be alcohol intake of >100 mL alcohol a week, taking drugs and/or food or vitamin supplements, diabetes or a physical or ECG injury that prevents engagement in any regular physical exercise. The elimination criteria will be attendance by the subject of <80% and/or 20% change in body weight during the study.

Enrolment, ethics and safety procedures

Overweight and obese students at the University of Ciudad Juárez, Mexico will be personally invited to participate. Participants will be informed of the purpose of the study and of the physical, clinical and biochemical procedures, benefits, physical intensity tests and risks before beginning the clinical trial. To guarantee the good physical and mental health of the participants at the beginning of the studies, a medical clinical laboratory study examination and an ECG at rest will be performed. The investigators are trained in cardiopulmonary resuscitation and the laboratory has the necessary communication tools and channels to perform emergency procedures. This study and all the procedures have been registered appropriately in the ClinicalTrials.gov database (NCT02837666) and approved by the Bioethics Committee of the Universidad Autónoma de Ciudad Juárez (see online Supplementary file). Participants' acceptance will be formalised by means of written informed consent and their anonymity and confidentiality will be strictly enforced. This study will be conducted in compliance with the principles of the Declaration of Helsinki.

Baseline measurements

On the day before beginning the treatment, subjects will visit the laboratory for baseline measurements. Body mass will be measured with subjects lightly dressed and barefoot using an electronic balance and standing height with a stadiometer. Body fat percentage will be measured by plethysmography (BodPod, COSMED, Rome, Italy), electric bioimpedance (in body160, BIOSPACE, Japan) and by dual-energy X-ray absorptiometry (DXA) (Prodigy v6.8; GE Lunar, Milwaukee, Wisconsin, USA) according to the manufacturers' guidelines. Anthropometric parameters will be taken by the standardised method of the International Society for the Advancement of Kinanthropometry (ISAK) and according to the ISAK Anthropometry Accreditation protocols.²⁷ Researchers are experienced in handling equipment and procedures.

Study design

The study consists of *S. maxima* treatment for 14 weeks in a randomised, double-blind, placebo-controlled,

Figure 1 Experimental design for independent and synergistic effect of *Spirulina maxima* and exercise. Same colour indicates the same group of participants.

counterbalanced crossover design aimed to eliminate interindividual bias as well as improve the reliability of the final results. This design is commonly used in clinical trials in order to have better results and conclusions.²⁸ Eligible participants ($n=80$) will be divided into two groups, one with systematic practice of physical exercise (GEx) and the other without it (GNEx). The participants of each group will be randomised in equal proportions between *S. maxima* and placebo treatment, 4.5 g daily in a non-transparent capsule for 12 weeks (6 weeks of the first treatment followed by a 2-week washout period, then 6 weeks of the second treatment) (figure 1). If any reported allergic reaction occurs the treatment will be discontinued.

To maintain the overall quality and legitimacy of the clinical trial, code interruptions should occur only in exceptional circumstances when knowledge of the actual treatment is essential for further management of the participant. Investigators are encouraged to discuss with a medical advisor if they believe that unblinding is necessary. Unblinding should not necessarily be a reason for study discontinuation.

A student outside the research team will feed data into the computer in separate datasheets so that the researchers can analyse the data without having access to information about the allocation treatments, storing all the participants' files in numerical order in a safe and accessible place. The participants' files will be stored for a period of 5 years after the end of the study.

The sample size was determined using the statistical programme G*Power,²⁹ selecting a sample of 73 subjects with $\alpha=0.05$ and $p=0.85$. This sample will be fixed to 80

subjects in order to anticipate possible withdrawals from the study.

Maximum intensity tests (MITs)

Each subject will participate in four stress tests performed at maximum intensity (MIT) during the study. During the MITs, oxygen consumed and carbon dioxide produced will be measured by a gas analyser (Cortex MetaLyzer 3B, Germany), HR with a Polar H7 sensor (Polar Electro, Lake Success, NY) and lactate in capillary blood samples will be assayed in a YSI 1500 Sport Lactate Analyzer (YSI Life Sciences, Ohio, USA). The MIT protocol consists of using a cycle ergometer (Monark Ergonomic 828 E; Monark Exercise AB, 105 Vansbro, Sweden) starting with a workload of 50–75 W with increments of 15–30 W every 3 min for 15 min or when the subject can no longer pedal more than 40 revolutions/min, finishing the test when 85–100% of the maximum HR reserve is reached. The MIT will not be valid if it lasts for less than 9 min

Figure 2 Design of maximum intensity tests. ↓ indicates the interval at which the resistance will be increased; in addition, physical perceived effort, heart rate and capillary blood samples will be measured.

(figure 2). At the end of each increment load (3 min), capillary blood samples will be taken to determine lactate and the physical perceived effort will be registered using the Borg scale. Measurements of HR, glucose and blood pressure will serve to check the subject's health during each MIT.

Before the first MIT, subjects will decide to be in the GEx or GNEEx and will then be divided randomly for treatment, consisting of a daily dose of 4.5 g of *Spirulina* or placebo in capsules which the subjects will be recommended to consume before each meal every day for 6 weeks. The day after the first supplementation period, subjects will come back to perform the second MIT with identical conditions to the first one. The third MIT will be performed after a washout period of 2 weeks to remove the effects of treatment and to avoid any possible delayed effect of *S. maxima* in the organism. The last MIT will be performed after the second treatment with identical conditions to the previous three occasions. General fitness parameters will be measured on the same day as the MITs to obtain accurate results.

Randomisation

All subjects who consent to participate and who meet the inclusion criteria will be randomly assigned by an investigator who will not have any other involvement in the clinical study; he will create a database with all the treatments throughout the duration of the study. Randomisation will therefore be carried out without any influence of the principal investigators. Participants will be randomly assigned to *S. maxima* or placebo treatment with a 1:1 allocation according to a computer-generated random schedule stratified by site and the baseline score of the Action Arm Research Test using permuted blocks of random sizes with an online central randomisation service (TENALEA). Block sizes will not be disclosed to ensure concealment.

Adherence assessments

Multiple methods will be used to assess treatment adherence including pill counting, reasons for non-compliance and use of the capsules. Participants will return each week to receive new capsules. All treatment data will be recorded on the appropriate case report form.

Dietary analysis

All participants will be subjected to a nutritional survey to define the daily calories required to establish a custom diet. Dietary intake will be monitored by retrospective methods including 24-hour recall (which investigates intake over a specific day) and a food frequency questionnaire (a summary of usual intake of different categories of foods). These accurate and validated techniques allow quantification of the types and quantities of foods and beverages consumed during the period of interest in the past. Dietary intake evaluations will be performed at each MIT of the study in order to record the possible variability in food consumption patterns.

Physical exercise protocol

General health assessment and physical activity questionnaires (PAR-Q and YOU) will be conducted to ensure that there are no physical impediments to exercise. GEx participants will exercise 5 days a week with the following protocol: 5–10 min of warm-up exercise, 20–30 min muscular endurance exercise and 20–30 min of aerobic exercise (cardiovascular exercise): walking, jogging, running and/or cycling. Three days a week aerobic intensities will be between 60% and 80% and 2 days between 70% and 90% of the maximum HR reserve, and five final minutes of stretching.

Sample collection and biochemical analysis

Blood samples (8 mL) will be collected before and after each MIT from the antecubital vein into ethylenediaminetetraacetic acid (EDTA) tubes after 10–12 hours of fasting; plasma will be obtained from the blood samples by refrigerated centrifugation (4°C) at 5000 g for 10 min. Plasma glucose TC, HDL-c, LDL-c and TAG concentrations will be analysed using standard enzymatic procedures (Jas Diagnostics, Mexico) with a spectrophotometer (Genesys 10 UV; Thermo Electron Corporation, USA). Plasma will be used to measure MDA and GSH content and activity of GPx and GR according to Jimenez-Osorio *et al.*³⁰ total antioxidant status (TAS) by the 2,2-diphenyl-1-picrylhydrazyl (DPPH) technique as previously described by Koren *et al.*³¹ and oxygen radical absorbance capacity following the methodology of Prior *et al.*³² The erythrocytes will be washed twice with cold 0.9% saline solution and used for SOD and catalase activity assays, as previously described by Jimenez-Osorio *et al.*³⁰ Plasma PON1 content will be measured following the method of Perez-Herrera *et al.*³³

Statistical analysis

To evaluate data distribution, normality and homoscedasticity tests will be conducted on each of the response variables analysed. In order to analyse differences between variables before and after the study and between the groups, an ANOVA test will be performed. To analyse possible associations between variables, a bivariate correlation analysis will be used. To analyse independence between variables, a multiple regression analysis will be performed. The software to be used is GraphPad Prism 6.

DISCUSSION

Cardiovascular diseases are the leading cause of mortality worldwide, with OxS, overweight, obesity and dyslipidaemias being the main risk factors. Many of the drugs aimed at reducing these risk factors often have collateral side effects when used, and many people often use complementary medicine instead. For this and other reasons, the pharmacological industry has focused on developing new treatment options.³⁴ *S. maxima* research has proven benefits to reduce CVD risk factors, alone or in combination with systematic practice of physical exercise, mainly in animal models.^{9 35 36}

The experimental design, procedures, compliance with ethical principles and statistical analyses of the study are summarised and described. Because of the double-blind design, the methodology described and a higher number of participants than other studies, it is expected to achieve better support for the hypothesis that *S. maxima* will have hypolipidaemic and antioxidant effects against intense exercise and obesity. It is noteworthy that not every overweight or obese patient has problems with lipid metabolism. Nowadays the association between the augmentation of body fat and lipid disturbances is well known.³⁷ For that reason, the primary focus of this protocol is prevention, which is why we do not consider dyslipidaemia as an inclusion criterion even though it could present in this condition.

Finally, it is known that the systematic practice of exercise and a balanced diet favours reduction of weight in obese people and generally decreases cardiometabolic risks,² so it probably generates an increase in antioxidant capacity.³⁸ However, no studies on the decrease in metabolic oxidation by the administration of *Spirulina* in obese people have been identified. Well-designed trials are needed to clarify the value of *S. maxima* supplementation in clinical practice³⁹ and its complementary effect with or without exercise against dyslipidaemias and OXS in overweight and obese people, a fact almost unknown at this time.⁴⁰ This establishes the importance of this study.

Author affiliations

¹Departamento de Ciencias Químico Biológicas, Instituto de Ciencias Biomédicas, Universidad Autónoma de Ciudad Juárez, Ciudad Juárez, Chihuahua, México

²Departamento de Ciencias de la Salud, Instituto de Ciencias Biomédicas, Universidad Autónoma de Ciudad Juárez, Ciudad Juárez, Chihuahua, México

³Departamento de Bioquímica, Facultad de Medicina, Universidad Nacional Autónoma de México, Ciudad de México, México

⁴Facultad de Ciencias de la Cultura Física, Universidad Autónoma de Chihuahua, Chihuahua, México

⁵Departamento de Biología, Facultad de Química, Universidad Nacional Autónoma de México, Ciudad de México, México

Correction notice This paper has been amended since it was published Online First. Owing to a scripting error, some of the publisher names in the references were replaced with 'BMJ Publishing Group'. This only affected the full text version, not the PDF. We have since corrected these errors and the correct publishers have been inserted into the references.

Acknowledgements We thank the Universidad Autonoma de Ciudad Juarez Institutional Bioethics Committee for the interest in this study, approving it and all its procedures. We also thank the Consejo Nacional de Ciencia y Tecnología (CONACyT) for providing a scholarship to tMAHL.

Contributors All authors contributed to the development of the study protocol and to the critical revision of the paper and approved the final version. RPHT, ARJ and JALD will be involved in patient recruitment; MAHL and AWM will analyse the data; MAHL and MAJO will do the lipid profile analyses; RUR, ARJ and MAHL will do the body composition measurements of all patients; LARC and JPC will do all the redox status analyses.

Competing interests None declared.

Patient consent Obtained.

Ethics approval Universidad Autonoma de Ciudad Juarez Bioethics Committee.

Provenance and peer review Not commissioned; externally peer reviewed.

Data sharing statement The protocol is an unpublished manuscript and is not under consideration by any other journal.

Open Access This is an Open Access article distributed in accordance with the Creative Commons Attribution Non Commercial (CC BY-NC 4.0) license, which permits others to distribute, remix, adapt, build upon this work non-commercially, and license their derivative works on different terms, provided the original work is properly cited and the use is non-commercial. See: <http://creativecommons.org/licenses/by-nc/4.0/>

© Article author(s) (or their employer(s) unless otherwise stated in the text of the article) 2017. All rights reserved. No commercial use is permitted unless otherwise expressly granted.

REFERENCES

1. World Health Organization. Factsheet No 317, 2015. <http://www.who.int/mediacentre/factsheets/fs317/en/>. (accessed 9 Nov 2015).
2. Genest J, Frohlich J, Fodor G, et al. Recommendations for the management of dyslipidemia and the prevention of cardiovascular disease: summary of the 2003 update. *CMAJ* 2003;169:921–4.
3. Ebbeling CB, Pawlak DB, Ludwig DS. Childhood obesity: public-health crisis, common sense cure. *Lancet* 2002;360:473–82.
4. Furukawa S, Fujita T, Shimabukuro M, et al. Increased oxidative stress in obesity and its impact on metabolic syndrome. *J Clin Invest* 2004;114:1752–61.
5. Li S, Chen W, Srinivasan SR, et al. Childhood cardiovascular risk factors and carotid vascular changes in adulthood: the Bogalusa Heart Study. *JAMA* 2003;290:2271–6.
6. Halliwell B, Cross CE. Oxygen-derived species: their relation to human disease and environmental stress. *Environ Health Perspect* 1994;102:5–12.
7. Schieber M, Chandel NS. ROS function in redox signaling and oxidative stress. *Curr Biol* 2014;24:R453–62.
8. Madamanchi NR, Vendrov A, Runge MS. Oxidative stress and vascular disease. *Arterioscler Thromb Vasc Biol* 2005;25:29–38.
9. Blé-Castillo JL, Rodríguez-Hernández A, Miranda-Zamora R, et al. Arthrospira maxima prevents the acute fatty liver induced by the administration of simvastatin, ethanol and a hypercholesterolemic diet to mice. *Life Sci* 2002;70:2665–73.
10. Estruch R, Ros E, Salas-Salvadó J, et al. Primary prevention of cardiovascular disease with a mediterranean diet. *N Engl J Med* 2013;368:1279–90.
11. Hosseini SM, Khosravi-Darani K, Mozafari MR. Nutritional and medical applications of spirulina microalgae. *Mini Rev Med Chem* 2013;13:1231–7.
12. Chamorro G, Salazar M, Favila L, et al. [Pharmacology and toxicology of Spirulina alga]. *Rev Invest Clin* 1996;48:389–99.
13. Iwata K, Inayama T, Kato T. Effects of spirulina platensis on plasma lipoprotein lipase activity in fructose-induced hyperlipidemic rats. *J Nutr Sci Vitaminol* 1990;36:165–71.
14. Riss J, Décorde K, Sutra T, et al. Phycobiliprotein C-phycocyanin from spirulina platensis is powerfully responsible for reducing oxidative stress and NADPH oxidase expression induced by an atherogenic diet in hamsters. *J Agric Food Chem* 2007;55:7962–7.
15. Torres-Duran PV, Ferreira-Hermosillo A, Juárez-Oropeza MA. Antihyperlipemic and antihypertensive effects of spirulina maxima in an open sample of Mexican population: a preliminary report. *Lipids Health Dis* 2007;6:33.
16. Plaza Pérez I, Villar Álvarez F, Mata López P, et al. Control de la colesterolemia en España, 2000. un instrumento para la prevención cardiovascular. *Rev Esp Cardiol* 2000;53:815–37.
17. Fisher-Wellman K, Bloomer RJ. Acute exercise and oxidative stress: a 30 year history. *Dyn Med* 2009;8:1.
18. Morillas-Ruiz JM, Villegas García JA, López FJ, et al. Effects of polyphenolic antioxidants on exercise-induced oxidative stress. *Clin Nutr* 2006;25:444–53.
19. Aoi W, Naito Y, Yoshikawa T. Role of oxidative stress in impaired insulin signaling associated with exercise-induced muscle damage. *Free Radic Biol Med* 2013;65:1265–72.
20. Gomez-Cabrera MC, Domenech E, Viña J. Moderate exercise is an antioxidant: upregulation of antioxidant genes by training. *Free Radic Biol Med* 2008;44:126–31.
21. Savini I, Catani MV, Evangelista D, et al. Obesity-associated oxidative stress: strategies finalized to improve redox state. *Int J Mol Sci* 2013;14:10497–538.
22. Gammone MA, Gemello E, Riccioni G, et al. Marine bioactives and potential application in sports. *Mar Drugs* 2014;12:2357–82.
23. Gammone MA, D'Orazio N. Anti-obesity activity of the marine carotenoid fucoxanthin. *Mar Drugs* 2015;13:2196–214.
24. Upasani CD, Khera A, Balaraman R. Effect of lead with vitamin E, C, or spirulina on malondialdehyde, conjugated dienes and hydroperoxides in rats. *Indian J Exp Biol* 2001;39:70–4.

25. Nagaoka S, Shimizu K, Kaneko H, *et al.* A novel protein C-phycocyanin plays a crucial role in the hypocholesterolemic action of *Spirulina platensis* concentrate in rats. *J Nutr* 2005;135:2425–30.
26. Thaakur SR, Jyothi B. Effect of spirulina maxima on the haloperidol induced tardive dyskinesia and oxidative stress in rats. *J Neural Transm* 2007;114:1217–25.
27. Stewart A, Marfell-Jones M, Olds T, *et al.* *International standards for anthropometric assessment*. Lower Hutt, New Zealand: ISAK, 2011.
28. Mao TK, Van de Water J, Gershwin ME. Effects of a spirulina-based dietary supplement on cytokine production from allergic rhinitis patients. *J Med Food* 2005;8:27–30.
29. Faul F, Erdfelder E, Lang AG, *et al.* G*Power 3: a flexible statistical power analysis program for the social, behavioral, and biomedical sciences. *Behav Res Methods* 2007;39:175–91.
30. Jiménez-Osorio AS, Picazo A, González-Reyes S, *et al.* Nrf2 and redox status in prediabetic and diabetic patients. *Int J Mol Sci* 2014;15:20290–305.
31. Koren E, Kohen R, Ginsburg I. Polyphenols enhance total oxidant-scavenging capacities of human blood by binding to red blood cells. *Exp Biol Med* 2010;235:689–99.
32. Prior RL, Hoang H, Gu L, *et al.* Assays for hydrophilic and lipophilic antioxidant capacity (Oxygen Radical Absorbance Capacity (ORAC(FL))) of plasma and other biological and food samples. *J Agric Food Chem* 2003;51:3273–9.
33. Pérez-Herrera N, May-Pech C, Hernández-Ochoa I, *et al.* PON1Q192R polymorphism is associated with lipid profile in Mexican men with mayan ascendancy. *Exp Mol Pathol* 2008;85:129–34.
34. Banach M, Stulc T, Dent R, *et al.* Statin non-adherence and residual cardiovascular risk: there is need for substantial improvement. *Int J Cardiol* 2016;225:184–96.
35. Cheong SH, Kim MY, Sok DE, *et al.* Spirulina prevents atherosclerosis by reducing hypercholesterolemia in rabbits fed a high-cholesterol diet. *J Nutr Sci Vitaminol* 2010;56:34–40.
36. Moura LP, Puga GM, Beck WR, *et al.* Exercise and spirulina control non-alcoholic hepatic steatosis and lipid profile in diabetic Wistar rats. *Lipids Health Dis* 2011;10:77.
37. Ugwuja E, Ogbonna N, Nwibo A, *et al.* Overweight and obesity, lipid profile and atherogenic indices among civil servants in Abakaliki, South Eastern Nigeria. *Ann Med Health Sci Res* 2013;3:13–18.
38. Fenster CP, Weinsier RL, Darley-Usmar VM, *et al.* Obesity, aerobic exercise, and vascular disease: the role of oxidant stress. *Obes Res* 2002;10:964–8.
39. Serban MC, Sahebkar A, Dragan S, *et al.* A systematic review and meta-analysis of the impact of Spirulina supplementation on plasma lipid concentrations. *Clin Nutr* 2016;35:842–51.
40. Hernández-Lepe MA, Wall-Medrano A, Juárez-Oropeza MA, *et al.* Spirulina y su efecto hipolipemiente y antioxidante en humanos: una revisión sistemática. *Nutr Hosp* 2015;32:494–500.