BMJ Open Respiratory syncytial virus: a systematic scientometric analysis of the global publication output and the gender distribution of publishing authors

Dörthe Brüggmann,^{1,2} Corinna Köster,² Doris Klingelhöfer,² Jan Bauer,² Daniela Ohlendorf,² Matthias Bundschuh,² David A Groneberg²

To cite: Brüggmann D, Köster C, Klingelhöfer D, *et al.* Respiratory syncytial virus: a systematic scientometric analysis of the global publication output and the gender distribution of publishing authors. *BMJ Open* 2017;7:e013615. doi:10.1136/bmjopen-2016-013615

Received 25 July 2016
Revised 1 March 2017
Accepted 8 March 2017

ABSTRACT

Objective Worldwide, the respiratory syncytial virus (RSV) represents the predominant viral agent causing bronchiolitis and pneumonia in children. To conduct research and tackle existing healthcare disparities, RSV-related research activities around the globe need to be described. Hence, we assessed the associated scientific output (represented by research articles) by geographical, chronological and socioeconomic criteria and analysed the authors publishing in the field by gender. Also, the 15 most cited articles and the most prolific journals were identified for RSV research.

Design Retrospective, descriptive study.

Setting The NewQIS (New Quality and Quantity Indices in Science) platform was employed to identify RSV-related articles published in the Web of Science until 2013. We performed a numerical analysis of all articles, and examined citation-based aspects (eg, citation rates); results were visualised by density equalising mapping tools.

Results We identified 4600 RSV-related articles. The USA led the field; US-American authors published 2139 articles (46.5% of all identified articles), which have been cited 83 000 times. When output was related to socioeconomic benchmarks such as gross domestic product or Research and Development expenditures, Guinea-Bissau, The Gambia and Chile were ranked in leading positions. A total of 614 articles on RSV (13.34% of all articles) were attributed to scientific collaborations. These were primarily established between high-income countries. The gender analysis indicated that male scientists dominated in all countries except Brazil.

Conclusions The majority of RSV-related research articles originated from high-income countries whereas developing nations showed only minimal publication productivity and were barely part of any collaborative networks. Hence, research capacity in these nations should be increased in order to assist in addressing inequities in resource allocation and the clinical burden of RSV in these countries.

INTRODUCTION

The human respiratory syncytial virus (RSV) is the predominant viral agent affecting the respiratory tract worldwide.^{1 2} It is

Strengths and limitations of this study

- ▶ This is the first concise depiction of the worldwide scientific productivity related to respiratory syncytial virus, which was assessed by geographical, chronological and socioeconomic criteria.
- ▶ The NewQIS (New Quality and Quantity Indices in Science) platform combines scientometric methods and 'density equalizing mapping projections' to evaluate the scientific output regarding quantitative and qualitative aspects, geographical and chronological developments, existing research networks and socioeconomic benchmarks in a reliable and standardised way.
- ▶ Since the Web of Science has a preference for English journals, we have to acknowledge a language bias associated with our analysis.
- ▶ Citation-based parameters were assessed, which have limitations since these rather reflect the recognition of the research in the scientific community than truly measure quality.

associated with 64 million infections that occur primarily in children under 5 years.³ RSV belongs to the Paramyxoviridae family and is a negative-sense, non-segmented, single-stranded RNA virus, which is mostly transmitted by droplets.⁴ RSV causes bronchiolitis, pneumonia, bronchitis and croup. It is linked to recurrent wheezing and paediatric asthma.^{5 6} There is no definitive treatment for RSV-related conditions. Although the development of effective and safe vaccines has remained unsuccessful to date, the variety of candidate vaccines is constantly growing in the last years.⁷ One prophylactic agent is commercially available, the neutralising anti-RSV antibody palivizumab. Its use is limited to preterm babies, chronic lung disease of prematurity and infants with congenital heart disease.⁸

RSV-associated morbidity and mortality depends on many factors such as the

CrossMark

¹Department of Obstetrics and Gynecology, Keck School of Medicine of USC, Los Angeles, California, USA

²Institute of Occupational Medicine, Social Medicine and Environmental Medicine, Goethe University Frankfurt, Frankfurt, Germany

Correspondence to

Dr Dörthe Brüggmann;
doerthe.brueggmann@med.usc.edu

geographic location, climate patterns, genetic susceptibility, socioeconomic factors and local virus strains.³ The percentage of children having contracted RSV by their second year of life approaches 100%.^{9,10} In the USA, over 2 million children aged 5 years and below need medical attention to treat their RSV infection every year.¹¹ On average, 0.3% of these children and 0.7% of infants younger than 6 months require hospitalisation.¹¹ The general RSV mortality rate in this demographic varies between 0% and 33%.¹⁰ Worldwide, up to 199 000 children die due to RSV infections. Ninety-nine per cent of these deaths occur in developing countries,³ hence RSV represents a substantial burden for community health in these nations. According to WHO field studies in 10 developing countries, RSV causes 70% of all acute respiratory tract infections in children below 5 years of age.¹² Here, the virus is identified in about 15%–40% of the hospitalised children with pneumonia or bronchiolitis.¹³ RSV epidemics occur during rainy seasons in tropical climates and during the winter months in temperate zones.⁴ Both virus strains, RSV-A and RSV-B, cocirculate during respective outbreaks. The A subtype is typically associated with more severe disease.¹⁴ Distinct genotypes of both strains (identified by the genetic classification of their G protein) are predominant in any given year. This pattern is highly flexible, varies by region and may shift to other prominent genotypes the following year.¹⁵ Epidemiological studies are dedicated to characterise the distribution of RSV strains and genotypes worldwide, often leading to the identification of new variants such as the RSV Thailand B strain.¹⁶

Although care of patients with RSV has been improved considerably and extensive studies have been launched to estimate viral spread and disease burden in developing countries, tremendous challenges still remain. In regard to the high prevalence, morbidity and mortality of RSV worldwide, we consider that further research and the implementation of related public health measures are crucial for future successes. However, global research funds are limited and their allocation becomes challenging. In this context, the assessment of the scientific performance is a prerequisite for the reasonable distribution of monetary support and the planning of future research endeavours by scientists interested in the field. Hence, we employed scientometric tools with the goals (1) to determine the RSV-related publication output among the extensive amount of biomedical publications and (2) to evaluate the scientific productivity of single countries in the framework of the global research landscape and in relation to socioeconomic and gender aspects. We also identified the 15 most cited landmark articles and the most prolific journals in the field of RSV research.

METHODS

Methodical platform

We used the NewQIS (New Quality and Quantity Indices in Science) platform to assess the worldwide RSV publication activity in a reliable and standardised way. This tool combines scientometric methods and ‘density equalizing mapping projections’ (DEMP)^{17–19} to evaluate the publication output regarding quantitative and qualitative aspects, geographical and chronological developments, existing research networks and socioeconomic benchmarks.

Density equalising mapping

DEMP is a state-of-the-art technique to visualise benchmarking processes by anamorphic maps.^{17,20} Based on the algorithm of Gastner and Newman, the size of each country was modified analogously to country-specific data on RSV research leading to a new geographic distribution of the global landscape.¹⁷

Data collection

For data collection, the Web of Science (WoS) Core Collection database (Thomson Reuters) was employed. The following search term was created: Title=(‘RSV’ OR ‘Respiratory Syncytial Virus’ OR ‘RS Virus*’) NOT Topic=(Rous Sarcoma). The Boolean operator NOT was used to exclude all publications concerning the Rous sarcoma (virus). We limited our search to original research articles. Following our protocol, the time frame was restricted from 1900 to 2013; publications in 2014 were not regarded due to incomplete data acquisition at the time the study was performed.

Data analysis

RSV-related articles were analysed regarding quantitative aspects such as the total number of publications, citation numbers, countries of article origin (defined as the ‘country where the institution is located each author, who worked on publishing the article, is affiliated’), institutions, languages, cited reference numbers and publication date. The number of authors publishing on RSV was only quantified for the years with 30 or more annual publications. Also, h-Indices, and the average citation number per item (citation rate, CR) were investigated. Regression analysis was used to investigate the chronological evolution of RSV research. We calculated the coefficient of determination (r^2) representing the slope of the growth in scientific output and citations.

Modified Hirsch-Index

The Hirsch-Index (h-Index) is a recognised semiquantitative proxy measure to assess the impact of one author’s research output on the scientific community.^{21,22} An h-Index of 12 indicates that out of 12 published papers each has been cited at least 12 times. In this study, we applied this concept to the RSV-specific research productivity of single countries and calculated a ‘modified country-specific’ h-Index.

Country-specific gender analysis

The proportionality of male and female researchers among authors publishing on RSV was analysed. Online name databases were used to identify the authors' genders.²³ If first names were not gender specific or quoted as initials, a manual search (using websites, corresponding addresses and social networks) was launched. We only evaluated countries where a minimum of 60 authors were affiliated and a minimum of 50% of authors were identified by gender to ensure a valid analysis. These thresholds were chosen arbitrarily based on previous studies.

Cooperation analysis

We identified all RSV-associated publications that were issued due to international collaborative efforts. The total count of collaborative items was related to the overall number of publications for each investigated country. In brief, if at least two authors originating from different institutions or countries (as identified by the affiliations in the article) contributed to one article, this publication was defined as a collaborative item. Publications with two or more authors affiliated to the same country were counted one time only towards the complete count of joint publications of this particular country. If an author had two affiliations, these were counted for every country mentioned in the affiliations. Connecting vectors visualised these cooperations; their width and shade of grey reflected the number of joint publications.

Journal analysis

The journals publishing on RSV were analysed regarding quantitative and qualitative aspects, for example, number of published RSV articles as well as citations these items received (citation rate, CR).

Analysis of economic key figures

Two quotients were calculated to assess the scientific output of a specific country: (1) in relation to the number of inhabitants (Q1); and (2) in relation to its economic power (as measured by the gross domestic product, GDP, Q2). Data regarding the population and GDP of investigated countries were obtained from 2012 from the *CIA World Factbook*.²⁴ We computed the quotients in the following way:

1. Articles/population index (Q1) = number of articles/population in millions
2. Articles/GDP index (Q2) = number of articles/GDP in billions

All countries were classified into high-income, upper-middle-income, lower-middle-income and low-income groups according to World Bank definitions.²⁵

We compared the total number of RSV articles to the gross domestic expenditure on Research and Development (R&D; in % of GDP) as well as to the number of researchers (per billion inhabitants) affiliated to the investigated countries.²⁶ The analysis was limited to

Figure 1 Chronological development of the number of articles.

countries that published a minimum of 30 articles in the field of RSV.

RESULTS

Number of published items

We identified 4600 articles on RSV published between 1900 and 2013; 2645 (57.5%) of these were issued after 2000. 1960 and onwards, the number of articles increased significantly over time as indicated by $r^2=0.9$ (figure 1). A total of 10 791 authors published in the field of RSV. The number of authors per article increased by 100% in the investigated time frame. We identified a mean of 3.8 and a median of 3 authors in 1978, which was the first year with more than 30 annual publications. Ten years later, a mean of 4.59 and a median of 4 authors were found, followed by a mean of 5.05 and a median of 5 authors in 2000, a mean of 5.95 and a median of 5 authors in 2006, and a mean of 7.6 and a median of 6 authors in 2013.

Analysis of research origin and citations

Although the publications originated from 92 of the 251 investigated countries and autonomous regions, the majority of the articles were written in English (4444 articles, 96.6% of all published RSV articles), followed by French (64 articles, 1.4% of all articles) and German (32 articles, 0.7% of all articles). More than 85% of the global publication output was issued by authors from high-income economies. The USA was the most productive nation (2139 articles, 46.5% of all published RSV articles), followed by the UK (583 articles, 12.7%), the Netherlands (231 articles, 5.0%), Canada (217 articles, 4.7%) and Germany (196 articles, 4.3%). Hence, the USA and UK dominated the cartogram, while major parts of Africa (with the exception of South Africa), Asia (with the exception of Japan, China, South Korea and India) and Central America occupied only minor areas (figure 2A).

The country-specific citation numbers and modified h-Indices showed a global distribution similar to the number of publications. Articles with US-American affiliation of the authors were cited most (83 000 citations, c),

Figure 2 Density equalising mapping projections. (A) Number of publications. (B) Modified h-Index.

followed by articles from the UK ($c=19240$), the Netherlands ($c=5587$), Canada ($c=5549$) and Germany ($c=5319$). Articles published by African, Asian and Middle American authors received hardly any citations.

The USA and the UK were the top-ranked countries with a modified h-Index of 121 (USA) and 68 (UK), followed by the Netherlands (h-Index=44), Germany (h-Index=43) and Canada (h-Index=40) (figure 2B).

Regarding the country-specific CR (average number of citations per total number of publications for each country with more than 30 articles, CR), Sweden (CR=40) dominated and was followed by the USA (CR=38.8), Finland (CR=34.9), the UK and Germany (CR=27.13).

Analysis of citation performance

The absolute citation count of all identified RSV-related articles resembled the growing volume of published papers in the investigated time frame. We documented a significant positive correlation between citation

numbers and the time of publication (with $r^2=0.72$ for the time frame from 1960 to 2008). The annual citation counts grew from 1960 to 1994 (eg, from 19 annual citations in 1960 to 2448 annual citations in 1994); after 1995, a rapid increase followed until 2003 (eg, 5274 annual citations in 1997 and 7790 annual citations in 2003) whereas a steep decline was noticed after 2006 (eg, 3147 annual citations in 2006). We identified visible peaks in citation activity for 1969 (2294 annual citations), 1986 (3898 annual citations), 2000 (7411 annual citations) and 2003 (7790 annual citations) (figure 3). Additionally, we compiled the 15 most-cited RSV articles, which constitute the publications that have sparked the most documented interest in the field to date (table 1).

Relation to economic parameters

Relating the number of publications to the population of a country, high-income countries such as Iceland ($Q1=34.59$), the Netherlands ($Q1=13.90$) and Denmark

Figure 3 Chronological development of annual citation numbers.

($Q1=12.25$) issued a higher number of publications per million inhabitants compared with low-income or lower-middle-income countries, for example, Nigeria ($Q1=0.03$) and Indonesia ($Q1=0.02$) (figure 4A). We identified two exceptions, which included the low-income countries Guinea-Bissau ($Q1=6.30$) and The Gambia ($Q1=4.76$). Both were represented among the 15 top-ranked countries in this analysis.

When looking at the publication activity in relation to GDP, two low-income countries outperformed high-income nations (figure 4B). Guinea-Bissau ($Q2=11.776$) and The Gambia ($Q2=8.721$) again held leading positions, followed by Iceland ($Q2=0.810$), Kenya ($Q2=0.516$) and Croatia ($Q2=0.319$). Besides these two exceptions, other low-income or lower-middle-income countries showed a weak performance (eg, Philippines with $Q2=0.004$ or Indonesia with $Q2=0.007$). Also, the USA did not remain under the top 10 nations and was found at position 18 ($Q2=0.132$).

Chile, the Netherlands, South Africa, UK and Argentina were leading the analysis when the total article count was related to the country-specific R&D expenditures (33.72, 13.32, 12.14, 11.46 and 9.56 articles per billion GDP spent on R&D, respectively). The USA dropped to position 11 with 4.52 articles per billion GDP spent on R&D (figure 5A, table 2).

When we related the article count to the number of researchers (per billion inhabitants) as a proxy measure for the active research community working on RSV and the productivity of these researchers, the USA was leading the field (532.27 articles per researcher per billion inhabitants), followed by the UK, China, Chile and South Africa (137.10, 83.55, 77.10 and 76.60 articles per number of researchers per billion inhabitants) (figure 5B, table 3).

Country-specific gender analysis

We identified the gender distribution among authors working on RSV in institutions affiliated to 92 countries. Our analysis indicated a larger proportion of male

scientists in almost all evaluated countries (figure 6). Brazil was an exception. Ninety-five authors with affiliation to Brazilian institutions were identified as working on RSV. Sixty-two of those were identifiable by name and gender. Here, men and women were almost equally represented (32 female authors, 51.6% of identifiable authors, 30 male authors, 48.4% of identifiable authors). By calculating the proportion of male to female scientists (m/f ratio), we documented the lowest ratio (m/f ratio=0.94) for Brazil, followed by Germany (m/f ratio=1.1). Japan had the highest score with m/f ratio=2.8.

International cooperation analysis

The first collaborative article on RSV was identified in 1973. It was published by researchers working in institutions located in Switzerland and the USA.²⁷ Since 1973, a total of 614 joint articles were published on RSV accounting for 13.34% of all articles. The USA was the preferred partner for international collaborations on RSV. The most productive cooperation was established between the USA and the UK (67 joint works, 10.9% of collaborative articles), followed by the USA and Canada (45 joint articles, 7.3% of collaborative articles), and the USA and Germany (34 joint articles, 5.5% of collaborative articles).

The most fruitful cooperation not involving the USA existed between the UK and Spain (33 joint articles, 5.4% of collaborative articles). The most productive cooperation between high-income and lower-middle-income or low-income economies was set up between Kenya and the UK (18 joint articles, 2.9% of collaborative articles), followed by Guinea-Bissau and Denmark (10 joint articles, 1.6% of collaborative articles). Overall, cooperations with low-income or lower-middle-income countries were less popular (figure 7).

Journal analysis

When we identified the most prolific journals in RSV research, the 'Journal of Virology' led the field (334

Table 1 The 15 most cited articles in the area of RSV-related research are displayed including their title, publication year, countries of article origin (defined as the countries where first, senior and coauthors are affiliated), citation count and journal

Title	Publication year	Country of article origin	Citations	Journal
Mortality associated with influenza and respiratory syncytial virus in the United States Thompson WW <i>et al</i>	2003	USA	1520	JAMA
Pattern recognition receptors TLR4 and CD14 mediate response to respiratory syncytial virus Kurt-Jones EA <i>et al</i>	2000	USA	856	Nat Immunol
Respiratory syncytial virus disease in infants despite prior administration of antigenic inactivated vaccine Kim HW <i>et al</i>	1969	USA	848	Am J Epidemiol
Respiratory syncytial virus in early life and risk of wheeze and allergy by age 13 years Stein RT <i>et al</i>	1999	Brazil, USA	719	Lancet
Palivizumab, a humanized respiratory syncytial virus monoclonal antibody, reduces hospitalization from respiratory syncytial virus infection in high-risk infants The Impact-RSV Study Group	1998	USA, Canada, UK	652	Pediatrics
An epidemiologic study of altered clinical reactivity to respiratory syncytial (RS) virus infection in children previously vaccinated with an inactivated RS virus vaccine Kapikian <i>et al</i>	1969	USA	552	Am J Epidemiol
Risk of primary infection and reinfection with respiratory syncytial virus Glezen <i>et al</i>	1986	USA	548	Am J Dis Child
Respiratory syncytial virus bronchiolitis in infancy is an important risk factor for asthma and allergy at age 7 Sigurs N <i>et al</i>	2000	Sweden, Iceland	488	Am J Respir Crit Care Med
Wheezing, asthma, and pulmonary dysfunction 10 years after infection with respiratory syncytial virus in infancy Pullan CR and Hey EN	1982	UK	449	Br Med J
Respiratory syncytial virus infection in elderly and high-risk adults Falsey AR <i>et al</i>	2005	USA	430	NEJM
Prophylactic administration of respiratory syncytial virus immune globulin to high-risk infants and young children Groothuis JR <i>et al</i>	1993	USA	427	NEJM
The development of respiratory syncytial virus-specific IgE and the release of histamine in nasopharyngeal secretions after infection Welliver RC <i>et al</i>	1981	USA	425	NEJM

Continued

Table 1 Continued					
Title	Publication year	Country of article origin	Citations	Journal	
Respiratory-syncytial-virus infections, re-infections and immunity—prospective, longitudinal study in young children Henderson FW <i>et al</i>	1979	USA	418	NEJM	
Generation of bovine respiratory syncytial virus (BRSV) from cDNA: BRSV NS2 is not essential for virus replication in tissue culture, and the human RSV leader region acts as a functional BRSV genome promoter Buchholz UJ <i>et al</i>	1999	Germany	417	J Virol	
Asthma and immunoglobulin-e antibodies after respiratory syncytial virus bronchiolitis—a prospective cohort study with matched controls Sigurs N <i>et al</i>	1995	Sweden	416	Pediatrics	

articles, CR of 45.49), followed by ‘The Pediatric Infectious Disease Journal’ (198 articles, CR=24.66) and the ‘Journal of Infectious Diseases’ (186 articles, CR 55.18) (figure 8). The highest CR was achieved by the ‘Proceedings of the National Academy of Sciences of the United States of America’ (42 articles, CR=81.43), followed by ‘Pediatrics’ (66 articles, CR=70.80), ‘Journal of Pediatrics’ (65 articles, CR=64.03), ‘American Journal of Respiratory Critical Care’ (47 articles, CR=62.55) and ‘Journal of Infectious Diseases’ (186 articles, CR=55.18).

DISCUSSION

In the WoS, we documented only 4600 RSV-related articles since 1900. The first item on the disease in humans was published in 1957.²⁸ This emphasises that research on RSV is a relatively new field considering the virus was initially isolated in 1956 from laboratory primates.²⁹ The predominance of English in the majority of identified articles aligns with the fact that this language is the recognised ‘scientific lingua franca’ and reflects the abundant research output of English-speaking countries such as the USA, Canada and the UK found in our study.

The increasing number of RSV publications over time is typical for most biomedical research, for example, on the John Cunningham virus, influenza or breast cancer.^{30 31} Overall, the steady growth of article numbers can be explained by the rising interest in the field due to the increasing relevance of RSV in pneumonia and child mortality.⁹ In the first few years after the detection of the virus, basic research was conducted. These endeavours, which aimed to characterise the virus, identify immunologic responses and develop vaccines, translated into a growing volume of articles on RSV.^{32–35} The continuous increase since the beginning of the 1990s may be attributed to the launch of a growing number of scientific journals providing a platform for publishing. The development of modern communication systems based on new computer technologies such as the World Wide Web made it easier to communicate, exchange ideas and publicise articles in central databases. Also, it is noticeable that the increasing publication output since the 1960s was paralleled by a globally growing funding volume allocated to the R&D sector. In the USA alone, a total of US\$13 711 million was allocated to R&D in 1960 as documented by the National Science Foundation. This amount increased to US\$26 271 million in 1970 and US\$452 556 million in 2012 (<https://www.nsf.gov/statistics/2015/nsf15315/>).

The chronological development in publication quantity (figure 1) was resembled by the steady increase of related citations (figure 3). Four prominent citation peaks in 1969, 1986, 2000 and 2003 coincided with milestone papers in the field. In 1969, adverse effects of the formalin-inactivated RSV vaccine in children were reported.³⁶ Large epidemiological studies investigated the risks of reinfection and the mortality associated with RSV in 1986 and 2003.^{37 38} The research on a prophylactic antibody

Figure 4 Density equalising mapping projections. (A) Articles/population index (Q1). (B) Articles/GDP index (Q2) (threshold ≥ 15 articles). GDP, gross domestic product.

licensed in 1999 and novel insights into immunologic responses involving pattern recognition receptors TLR4 and CD14 may be responsible for the peaking number of publications and citations in 2000.³⁹ The citation decline after 2006 can be attributed to the short timespan articles had allotted to receive recognition within the scientific community and obtain the appropriate citation number reflecting their true impact.⁴⁰ Hence, we expect this trend to be reversed in the future.

The USA dominated RSV research with regard to overall publication quantity, citation numbers and h-Index. This corresponds with a previous biomedical benchmarking study. Here, the USA was the most productive nation and authored 1893800 of 5527558 publications related to 22 organ systems from 1961 to 2007.⁴¹ The leading role of the USA might be linked to major financial resources

this nation dedicates to research. The US-American National Institutes of Health is by far the biggest biomedical funder in the world (eg, with a funding volume of US\$26.08 billion in 2013 compared with the biggest funding source in the European Union, the European Commission, with a funding volume of US\$3.71 billion in 2013). Also, the Department of Defense constitutes another large US-American funding organisation with a volume of US\$1.017 billion (2013), followed by private philanthropic institutions such as the Howard Hughes Medical Institute or the Bill & Melinda Gates Foundation with impressive funding volumes of US\$752.0 million (2013) and US\$462.6 million (2011), respectively.⁴² These funds can support manpower and an outstanding scientific infrastructure illustrated by the fact that the majority of institutions working on RSV were

Figure 5 Density equalising mapping projections. (A) Articles/R&D expenditure in billion USD index. (B) Articles/researcher (per billion inhabitants) index (threshold ≥ 30 articles). R&D, Research and Development; USD, US dollars.

identified in the USA, and that this nation is a preferred partner for national and international cooperations.

The cluster of the USA, Western-European countries (eg, UK or Germany) and Japan dominated the overall publication output and analyses of citation-based benchmarks. This finding corresponds with other scientometric studies (eg, on influenza, Ebola or hepatitis B).^{30 43} Although Sweden and Finland published relatively low numbers of articles, they are characterised by the highest CRs in our analysis, indicating the outstanding quality and high recognition their articles received in the scientific community. Furthermore, it was striking that most African, Asian and Central American countries afflicted with a considerable RSV-related burden did not play a

visible role in the field. Methodologically, we included only countries in the CR analysis that published more than 30 RSV articles aiming to generate a better validity of the investigation by avoiding overestimation of few but frequently cited articles. Nevertheless, we want to stress that the absolute number of citations as well as the CR should be viewed critically. These parameters can be affected by self-citation and inaccurate citations. Also, the Matthew effect might influence citation-based variables. Here, scientists prefer to cite articles issued by well-known researchers to papers by junior scientists leading to a disproportional increase in the related citation counts.^{44 45} Hence, we also evaluated the modified h-Index since it is less influenced by outstanding, frequently or rarely cited

Table 2 Total number of publications related to gross domestic expenditures on Research and Development in billion USD of countries that have published more than 30 items on RSV

Rank	Country	No. of articles	R&D expenditure in % GDP	GDP in billion USD	R&D expenditure in billion USD	Articles/billion USD of R&D expenditure
1	Chile	33	0.38	258	0.98	33.72
2	Netherlands	231	1.97	879	17.34	13.32
3	South Africa	31	0.73	349	2.55	12.14
4	UK	583	1.70	2990	50.85	11.46
5	Argentina	31	0.61	529	3.24	9.56
6	Spain	160	1.23	1381	17.01	9.41
7	Canada	217	1.61	1783	28.73	7.55
8	Denmark	68	3.08	346	10.67	6.37
9	Sweden	99	3.16	571	18.05	5.49
10	Finland	46	3.17	272	8.63	5.33
11	USA	2139	2.73	17 348	472.78	4.52
12	Belgium	56	2.46	531	13.09	4.28
13	Italy	84	1.29	2138	27.52	3.05
14	Switzerland	62	2.97	701	20.80	2.98
15	France	189	2.26	2829	63.82	2.96
16	Australia	86	2.20	1454	31.93	2.69
17	Israel	33	4.11	305	12.53	2.63
18	Brazil	75	1.24	2417	29.87	2.51
19	Germany	196	2.87	3868	110.96	1.77
20	Russia	37	1.19	2030	24.09	1.54
21	Japan	192	3.58	4596	164.73	1.17
22	South Korea	36	4.29	1411	60.55	0.59
23	China*	93	2.05	10 351	211.79	0.44

GDP, gross domestic product; RSV, respiratory syncytial virus; R&D, Research and Development; USD, US dollars. *China without Taiwan.

articles skewing the CR value.^{21 22} Furthermore, all citation-based variables have limitations in assessing the quality of the identified articles because they rather reflect the recognition of the research in the scientific community than measure quality.

We identified an over-representation of male authors in the majority of evaluated nations. This corresponds to the study of Head *et al* who documented the preferential funding of male researchers by UK institutions in the area of global infectious disease research. Between 1997 and 2010, funding agencies supported fewer studies of female PIs and awarded less monetary support to research supervised by women. Particularly for RSV, male researchers received five times more funding than female scientists, who spearheaded only half the funded studies compared with their male counterparts in the field.⁴⁶ In our gender analysis, Brazil was an exception with a majority of women authoring RSV research (figure 6). This result correlates with previous investigations on yellow fever and rotavirus infections⁴⁷ as well as with gender benchmarking studies (eg, conducted by the 'Konrad-Adenauer Foundation' or the 'Organization for Women in Science for the

Developing World'), which suggests that Brazil pioneers in the support and participation of women in science.^{48 49}

Our analysis of RSV publication outputs changed in relation to economic capabilities (figure 4). Two developing, low-income nations, The Gambia and Guinea-Bissau, occupied the leading positions. This finding points towards the fact that both prioritise RSV research and might be connected to existing cooperations with a long-standing shared history between collaborating nations and their focused support of RSV scientific activities. For example, the UK's 'Medical Research Council The Gambia Unit' and the Danish 'Bandim Health Project' encourage medical research in Guinea-Bissau and The Gambia.^{50 51} Also, research in the field is promoted by single researchers with a strong dedication to conduct research in African sites. Here, Sir Brian Greenwood has spearheaded RSV-related research very successfully for decades while being faculty at the London School of Hygiene and Tropical Medicine as well as the Director of the Medical Research Council in The Gambia.^{13 50–58}

If the RSV article counts were related to R&D expenditures and number of researchers in specific countries, a

Table 3 Total number of publications related to articles per researcher (per billion inhabitants) of countries that have published more than 30 items on RSV

Rank	Country	Number of RSV articles	Number of researchers per billion inhabitants	Articles/researcher (per billion inhabitants)
1	USA	2139	4.02	532.27
2	UK	583	4.25	137.10
3	China*	93	1.11	83.55
4	Chile	33	0.42	77.11
5	South Africa	31	0.40	76.60
6	Spain	160	2.64	60.55
7	Netherlands	231	4.48	51.58
8	Canada	217	4.52	48.02
9	France	189	4.20	44.99
10	Germany	196	4.38	44.74
11	Italy	84	2.01	41.86
12	Japan	192	5.39	35.65
13	Argentina	31	1.20	25.79
14	Sweden	99	6.87	14.41
15	Switzerland	62	4.48	13.83
16	Belgium	56	4.18	13.41
17	Russia	37	3.10	11.93
18	Denmark	68	7.20	9.45
19	Finland	46	7.00	6.58
20	South Korea	36	6.90	5.22
21	Israel	33	8.26	4.00

RSV, respiratory syncytial virus. *China without Taiwan.

different, more refined picture emerged compared with the assessments based on absolute publication numbers or related to socioeconomic variables. Here, two Latin American countries gained importance, Chile and Argentina. It

appears that these nations invest funding very efficiently in RSV research, with Chile ranked first position with 33.72 RSV articles per billion US dollars in R&D expenditures followed by Argentina in fifth position (9.56 articles

Figure 6 Country-specific gender analysis of the authors publishing articles referring to respiratory syncytial virus of countries (threshold: >50% definable genders, >60 authors per country).

per billion US dollars in R&D). Both nations' interest to fund RSV research might be linked to the fact that respiratory infections and RSV in particular impose a heavy burden on the local paediatric population. Respiratory infections constitute the second leading cause of death in Latin American children aged 5 years or below with RSV

as the causative agent in 70.0% of these infections.^{3 4 59 60} In the temperate climate of Chile and Argentina, RSV causes predictable outbreaks during the summer months. A particular high RSV burden of up to 70% was reported in Chilean children aged 0–11 months with lower respiratory tract infection. This was substantially higher than

in other Latin American countries (eg, 18.2% in Argentina and 44% in Brazil).⁶¹ Hence, RSV is constantly in the focus of the local health authorities, which routinely monitor and report the trends in RSV infections to better allocate resources for paediatric patients and limit related morbidity and mortality.^{61 62}

Collaborations are becoming increasingly important in the field of RSV research as indicated by existing tight-knit networks and the growing numbers of authors per article over time. We link this development to the globalisation process, which connects scientists worldwide to exchange ideas, resources and knowledge facilitated by the growing availability of information technology. Further, it is noticeable that countries such as the USA or European Nations play a more prominent role in international collaborations compared with low-income and lower-middle-income countries. As exceptions, we could identify productive cooperations between the UK and Kenya as well as Denmark and Guinea-Bissau. The relation between Kenya and the UK might be based on their shared history and facilitated by implemented programmes such as the ‘KEMRI Wellcome Trust Research Program’ between the Kenya Medical Research Institute and the University of Oxford.⁶³ As revealed by Fitchett *et al*,⁶⁴ a substantial funding volume goes to infectious disease research in countries with colonial ties to the UK such as Kenya and The Gambia. From 1997 to 2010, these countries received £13.13 million (The Gambia) and £12.92 million (Kenya) of biomedical funding by UK-based institutions. Research activity on RSV in Guinea-Bissau is also supported by the aforementioned Danish ‘Bandim Health Project,’ which was founded by the anthropologist Sir Peter Aaby in the 1970s. It gathers local epidemiological data on more than 200 000 individuals. Since its foundation, this group has published more than 600 items on vaccines, maternal mortality and childhood infections such as RSV.⁵⁰

Timing, intensity and clinical impact of RSV infections vary worldwide.³ Hence, research is still needed to alleviate the burden related to RSV in high-risk populations. Interventions should focus on data collection via established surveillance systems (eg, aiming to define local morbidity and mortality, assist disease modelling and guide prophylactic measures and vaccine development).⁶⁵ In this context, our study revealed a striking discrepancy in scientific productivity and collaborative involvement between high-income and low-income countries. Also, attention should be drawn to a further problem concerning low-income nations or countries whose researchers have a limited financial budget to pay for publication in renowned open access journals. This issue increases the apparent discrepancies regarding publication activities even further. To minimise this problem, a number of waiver programmes currently exist, that is, for journals such as *PLOS*, *Biomed Central* or *BMJ OPEN*,^{66–68} but these should be expanded more broadly. However, we can deduce from our findings that developing nations—although experiencing the most significant consequences of RSV epidemics—cannot

compete equally in the field of RSV research due to the lack of funding and infrastructure. Therefore, we want to emphasise the need—an almost ethical responsibility—to involve these nations in funding programmes and successful international collaborations as seen for Guinea-Bissau, Kenya and The Gambia. We acknowledge that the establishment of collaborations between high-income and low-income nations is challenging due to the lack of resources, manpower and funding opportunities, the political climate, cultural differences between the potential partners, and an unrealistic assessment of the local research capacity and resources.⁶⁹ Also, existing collaborations and funding streams need to be viewed critically since they should rather reflect local disease burden, apparent healthcare disparities and scientific capability than being allocated based on a shared language or history between countries (eg, guided by former colonial ties). Therefore, funding institutions should revise their policies appropriately.⁶⁴ Also, global investment surveillance systems need to be established such as the ‘Research Fairness Initiative’ led by Cohred to guide and monitor sustainable, transparent and effective partnerships in research (<http://rfi.cohred.org/origin-of-the-rfi/>). Nevertheless, tight-knit networks would be key for developing countries to participate in the international exchange of data, resources and knowledge, and to facilitate their involvement in high-quality research efforts despite an unequal starting point.

Study limitations

Our study has several limitations. Using the WoS to conduct this analysis is associated with an important strength and with a weakness of the study. The WoS enabled us to assess quantitative and qualitative aspects of the publication output related to RSV research. This is a unique feature allowing a multifaceted evaluation of the publication productivity. On the other hand, the WoS displays a strong preference for English journals. Therefore, not all articles ever published on RSV could be analysed here. However, we regard this bias as limited as the majority of high-quality data are commonly published in international journals indexed by the WoS and therefore definitely included in our search. Also, we acknowledge that not all eligible RSV publications were detected by the conducted ‘TITLE’ search. This approach was preferred to a ‘TOPIC’ search, which identifies the search term in the abstract and the keywords leading to a significant amount of off-topic publications compromising the validity of our data collection. We concentrated our study on original articles to focus on published ‘cutting edge research’ in the field of RSV. We acknowledge that this strategy narrowed down the focus on the topic since other publication types such as commentaries, reviews, case reports or meeting reports were not included. Further, we did not employ any additional platforms such as PubMed, Google Scholar or Scopus to collect bibliometric data because the management, organisation and the scope of data are slightly different among these databases. This

would affect triangulating, comparing and integrating data related to RSV research in a meaningful way.

We identified the first collaborative article on RSV in 1973, which would indicate that researchers from different countries did not work together on RSV before this point in time. This assumption is not necessarily true. In 1972, the WoS indexed author affiliations for the first time, which indicates that articles published in a joint effort before 1972 would not have been detected by our methodological approach. The evaluation of country-specific publication performance by gender is meaningful but should be evaluated critically. Not all first names were included in our analysis since some were gender neutral, not listed in name databases or displayed as initials. Therefore, the threshold of at least 60 publishing scientists and 50% gender definability was implemented to include only countries providing meaningful and valid data. Further, we identified 'Mortality associated with influenza and respiratory syncytial virus in the United States' by Thompson *et al* as the most cited journal article in the field of RSV. It received 1520 citations representing its outstanding recognition in the scientific community. The publication covers the topics RSV as well as influenza, so the limitation has to be mentioned that our computed approach did not differentiate if the article was cited in 'influenza' or 'RSV' papers. Employing a manual analysis, we found that 70.3% of citing articles can be attributed to influenza, only 15.8% to RSV, and 2.2% were covering both topics. Hence, the impact of this particular paper on the field of RSV must be considered as less than it initially appeared based on the citation count alone.

Conclusion

We evaluated the worldwide RSV-related research output and demonstrated large differences between high-income, middle-income or low-income nations regarding most scientometric variables. These discrepancies partly diminished when country-specific scientific activities were related to economic key measures; here, the leading position of the USA in science was challenged by other nations. Hence, calculating these quotients is beneficial for the comparison of countries with unequal conditions and different scientific infrastructures. However, we can deduce from our study that research efforts of middle-income or low-income nations have to be strengthened, for example, by the reduction of journal fees and inclusion in international collaborations, so apparent disparities can be minimised and higher mortality rates related to RSV in developing nations can be tackled successfully.

Acknowledgements We thank Cristian Scutaru for the development and provision of the NewQIS analysing tools. We also thank Mario Schwarzer, MD, for supporting the study and helpful discussions as well as Jenny M. Jaque, MD, for thoroughly editing our manuscript.

Contributors DB, CK, DK, DAG, DO, JB and MB have made substantial contributions to the conception and design of the study, acquisition of the study data and have been involved in drafting and revising the manuscript. All authors have read and approved the final manuscript.

Funding This research received no specific grant from any funding agency in the public, commercial or not-for-profit sectors.

Competing interests None declared.

Provenance and peer review Not commissioned; externally peer reviewed.

Data sharing statement Datasets of this study are available from the corresponding author upon request.

Open Access This is an Open Access article distributed in accordance with the Creative Commons Attribution Non Commercial (CC BY-NC 4.0) license, which permits others to distribute, remix, adapt, build upon this work non-commercially, and license their derivative works on different terms, provided the original work is properly cited and the use is non-commercial. See: <http://creativecommons.org/licenses/by-nc/4.0/>

© Article author(s) (or their employer(s) unless otherwise stated in the text of the article) 2017. All rights reserved. No commercial use is permitted unless otherwise expressly granted.

REFERENCES

- Murphy BR, Prince GA, Collins PL, *et al*. Current approaches to the development of vaccines effective against parainfluenza and respiratory syncytial viruses. *Virus Res* 1988;11:1–15.
- Collins PL, Graham BS. Viral and host factors in human respiratory syncytial virus pathogenesis. *J Virol* 2008;82:2040–55.
- Nair H, Nokes DJ, Gessner BD, *et al*. Global burden of acute lower respiratory infections due to respiratory syncytial virus in young children: a systematic review and meta-analysis. *Lancet* 2010;375:1545–55.
- Borchers AT, Chang C, Gershwin ME, *et al*. Respiratory syncytial virus—a comprehensive review. *Clin Rev Allergy Immunol* 2013;45:331–79.
- Blanken MO, Rovers MM, Molenaar JM, *et al*. Respiratory syncytial virus and recurrent wheeze in healthy preterm infants. *N Engl J Med* 2013;368:1791–9.
- Wu P, Dupont WD, Griffin MR, *et al*. Evidence of a causal role of winter virus infection during infancy in early childhood asthma. *Am J Respir Crit Care Med* 2008;178:1123–9.
- Graham BS. Vaccines against respiratory syncytial virus: the time has finally come. *Vaccine* 2016;34:3535–41.
- Polack FP. The changing landscape of respiratory syncytial virus. *Vaccine* 2015;33:6473–8.
- Centers for Disease Control and Prevention. Secondary respiratory syncytial virus infection (RSV). 2016 <http://www.cdc.gov/rsv/about/index.html>.
- Welliver RC, Checchia PA, Bauman JH, *et al*. Fatality rates in published reports of RSV hospitalizations among high-risk and otherwise healthy children. *Curr Med Res Opin* 2010;26:2175–81.
- Hall CB, Weinberg GA, Iwane MK, *et al*. The burden of respiratory syncytial virus infection in young children. *N Engl J Med* 2009;360:588–98.
- Selwyn BJ. The epidemiology of acute respiratory tract infection in young children: comparison of findings from several developing countries. Coordinated Data Group of BOSTID Researchers. *Rev Infect Dis* 1990;12(Suppl 8):S870–88.
- Weber MW, Mulholland EK, Greenwood BM. Respiratory syncytial virus infection in tropical and developing countries. *Trop Med Int Health* 1998;3:268–80.
- Walsh EE, McConnochie KM, Long CE, *et al*. Severity of respiratory syncytial virus infection is related to virus strain. *J Infect Dis* 1997;175:814–20.
- Peret TC, Hall CB, Schnabel KC, *et al*. Circulation patterns of genetically distinct group A and B strains of human respiratory syncytial virus in a community. *J Gen Virol* 1998;79:2221–9.
- Auksornkitti V, Kamprasert N, Thongkomplew S, *et al*. Molecular characterization of human respiratory syncytial virus, 2010–2011: identification of genotype ON1 and a new subgroup B genotype in Thailand. *Arch Virol* 2014;159:499–507.
- Gastner MT, Newman ME. From the Cover: diffusion-based method for producing density-equalizing maps. *Proc Natl Acad Sci USA* 2004;101:7499–504.
- Scutaru C, Quarcoo D, Sakr M, *et al*. Density-equalizing mapping and scientometric benchmarking of European allergy research. *J Occup Med Toxicol* 2010;5:2.
- Groneberg-Kloft B, Quarcoo D, Scutaru C. Quality and quantity indices in science: use of visualization tools. *EMBO Rep* 2009;10:800–3.
- Gerber A, Klingelhoefer D, Groneberg D, *et al*. Antineutrophil cytoplasmic antibody-associated vasculitides: a scientometric

- approach visualizing worldwide research activity. *Int J Rheum Dis* 2014;17:796–804.
21. Hirsch JE. An index to quantify an individual's scientific research output. *Proc Natl Acad Sci USA* 2005;102:16569–72.
 22. Hirsch JE. Does the H index have predictive power? *Proc Natl Acad Sci USA* 2007;104:19193–8.
 23. Namepedia. Secondary namepedia. 2016 <http://www.namepedia.org/>.
 24. World Economic Outlook Database. Secondary World Economic Outlook Database. 2013;2013 <http://www.imf.org/external/pubs/ft/weo/2013/02/weodata/weorept.aspx?pr.x=75&pr.y=10&sy=2012&ey=2012&scsm=1&ssd=1&sort=country&ds=.&br=1&c=193%2C223%2C924%2C132%2C134%2C146%2C136%2C158%2C112%2C111&s=NGDPD&grp=0&a=>.
 25. World Bank. Secondary country and lending groups. 2015 <http://data.worldbank.org/about/country-and-lending-groups>.
 26. OECD. Secondary main science and technology indicators. 2013 http://www.oecd.org/sti/2013_1_documentation_e.pdf.
 27. Bächli T, Howe C. Morphogenesis and ultrastructure of respiratory syncytial virus. *J Virol* 1973;12:1173–80.
 28. Chanock R, ROIZMAN B, Myers R. Recovery from infants with respiratory illness of a virus related to chimpanzee coryza agent (CCA). I. isolation, properties and characterization. *Am J Hyg* 1957;66:281–90.
 29. Blount RE, Morris JA, Savage RE. Recovery of cytopathogenic agent from chimpanzees with coryza. *Proc Soc Exp Biol Med* 1956;92:544–9.
 30. Fricke R, Uibel S, Klingelhoefer D, et al. Influenza: a scientometric and density-equalizing analysis. *BMC Infect Dis* 2013;13:454.
 31. Zheng HC, Yan L, Cui L, et al. Mapping the history and current situation of research on John Cunningham virus - a bibliometric analysis. *BMC Infect Dis* 2009;9:28.
 32. Simoes EA. Respiratory syncytial virus infection. *Lancet* 1999;354:847–52.
 33. Collins PL, Hill MG, Camargo E, et al. Production of infectious human respiratory syncytial virus from cloned cDNA confirms an essential role for the transcription elongation factor from the 5' proximal open reading frame of the M2 mRNA in gene expression and provides a capability for vaccine development. *Proc Natl Acad Sci USA* 1995;92:11563–7.
 34. Cranage MP, Gardner PS. Systemic cell-mediated and antibody responses in infants with respiratory syncytial virus infections. *J Med Virol* 1980;5:161–70.
 35. Fulginiti VA, Eller JJ, Sieber OF, et al. Respiratory virus immunization. I. A field trial of two inactivated respiratory virus vaccines; an aqueous trivalent parainfluenza virus vaccine and an alum-precipitated respiratory syncytial virus vaccine. *Am J Epidemiol* 1969;89:435–48.
 36. Kim HW, Canchola JG, Brandt CD, et al. Respiratory syncytial virus disease in infants despite prior administration of antigenic inactivated vaccine. *Am J Epidemiol* 1969;89:422–34.
 37. Glezen WP, Taber LH, Frank AL, et al. Risk of primary infection and reinfection with respiratory syncytial virus. *Am J Dis Child* 1986;140:543–6.
 38. Thompson WW, Shay DK, Weintraub E, et al. Mortality associated with influenza and respiratory syncytial virus in the United States. *JAMA* 2003;289:179–86.
 39. Kurt-Jones EA, Popova L, Kwinn L, et al. Pattern recognition receptors TLR4 and CD14 mediate response to respiratory syncytial virus. *Nat Immunol* 2000;1:398–401.
 40. Testa J. The Thomson Scientific journal selection process. *Int Microbiol* 2006;9:135–8.
 41. Groneberg-Kloft B, Scutaru C, Kreiter C, et al. Institutional operating figures in basic and applied sciences: scientometric analysis of quantitative output benchmarking. *Health Res Policy Syst* 2008;6:6:6.
 42. Viergever RF, Hendriks TC. The 10 largest public and philanthropic funders of health research in the world: what they fund and how they distribute their funds. *Health Res Policy Syst* 2016;14:12.
 43. Schmidt S, Bundschuh M, Scutaru C, et al. Hepatitis B: global scientific development from a critical point of view. *J Viral Hepat* 2014;21.
 44. Fassoulaki A, Paraskeva A, Papilas K, et al. Self-citations in six anaesthesia journals and their significance in determining the impact factor. *Br J Anaesth* 2000;84:266–9.
 45. Merton RK. The Matthew effect in science: the reward and communication systems of science are considered. *Science* 1968;159:56–63.
 46. Head MG, Fitchett JR, Cooke MK, et al. Differences in research funding for women scientists: a systematic comparison of UK investments in global infectious disease research during 1997–2010. *BMJ Open* 2013;3:e003362.
 47. Köster C, Klingelhoefer D, Groneberg DA, et al. Rotavirus - Global research density equalizing mapping and gender analysis. *Vaccine* 2016;34:90–100.
 48. Konrad-Adenauer-Stiftung. Secondary frauen in Brasilien. 2014 http://www.kas.de/wf/doc/kas_17800-1522-1-30.pdf?091024002708.
 49. Huyer S, Hafkin N. Scorecard on gender equality in the knowledge society. secondary scorecard on gender equality in the knowledge society. 2014 <http://www.elsevier.com/connect/brazilian-women-lead-in-science-technology-and-innovation-study-shows>.
 50. Bandim Health Project. About BHP. Secondary about BHP. <http://www.bandim.org/about-bhp.aspx>.
 51. Council. MR. Medical research council: the Gambia unit. The Gambia Unit: Secondary Medical Research Council. <http://www.mrc.gm/our-research/themes>.
 52. Adegbola RA, Falade AG, Sam BE, et al. The etiology of pneumonia in malnourished and well-nourished Gambian children. *Pediatr Infect Dis J* 1994;13:975–82.
 53. Suara RO, Piedra PA, Glezen WP, et al. Prevalence of neutralizing antibody to respiratory syncytial virus in sera from mothers and newborns residing in the Gambia and in the United States. *Clin Diagn Lab Immunol* 1996;3:477–9.
 54. Weber MW, Dackour R, Usen S, et al. The clinical spectrum of respiratory syncytial virus disease in the Gambia. *Pediatr Infect Dis J* 1998;17:224–30.
 55. Weber MW, Milligan P, Hilton S, et al. Risk factors for severe respiratory syncytial virus infection leading to hospital admission in children in the western region of the Gambia. *Int J Epidemiol* 1999;28:157–62.
 56. Weber MW, Milligan P, Sanneh M, et al. An epidemiological study of RSV infection in the Gambia. *Bull World Health Organ* 2002;80:562–8.
 57. Weber MW, Milligan P, Giadom B, et al. Respiratory illness after severe respiratory syncytial virus disease in infancy in the Gambia. *J Pediatr* 1999;135:683–8.
 58. Loscertales MP, Roca A, Ventura PJ, et al. Epidemiology and clinical presentation of respiratory syncytial virus infection in a rural area of southern Mozambique. *Pediatr Infect Dis J* 2002;21:148–55.
 59. Salomão Junior JB, Gardinassi LG, Simas PV, et al. Human respiratory syncytial virus in children hospitalized for acute lower respiratory infection. *J Pediatr* 2011;87:219.
 60. Piñeros JG, Baquero H, Bastidas J, et al. Respiratory syncytial virus infection as a cause of hospitalization in population under 1 year in Colombia. *J Pediatr* 2013;89:544–8.
 61. Bardach A, Rey-Ares L, Cafferata ML, et al. Systematic review and meta-analysis of respiratory syncytial virus infection epidemiology in Latin America. *Rev Med Virol* 2014;24:76–89.
 62. Avendaño LF, Palomino MA, Larrañaga C. Surveillance for respiratory syncytial virus in infants hospitalized for acute lower respiratory infection in Chile (1989 to 2000). *J Clin Microbiol* 2003;41:4879–82.
 63. KEMRI Wellcome Trust Research Programm. Secondary KEMRI Wellcome Trust Research Programm. <http://www.kemri-wellcome.org>.
 64. Fitchett JR, Head MG, Atun R. Infectious disease research investments follow colonial ties: questionable ethics. *Int Health* 2014;6:74–6.
 65. Haynes AK, Manangan AP, Iwane MK, et al. Respiratory syncytial virus circulation in seven countries with global disease detection Regional Centers. *J Infect Dis* 2013;208(Suppl 3):S246–54.
 66. PLOS. Secondary publication fees. <http://www.plos.org/publications/publication-fees>.
 67. Bio Med Central. Can charges be waived if the author cannot pay? Secondary can charges be waived if the author cannot pay? <http://www.biomedcentral.com/about/apcfaq/waivers>.
 68. BMJOpen. Secondary instructions for authors. <http://bmjopen.bmj.com/site/about/guidelines.xhtml>.
 69. Akinremi TO. Research collaboration with low resource countries: overcoming the challenges. *Infect Agent Cancer* 2011;6(Suppl 2):S3.