

# BMJ Open Suboptimal vitamin D status in Korean adolescents: a nationwide study on its prevalence, risk factors including cotinine-verified smoking status and association with atopic dermatitis and asthma

Eun Jung Byun,<sup>1</sup> Jinyoung Heo,<sup>2</sup> Sang Hyun Cho,<sup>1</sup> Jeong Deuk Lee,<sup>1</sup> Hei Sung Kim<sup>1</sup>

**To cite:** Byun EJ, Heo J, Cho SH, *et al.* Suboptimal vitamin D status in Korean adolescents: a nationwide study on its prevalence, risk factors including cotinine-verified smoking status and association with atopic dermatitis and asthma. *BMJ Open* 2017;7:e016409. doi:10.1136/bmjopen-2017-016409

► Prepublication history for this paper is available online. To view these files, please visit the journal online (<http://dx.doi.org/10.1136/bmjopen-2017-016409>).

Received 12 February 2017  
Revised 23 May 2017  
Accepted 25 May 2017



CrossMark

<sup>1</sup>Department of Dermatology, Incheon St. Mary's Hospital, College of Medicine, The Catholic University of Korea, Incheon, Republic of Korea

<sup>2</sup>Department of Preventive Medicine, The Catholic University of Korea, Seoul, Republic of Korea

#### Correspondence to

Prof. Hei Sung Kim;  
[hazelkimhoho@gmail.com](mailto:hazelkimhoho@gmail.com)

## ABSTRACT

**Objective** To determine the prevalence and risk factors for suboptimal vitamin D status in Korean adolescents and to assess its relationship with atopic dermatitis (AD) and asthma at a national level.

**Design** This is a cross-sectional study with data from the Korean National Health and Nutrition Examination Survey. Information regarding socioeconomic characteristics, clinical data and environmental factors was collected. Blood and urine samples were taken for vitamin D and cotinine, respectively. Descriptive and multivariable logistic regression was performed on the data.

**Setting** South Korea (nationwide).

**Participants** 2515 individuals aged 10–18 years who participated in the Korean National Health and Nutrition Examination Survey from 2008 to 2011.

**Main outcome measures** Vitamin D status was determined through measurement of serum 25-hydroxyvitamin D (25OHD). Smoking status was classified based on the urine cotinine level. Physician diagnosed with AD and asthma were assessed using a questionnaire.

**Results** Overall, 73.3% of the subjects were vitamin D deficient (25OHD <20 ng/mL) and 24.4% of the subjects were vitamin D insufficient (25OHD, 20–29.9 ng/mL). Older age ( $p<0.001$ ), female gender ( $p<0.001$ ), urban residence ( $p=0.019$ ), higher body mass index ( $p=0.003$ ) and sampling in winter months (November–March) ( $p<0.001$ ) were independently associated with low serum 25OHD levels. With cotinine verification, 18.2% of the participants were classified as active smokers, and 43.4% were classified as passive smokers. After adjusting for potential confounders, serum 25OHD status showed no association with AD or asthma.

**Conclusion** Vitamin D deficiency is highly prevalent in Korean adolescents. Cotinine-verified prevalence of smoking was also high, but its relationship with vitamin D deficiency was not confirmed in our study. Above all, our results provide epidemiological evidence against the association of vitamin D status with AD and asthma at the national level among Korean adolescents.

## Strengths and limitations of this study

- The strength of this study is that a nationally representative sample was used as the study population.
- This study is unique in that a cotinine-verified smoking status was adopted. Cotinine (a metabolite of nicotine) is a specific marker of smoking with high sensitivity.
- The limitation of the study is its cross-sectional design and the lack of data on the population's dietary and supplemental vitamin D intake. Also, the study has recall bias because the diagnoses of allergic conditions (atopic dermatitis (AD) and asthma) were self-reported.
- The definition of asthma and AD was physician-diagnosed asthma and AD which may have missed some children with symptomatic asthma and AD who have not been diagnosed.

## INTRODUCTION

Vitamin D is essential for skeletal health and bone growth and its deficiency has been associated with skeletal deformities in children and fracture risk in adults. Recent studies have also identified the previously unanticipated roles of vitamin D in the immune system, cardiovascular system and cancer prevention.<sup>1 2</sup> Despite growing health concerns over low vitamin D status, few nationwide studies have been performed to evaluate vitamin D deficiency in the Korean paediatric population.<sup>3 4</sup> Suboptimal vitamin D level was reported in 70% of the US children (9% vitamin D deficient, 61% vitamin D insufficient),<sup>5</sup> and it was found in 41.7% of the children from New Zealand.<sup>6</sup> Korean adolescents are theoretically at increased

risk for vitamin D deficiency because of the high latitude (34–38°N), culturally vigorous sun protection, reduced outdoor activity and lack of vitamin D-fortified food.

The increasing prevalence of allergic diseases is a worldwide phenomenon and it is strikingly more evident in the younger population compared with adults.<sup>7</sup> Vitamin D has immunomodulatory functions, and its relationship with allergic disease has been evaluated in a number of studies.<sup>8–15</sup> While some authors have reported about the protective role of vitamin D in atopic dermatitis (AD), asthma, allergic rhinitis and allergic sensitisation in childhood, other authors have shown a deleterious effect. Although the exact cause of such conflicting results is not known, racial difference may be a contributing factor. Unfortunately, nationwide studies on vitamin D and allergic disease are limited and are mostly from western countries.

In this study, we aimed to identify the prevalence and risk factors for vitamin D deficiency in Korean adolescents and to assess its relationship with AD and asthma at the national level. Cotinine-verified smoking status was adopted in this study.

## METHODS

### Study population

This study was based on data acquired from the Korean National Health and Nutrition Examination Survey (KNHANES), a survey conducted by the Korea Centers for Disease Control and Prevention to provide nationally representative and reliable statistical data regarding the health, behaviour associated with health, nutrition and food intake status of the Korean population. Data were collected from 2008 to 2011, which corresponds to the second and third year of KNHANES IV (2007–2009) and the first and second year of KNHANES V (2010–2012). The survey included a health interview, a nutritional survey, physical examination and blood and urine tests. The institutional review board at the Korea Centers for Disease Control and Prevention/Incheon St. Mary's Hospital, The Catholic University of Korea (IRB number: OC17ZESI0055) approved the protocol, and all participants and their parents signed informed consent forms.

Both KNHANES IV and V adopted the stratified multi-stage cluster sampling design by using the rolling-survey sampling method. The rolling sample collected each year is the probability sample representing the general Korean population, and it is homogeneous and independent of each other. In 2008, 2009, 2010 and 2011, a total of 37753 individuals were sampled. The study population was further limited to 4598 adolescents aged 10–18 years. Among the 4598 participants, we subsequently excluded the following participants: those whose serum 25-hydroxyvitamin D (25OHD) levels were not measured; those without urine cotinine levels; those who did not completely answer the questions regarding AD or asthma; those without body mass index (BMI) measurement and those who had a chronic disease that may affect vitamin

D metabolism. Finally, a total of 2515 participants (1314 males and 1201 females) were eligible for analysis.

### Study variables

Blood samples were collected from the antecubital vein, refrigerated immediately, transported to the central testing facility in cold storage and analysed within 24 hours of sampling. Serum 25OHD levels were measured as described previously<sup>16</sup> and categorised as sufficient ( $\geq 30$  ng/mL), insufficient (20–29.9 ng/mL) or deficient ( $< 20$  ng/mL).<sup>17</sup>

Factors were categorised to analyse the risk factors for vitamin D deficiency. Age and BMI were continuous variables while season of sampling was categorised into winter months (November–March) and summer months (April–October).<sup>18</sup> The region of residence of each participant was grouped as follows: urban (Seoul, Gyeonggi, Busan, Daegu, Incheon, Gwangju, Daejeon and Ulsan) and rural (Gangwon, Chungbuk, Chungnam, Jeonbuk, Jeonnam, Gyeongbuk, Gyeongnam and Jeju).<sup>19</sup> Monthly income was standardised according to the number of family members (monthly income/number of family members) and it was divided into the following four quartile groups: lowest, lower middle, higher middle and highest. Participants who performed moderate physical activity for more than 30 min per day on more than 5 days a week and/or strenuous physical activity for more than 20 min per day on more than 3 days a week were assigned to the regular exercise group. Regular walking was designated as 'yes' for those who walked for more than 30 min per day on more than 5 days a week.<sup>19</sup> Smoking status was divided into three groups based on the urine cotinine level, which are as follows: non-smokers ( $< 5$  ng/mL), passive smokers (secondhand smoking) (5–100 ng/mL) and active smokers ( $> 100$  ng/mL).<sup>19–21</sup> Urine cotinine level was measured by chromatography mass spectrometry using the Perkin Elmer Clarus 600T (PerkinElmer, Turku, Finland). All data were measured in a standardised manner and reviewed by the central quality control centre.

The following question was used to assess physician-diagnosed AD in each participant: "Have you been diagnosed with AD by a doctor?" or "Have you been told by a doctor that (your child) had AD?" Physician-diagnosed asthma was also determined using similar questions.

### Statistical analysis

Statistical analyses were performed using an SAS survey procedure V.9.2 (SAS Institute), and two-sided p values of less than 0.05 were considered statistically significant. To produce unbiased national estimates representing the general Korean population, we used KNHANES sample weights accounting for the complex sampling design to each participant.

To compare the mean serum 25OHD levels among categories of each possible predicting factor, Student's t-test or analysis of variance (ANOVA; followed by Tukey-Kramer for multiple comparison) was used.

Univariate analysis was performed to evaluate the association of the possible predicting factors, AD and asthma with vitamin D deficiency. Participants' characteristics were described using means and standard errors for continuous variables and numbers and percentages for categorical variables. Here, the Student's *t*-test or ANOVA was used for comparing continuous variables, as appropriate, and Rao-Scott  $\chi^2$  test was used for comparing categorical variables. Variables with a *p* value <0.05 in univariate analyses were included in the multivariate regression model for exploring factors associated with serum 25OHD.

To estimate the mean serum 25OHD levels in participants with and without AD and asthma, we performed simple and multiple linear regression analyses using the generalised linear model for a complex survey design. The estimated means were calculated as follows: no adjustment for potential confounders and confounder adjustment for age, gender, region, BMI, smoking and season of sampling.

To estimate the ORs for AD and asthma according to quartiles of serum 25OHD levels, we conducted simple and multivariate logistic regression analyses by using the generalised linear model for a complex survey design. The ORs and 95% CIs were calculated in the following ways: no adjustment for potential confounders and confounder adjustment for age, gender, region, BMI, smoking and season of sampling.

## RESULTS

### General characteristics

A total of 2515 subjects (1314 males and 1201 females, age: 14.4±0.1 years) were included in the study. Table 1 summarises the baseline data of the participants. The mean BMI of the study population was 20.9 kg/m<sup>2</sup> (SE: 0.1). Blood samples were drawn more frequently in April–October (58.4%) than in November–March (41.6%). The urine cotinine-verified smoking statuses were as follows: active smoking 18.2%, passive smoking 43.4% and non-smoking 38.4%. The prevalence of AD and asthma in the study population were 10.6% and 4.1%, respectively.

### Serum 25OHD levels in the study population

The mean concentration of serum 25OHD in the 2515 subjects was 16.7 ng/mL (SE: 0.2), with a range of 3.0 to 46.2 ng/mL. Overall, 1843 subjects (73.3%) were vitamin D deficient, 613 subjects (24.4%) were vitamin D insufficient and 59 subjects (2.3%) were vitamin sufficient (figure 1). Girls had a significantly lower mean serum 25OHD level than boys (16.0±0.2 ng/mL vs 17.3±0.2 ng/mL, *p*<0.001), and subjects sampled in winter months (November–March) had lower serum 25OHD levels than those sampled in summer months (April–October) (14.7±0.2 ng/mL vs 18.1±0.2 ng/mL, *p*<0.001). Passive smoking (vs non-smoking) and urban residence (vs rural) were also associated with statistically lower serum 25OHD levels (*p*=0.038 and *p*=0.002, respectively) (table 2).

### Univariate and multivariate analyses of factors associated with serum 25OHD

Univariate analysis revealed that age (*p*<0.001), gender (*p*=0.003), region of residence (*p*=0.011), season of sampling (*p*<0.001), BMI (<0.001) and smoking status (*p*=0.008) were associated with vitamin D deficiency (table 1).

Multivariate linear regression analysis of serum 25OHD levels was performed using variables that had a *p* value <0.05 in univariate analyses. Older age (*p*<0.001), female gender (*p*<0.001), urban residence (*p*=0.019), higher BMI (*p*=0.003) and sampling in winter months (November–March) (*p*<0.001) were independently associated with low serum 25OHD levels (table 3). Multivariate ordinal logistic regression analysis of serum 25OHD showed similar results (table 4).

### Relationship of serum 25OHD with AD and asthma

Univariate analysis demonstrated that vitamin D deficiency was not significantly associated with AD (*p*=0.468) or asthma (*p*=0.538) (table 1).

Multivariate logistic regression showed that vitamin D deficiency had no association with AD and asthma after adjusting for age, gender, region of residence, BMI, smoking and season of sampling. The adjusted OR for AD and asthma increased across categories of serum 25OHD (1.00 (reference) for ≥30 ng/mL, 1.14 and 4.87 each for 20–29.9 ng/mL, and 1.52 and 5.31, respectively for <20 ng/mL), but it was not statistically significant (*p*=0.8078 (AD) and *p*=0.1360 (asthma) for 20–29.9 ng/mL and *p*=0.4271 (AD) and *p*=0.1042 (asthma) for <20 ng/mL) (table 5).

Univariate or multivariate analyses with serum 25OHD levels as a continuous variable showed similar results, regardless of the adjustment (table 6).

## DISCUSSION

In our cross-sectional study of Korean adolescents, Vitamin D deficiency was very common, where the mean concentration of serum 25OHD was 16.7 ng/mL with 73.3% of the subjects being vitamin D deficient (serum 25OHD <20 ng/mL). Potential explanations for this remarkably low vitamin D status observed in Korean children include, but are not limited to, pigmented skin (skin pigmentation is known to reduce the skin's production of vitamin D in East Asian populations),<sup>3</sup> vigorous use of sunscreens and possibly dietary factors. The Dietary Reference Intakes for vitamin D from the Institute of Medicine (2010) and the American Academy of Pediatrics state that the adequate intake (AI) of vitamin D is 15 µg/day for children.<sup>22 23</sup> However, in the Dietary Reference Intakes for vitamin D for Koreans published in 2010, the AI of vitamin D for children is claimed to be 5 µg/day,<sup>24</sup> which we feel is too low. Exposure to sunlight, specifically ultraviolet B (UVB), leads to vitamin D synthesis in the skin, which provides most of the vitamin D requirement of an individual. In Korea (latitude: 34–38°N), vitamin D is

Table 1 Characteristics of the study population


	Vitamin D level				p Value
	Total (n=2515)	Deficiency (<20 ng/mL)	Insufficiency (≥20 to <30 ng/mL)	Sufficiency (≥30 ng/mL)	
Age	14.39±0.07	14.66±0.08	13.46±0.15	14.07±0.52	<0.0001
Sex					
Male	1314 (53.39)	914 (51.20)	359 (59.74)	41 (71.46)	0.0031
Female	1201 (46.61)	929 (37.45)	254 (40.26)	18 (28.54)	
Region of residence					
Urban	1709 (71.98)	1297 (73.59)	379 (67.94)	33 (50.84)	0.0108
Rural	806 (28.02)	546 (26.41)	234 (32.06)	26 (49.16)	
Regular exercise					
No	1861 (73.83)	1375 (74.94)	449 (70.20)	37 (69.98)	0.1912
Yes	654 (26.17)	468 (25.06)	164 (29.80)	22 (30.02)	
Regular walking					
No	1354 (53.23)	981 (52.98)	339 (54.37)	34 (50.44)	0.8708
Yes	1161 (46.77)	862 (47.02)	274 (45.63)	25 (49.56)	
Income					
Lowest	306 (13.87)	229 (14.33)	71 (12.78)	6 (6.83)	0.0996
Lower middle	604 (27.04)	452 (27.66)	133 (24.41)	19 (31.94)	
Higher middle	803 (30.39)	580 (30.92)	210 (29.54)	13 (17.44)	
Highest	802 (28.71)	582 (27.09)	199 (33.27)	21 (43.79)	
Season					
April–October	1465 (58.38)	930 (51.36)	485 (81.49)	50 (82.62)	<0.0001
November–March	1050 (41.62)	913 (48.64)	128 (18.51)	9 (17.38)	
BMI	20.88±0.10	21.12±0.13	20.08±0.19	20.12±0.47	<0.0001
Smoking					
Active	382 (18.22)	293 (19.08)	75 (13.99)	14 (32.22)	0.0075
Passive	1200 (43.39)	913 (44.36)	261 (39.98)	26 (42.56)	
No	933 (38.40)	637 (36.56)	277 (46.03)	19 (25.22)	
Atopic dermatitis					
No	2247 (89.44)	1648 (88.99)	546 (90.83)	53 (92.28)	0.4675
Yes	268 (10.56)	195 (11.01)	67 (9.17)	6 (7.72)	
Asthma					
No	2412 (95.87)	1771 (95.85)	583 (95.68)	58 (99.26)	0.5376

Continued

**Table 1** Continued

Vitamin D level					
	Total (n=2515)	Deficiency (<20 ng/mL)	Insufficiency (≥20 to <30 ng/mL)	Sufficiency (≥30 ng/mL)	p Value
Yes	103 (4.13)	72 (4.15)	30 (4.32)	1 (0.74)	

Univariate and multivariate analyses of factors associated with serum 25(OH)D. Data are presented as mean±SE, n (weighted %). Statistics were carried out using Rao-Scott  $\chi^2$  test. 25OHD, 25-hydroxyvitamin D; BMI, body mass index.



**Figure 1** Histogram of serum 25-hydroxyvitamin D (25OHD) levels.

synthesised mostly between April and October (summer months),<sup>25</sup> which was apparent in our study.

The finding of an inverse relationship of age, female gender, urban residence and BMI with vitamin D status is in accordance with previous studies.<sup>3 4 26–28</sup> It has been previously shown that there is a consistent decline in physical outdoor activity over the school age years. The prevalence of physical inactivity is higher in girls than in boys, and it is greater in obese children than in their normal-weight peers. The lifestyle in big cities is also sedentary, and it is therefore possible that age, gender, place of residence and BMI may act as surrogate indicators for sunlight exposure through their association with physical activity. In addition to limited sunlight exposure from physical inactivity, the inverse association of BMI and urban residence with vitamin D levels can be explained by sequestration of vitamin D in fat tissues<sup>29</sup> and UVB blockade due to air pollution.

This nationwide study is particularly meaningful as it is the first to measure both serum 25OHD and urine cotinine in a population of 2515 Korean adolescents. Smoking has been quoted as a significant determinant of serum 25OHD in a number of studies,<sup>20 21 28 30</sup> including one from Taiwan,<sup>28</sup> where passive smoking was independently associated with low serum 25OHD. To identify the effect of smoking on serum 25OHD in Koreans, we adopted urine cotinine, which is presently the biomarker of choice for assessing tobacco smoke exposure.<sup>31–34</sup> KNHANES normally assesses a person’s smoking status via a survey. For those who are aged 19 years and above, detailed questions are asked regarding the presence and history of active and passive smoking, but for those who are aged less than 19 years, the questions are much more limited. There is currently no data with regards to passive smoking in Korean adolescents and the survey only determines the presence of active smoking in subjects aged between 12 and 18 years. Omissions and false responses are common in this age group<sup>35</sup> which impede accurate assessment of the smoking status, and to improve accuracy, we adopted the cotinine-verified smoking status in

**Table 2** Vitamin D levels in the study population

	Vitamin D	p Value*	p Value†
Total	16.68±0.17		
Sex			
Male	17.26±0.22	<0.0001	
Female	16.01±0.21		
Region of residence			
Urban	16.34±0.20	0.0024	
Rural	17.53±0.34		
Regular exercise			
No	16.52±0.18	0.0565	
Yes	17.13±0.31		
Regular walking			
No	16.62±0.21	0.687	
Yes	16.74±0.24		
Income			
Lowest	16.09±0.41	0.0043	Lowest versus highest: 0.0218
Lower middle	16.63±0.29		Lower middle versus highest: 0.1576
Higher middle	16.47±0.24		Higher middle versus highest: 0.0449
Highest	17.22±0.31		
Season			
April–October	18.11±0.21	<0.0001	
November–March	14.67±0.21		
Smoking			
Active	16.52±0.42	0.0062	Active versus passive: 0.9439
Passive	16.37±0.23		Active versus no: 0.4260
No	17.10±0.23		Passive versus no: 0.0382
Atopic dermatitis			
No	16.68±0.18	0.8887	
Yes	16.63±0.32		
Asthma			
No	16.70±0.18	0.4507	
Yes	16.22±0.62		

Data are presented as mean±SE, n (weighted %).

\*p Value, t-test or analysis of variance (variables: income, smoking) result.

†p Value, multiple comparison: Tukey-Kramer.

**Table 3** Multivariate linear regression analyses of serum 25OHD levels

	Coefficient	95% CI	p Value
Age	-0.52	-0.65 to -0.40	<0.0001
Sex			
Male	Reference		<0.0001
Female	-1.40	-1.88 to 0.91	
Region of residence			
Urban	-0.85	-1.55 to 0.14	0.0189
Rural	Reference		
BMI	-0.10	-0.17 to 0.03	0.0034
Smoking			
Active	0.47	-0.42 to 1.36	0.3017
Passive	-0.04	-0.62 to 0.53	0.8842
No	Reference		
Season			
April–October	Reference		<0.0001
November–March	-3.38	-3.97 to 2.79	

Statistics were carried out using multivariable linear regression. Model: adjusted for age, sex, region, BMI, smoking, season. 25OHD, 25-hydroxyvitamin D; BMI, body mass index.

**Table 4** Multivariate ordinal logistic regression analyses of categories of serum 25OHD levels

	OR	95% CI	p Value
Age	1.22	1.13 to 1.31	<0.0001
Sex			
Male	Reference		0.0002
Female	1.68	1.28 to 2.21	
Region of residence			
Urban	1.32	0.96 to 1.81	0.0890
Rural	Reference		
BMI	1.07	1.03 to 1.12	0.0011
Smoking			
Active	0.89	0.54 to 1.47	0.6539
Passive	1.01	0.74 to 1.37	0.9535
No	Reference		
Season			
April–October	Reference		
November–March	4.49	3.25 to 6.22	<0.0001

Data are presented OR (95% CI).

Statistics were carried out using logistic regression.

Serum 25OHD levels were categorised into three ordinal categories of vitamin D status (<20: deficiency, ≥20 to <30: insufficiency, 30≥: sufficiency).

Model: adjusted for age, sex, region, BMI, smoking, season. 25OHD, 25-hydroxyvitamin D; BMI, body mass index.

**Table 5** ORs and 95% CIs of AD and asthma according to serum 25OHD levels

		Unadjusted		Adjusted	
		OR (95% CI)	p Value	OR (95% CI)	p Value
AD	Vitamin D Level				
	Sufficiency	Reference		Reference	
	Insufficiency	1.21 (0.43 to 3.39)	0.7228	1.14 (0.41 to 3.19)	0.8078
	Deficiency	1.48 (0.54 to 4.02)	0.4441	1.52 (0.54 to 4.22)	0.4271
Asthma	Vitamin D Level				
	Sufficiency	Reference		Reference	
	Insufficiency	6.02 (0.75 to 48.25)	0.0908	4.87 (0.61 to 39.05)	0.1360
	Deficiency	5.79 (0.76 to 43.96)	0.0898	5.31 (0.71 to 39.86)	0.1042

Data are presented OR (95% CI).

Statistics were carried out using logistic regression.

Adjusted for age, sex, region, BMI, smoking, season.

25OHD, 25-hydroxyvitamin D; AD, atopic dermatitis; BMI, body mass index.

our study. Urine cotinine level was measured for a limited time period (2008–2011), and this measurement has only been performed for those aged 10 years and above, which resulted in an inevitable reduction of our study population. We used a urine cotinine cut-off value of more than 100 ng/mL to discriminate smokers from non-smokers and 5 ng/mL as a threshold for non-smokers exposed to secondhand smoking based on published values.<sup>31–33</sup>

With cotinine verification, 18.2% of the participants were classified as active smokers, 43.4% of the participants were classified as passive smokers, and 38.4% of the participants were classified as non-smokers, where the percentage of smokers (active and passive) was higher than what we expected.

Smokers have been reported to have a significantly higher risk of vitamin D deficiency,<sup>20 21 28 30</sup> but this was not confirmed in our study. Since active smoking is usually done outdoors, this can result in greater exposure to sunlight which may alter the true effect of smoking on serum 25OHD. Passive smoking in adolescents is more likely to occur indoors and interestingly enough, passive smoking was reported to be independently associated with

low serum 25OHD in Taiwanese children/adolescents.<sup>28</sup> However, results can also be influenced by cotinine cut-off values and higher thresholds such as 10 or 30 ng/mL may be considered in the future when testing children as this population can have higher cotinine concentrations than adults due to differences in body distribution and nicotine metabolism.<sup>36</sup>

Some studies have indicated that an adequate concentration of vitamin D is protective against allergic disease in children.<sup>8 9 15 37 38</sup> However, most of these findings have been reported in western countries and only a few studies have been performed in Asia. We consider our investigation meaningful because differences among ethnicities and subjects of various age groups can possibly affect the results.

In this national level study, the risk of AD and asthma were not dependent on serum 25OHD. The present study results, as well as prior studies which showed little association between vitamin D deficiency and AD,<sup>10–12 28</sup> raise questions on the suggestion that vitamin D may be used to prevent or treat AD. In fact, a Cochrane review found no evidence for an effect of vitamin D supplementation

**Table 6** The estimated mean serum 25OHD levels and their differences according to AD and asthma

		Unadjusted			Adjusted		
		Estimated mean	Difference (estimated mean)	p Value	Estimated mean	Difference (estimated mean)	p Value
AD	Yes	16.68±0.18	0.05±0.34	0.8887	16.87±0.16	0.19±0.33	0.5776
	No	16.63±0.32			16.68±0.32		
Asthma	Yes	16.70±0.18	0.48±0.63	0.4507	16.88±0.15	0.72±0.48	0.1313
	No	16.22±0.62			16.15±0.47		

Data are presented as numeric, mean±SD.

Statistics were carried out using simple linear regression and multivariable linear regression.

Adjusted for age, sex, region, BMI, smoking, season.

25OHD, 25-hydroxyvitamin D; AD, atopic dermatitis; BMI, body mass index.

on AD.<sup>39</sup> The beneficial effect of sunlight on AD has been well documented, but this is probably due to the anti-inflammatory and antimicrobial effects of UV and not those of vitamin D.

As in AD, there has been considerable controversy over the relationship between vitamin D levels and asthma in children. While some studies showed an inverse relationship between vitamin D levels and current wheeze and asthma severity,<sup>8</sup> others such as Hollams *et al*<sup>40</sup> showed no significant cross-sectional association between serum vitamin D levels and current asthma. Our study too did not find any relationship between vitamin D level and asthma, but further studies would be needed to determine a causal relationship and its mechanism.

This study has several notable strengths. The sampling of adolescents across a broad age range, a large sample size with robust data collection, incorporation of objective markers of smoking status, and a thorough analysis strengthen the results of this study. Limitations of the study include its cross-sectional design and lack of data on dietary and supplemental vitamin D intake. Also, the study is prone to recall bias because diagnoses of allergic conditions (AD and asthma) were self-reported. The definition of asthma and AD were physician-diagnosed asthma and AD, which may have missed some children with symptomatic asthma and AD in the population who have not been diagnosed. Although season adjusted, deseasonalisation of vitamin D levels with an algorithm<sup>41</sup> may have further increased the accuracy and strength of our study.

In conclusion, a high prevalence of vitamin D deficiency was noted in Korean adolescents aged 10 to 18 years. For high-risk individuals of vitamin D deficiency, we suggest vitamin D supplementation and food fortification. Adequate outdoor activity should also be emphasised in Korean adolescents, especially high school students, via media promotion and public health plans by schools and government. We found no apparent association between the cotinine-verified smoking status, AD and asthma with vitamin D deficiency, which merits further study.

**Contributors** EJB, SHC, JDL and HSK made substantial contributions to the conception or design of the work. JYH and HSK had role in the acquisition, analysis and interpretation of data for the work. EJB, SHC, JDL and HSK drafted and revised the work and all authors gave final approval and agreed to be accountable for all aspects of the work.

**Funding** This study was supported by a grant from the Korean Healthcare Technology R&D project, Ministry of Health and Welfare, Republic of Korea (grant no HN15C0105).

**Disclaimer** The funders of this study had no role in study design, collection, analysis and interpretation of data, writing of the manuscript or decision to submit for publication.

**Competing interests** None declared.

**Patient consent** Detail has been removed from this case description/these case descriptions to ensure anonymity. The editors and reviewers have seen the detailed information available and are satisfied that the information backs up the case the authors are making.

**Ethics approval** The institutional review board at the Korea Centers for Disease Control and Prevention/Incheon St. Mary's Hospital, The Catholic University of Korea (IRB number: OC17ZESI0055) approved the protocol.

**Provenance and peer review** Not commissioned; externally peer reviewed.

**Data sharing statement** All data from the study, published and unpublished, are available with the principle investigators. The data are managed under the Department of Dermatology, Incheon St. Mary's Hospital, The Catholic University of Korea, Incheon, Korea.

**Open Access** This is an Open Access article distributed in accordance with the Creative Commons Attribution Non Commercial (CC BY-NC 4.0) license, which permits others to distribute, remix, adapt, build upon this work non-commercially, and license their derivative works on different terms, provided the original work is properly cited and the use is non-commercial. See: <http://creativecommons.org/licenses/by-nc/4.0/>

© Article author(s) (or their employer(s) unless otherwise stated in the text of the article) 2017. All rights reserved. No commercial use is permitted unless otherwise expressly granted.

## REFERENCES

- Holick MF. Vitamin D deficiency. *N Engl J Med Overseas Ed* 2007;357:266–81.
- Cheng HM, Kim S, Park GH, *et al*. Low vitamin D levels are associated with atopic dermatitis, but not allergic rhinitis, asthma, or IgE sensitization, in the adult Korean population. *J Allergy Clin Immunol* 2014;133:1048–55.
- Kim SH, Oh MK, Namgung R, *et al*. Prevalence of 25-hydroxyvitamin D deficiency in Korean adolescents: association with age, season and parental vitamin D status. *Public Health Nutr* 2014;17:122–30.
- Lee YA, Kim HY, Hong H, *et al*. Risk factors for low vitamin D status in Korean adolescents: the Korea National Health and Nutrition Examination Survey (KNHANES) 2008–2009. *Public Health Nutr* 2014;17:764–71.
- Kumar J, Muntner P, Kaskel FJ, *et al*. Prevalence and associations of 25-hydroxyvitamin D deficiency in US children: NHANES 2001–2004. *Pediatrics* 2009;124:e362–70.
- Rockell JE, Green TJ, Skeaff CM, *et al*. Season and ethnicity are determinants of serum 25-hydroxyvitamin D concentrations in New Zealand children aged 5–14 y. *J Nutr* 2005;135:2602–8.
- Asher MI, Montefort S, Bjorksten B, *et al*. Worldwide time trends in the prevalence of symptoms of asthma, allergic rhino-conjunctivitis, and eczema in childhood: ISAAC Phases one and three repeat multi-country cross-sectional surveys. *Lancet* 2006;26:733–43.
- Bener A, Ehlayel MS, Tulic MK, *et al*. Vitamin D deficiency as a strong predictor of asthma in children. *Int Arch Allergy Immunol* 2012;157:168–75.
- El Taieb MA, Fayed HM, Aly SS, *et al*. Assessment of serum 25-hydroxyvitamin D levels in children with atopic dermatitis: correlation with SCORAD index. *Dermatitis* 2013;24:296–301.
- Berents TL, Lødrup Carlsen KC, Mowinckel P, *et al*. Vitamin D levels and atopic eczema in infancy and early childhood in Norway: a cohort study. *Br J Dermatol* 2016;175:95–101.
- Barman M, Jonsson K, Hesselmar B, *et al*. No association between allergy and current 25-hydroxy vitamin D in serum or vitamin D intake. *Acta Paediatr* 2015;104:405–13.
- Chiu YE, Havens PL, Siegel DH, *et al*. Serum 25-hydroxyvitamin D concentration does not correlate with atopic dermatitis severity. *J Am Acad Dermatol* 2013;69:40–6.
- Yang HK, Choi J, Kim WK, *et al*. The association between hypovitaminosis D and pediatric allergic diseases: a Korean nationwide population-based study. *Allergy Asthma Proc* 2016;37:64–9.
- Lee SJ, Kang BH, Choi BS. Vitamin D serum levels in children with allergic and vasomotor rhinitis. *Korean J Pediatr* 2015;58:325–9.
- Cheon BR, Shin JE, Kim YJ, *et al*. Relationship between serum 25-hydroxyvitamin D and interleukin-31 levels, and the severity of atopic dermatitis in children. *Korean J Pediatr* 2015;58:96–101.
- Nam GE, Kim DH, Cho KH, *et al*. 25-Hydroxyvitamin D insufficiency is associated with cardiometabolic risk in Korean adolescents: the 2008–2009 Korea National Health and Nutrition Examination survey (KNHANES). *Public Health Nutr* 2014;17:186–94.
- Holick MF, Binkley NC, Bischoff-Ferrari HA, *et al*. Evaluation, treatment, and prevention of vitamin D deficiency: an Endocrine Society Clinical Practice guideline. *J Clin Endocrinol Metab* 2011;96:1911–30.
- McKinney K, Breitkopf CR, Berenson AB. Association of race, body fat and season with vitamin D status among young women: a cross-sectional study. *Clin Endocrinol* 2008;69:535–41.
- Choi HS, Oh HJ, Choi H, *et al*. Vitamin D insufficiency in Korea—a greater threat to younger generation: the Korea National Health and



- Nutrition Examination Survey (KNHANES) 2008. *J Clin Endocrinol Metab* 2011;96:643–51.
20. Kassi EN, Stavropoulos S, Kokkoris P, et al. Smoking is a significant determinant of low serum vitamin D in young and middle-aged healthy males. *Hormones* 2015;14:245–50.
  21. Manavi KR, Alston-Mills BP, Thompson MP, et al. Effect of serum cotinine on vitamin D serum concentrations among american females with different ethnic backgrounds. *Anticancer Res* 2015;35:1211–8.
  22. Wagner CL, Greer FR; American Academy of Pediatrics Section on Breastfeeding/American Academy of Pediatrics Committee on Nutrition. Prevention of rickets and vitamin D deficiency in infants, children, and adolescents. *Pediatrics* 2008;122:1142–52.
  23. Ross AC, Manson JE, Abrams SA, et al. The 2011 report on dietary reference intakes for calcium and vitamin D from the institute of medicine: what clinicians needs to know. *J Clin Endocrinol Metab* 2011;96:53–8.
  24. Ryoo E. Adolescent nutrition: what do pediatricians do? *Korean J Pediatr* 2011;54:287–91.
  25. Chung IH, Kim HJ, Chung S, et al. Vitamin D deficiency in Korean children: prevalence, risk factors, and the relationship with parathyroid hormone levels. *Ann Pediatr Endocrinol Metab* 2014;19:86–90.
  26. Carpenter TO, Herreros F, Zhang JH, et al. Demographic, dietary, and biochemical determinants of vitamin D status in inner-city children. *Am J Clin Nutr* 2012;95:137–46.
  27. Tolppanen AM, Fraser A, Fraser WD, et al. Risk factors for variation in 25-hydroxyvitamin D3 and D: concentrations and vitamin D deficiency in children. *J Clin Endocrinol Metab* 2012;97:1202–10.
  28. Yao TC, Tu YL, Chang SW, et al. Suboptimal vitamin D status in a population-based study of Asian children: prevalence and relation to allergic diseases and atopy. *PLoS One* 2014;9:e99105.
  29. Wortsman J, Matsuoka LY, Chen TC, et al. Decreased bioavailability of vitamin D in obesity. *Am J Clin Nutr* 2000;72:690–3.
  30. Cutillas-Marco E, Fuertes-Prosper A, Grant WB, et al. Vitamin D deficiency in South Europe: effect of smoking and aging. *Photodermatol Photoimmunol Photomed* 2012;28:159–61.
  31. Jeong BY, Lim MK, Yun EH, et al. Tolerance for and potential indicators of second-hand smoke exposure among nonsmokers: a comparison of self-reported and cotinine verified second-hand smoke exposure based on nationally representative data. *Prev Med* 2014;67:280–7.
  32. Jung S, Lee IS, Kim SB, et al. Urine cotinine for assessing tobacco smoke exposure in Korea: analysis of the Korea National Health and Nutrition Examination survey (KNHANES). *Tuberc Respir Dis* 2014;73:210–8.
  33. Florescu A, Ferrence R, Einarson T, et al. Methods for quantification of exposure to cigarette smoking and environmental tobacco smoke: focus on developmental toxicology. *Ther Drug Monit* 2009;31:14–30.
  34. Kim Y, Choi YJ, Oh SW, Sw O, et al. Discrepancy between self-reported and urine-cotinine verified smoking status among Korean male adults: analysis of health check-up data from a single private hospital. *Korean J Fam Med* 2016;37:171–6.
  35. Park MB, Nam EW, Lee SK, et al. The correlation of different cotinine levels with questionnaire results: a comparative study for different measurement methods of the adolescent smoking rate in Korea. *Asia Pac J Public Health* 2015;27:542–50.
  36. Haufroid V, Lison D. Urinary cotinine as a tobacco-smoke exposure index: a minireview. *Int Arch Occup Environ Health* 1998;71:162–8.
  37. Paul G, Brehm JM, Alcorn JF, et al. Vitamin D and asthma. *Am J Respir Crit Care Med* 2012;185:124–32.
  38. Bozzetto S, Carraro S, Giordano G, et al. Asthma, allergy and respiratory infections: the vitamin D hypothesis. *Allergy* 2012;67:10–17.
  39. Bath-Hextall FJ, Jenkinson C, Humphreys R, et al. Dietary supplements for established atopic eczema. *Cochrane Database Syst Rev* 2012;15:CD005205.
  40. Hollams EM, Hart PH, Holt BJ, et al. Vitamin D and atopy and asthma phenotypes in children: a longitudinal cohort study. *Eur Respir J* 2011;38:1320–7.
  41. van der Mei IA, Ponsonby AL, Dwyer T, et al. Vitamin D levels in people with multiple sclerosis and community controls in Tasmania, Australia. *J Neurol* 2007;254:581–90.