Correction: Factors predicting self-reported medication low adherence in a large sample of adults in the US general population: a cross-sectional study

Feehan M, Morrison MA, Tak C, *et al.* Factors predicting self-reported medication low adherence in a large sample of adults in the US general population: a cross-sectional study. *BMJ Open* 2017;7:e014435. doi: 10.1136/bmjopen-2016-014435

The description of the cutoff of the Morisky scale was listed as '6 or 7' but have been '6 to <8'. Therefore, the sentences:

'MMAS-8 scores can range from 0 to 8 with low adherence defined as a score <6, medium adherence as scores of 6 or 7 and high adherence with a score of 8.'

should have read

'MMAS-8 scores can range from 0 to 8 with low adherence defined as a score <6, medium adherence as scores of 6 to <8 and high adherence with a score of 8.'

and

'Using the standard cutoffs for the Morisky scale, 3862 (42%) respondents had "low" self-reported adherence (<6 on the scale), 2706 (29.4%) had "medium" adherence (6 or 7 on the scale) and 2635 (28.6) had 'high' adherence (score of 8).'

should have read:

'Using the standard cutoffs for the Morisky scale, 3862 (42%) respondents had "low" self-reported adherence (<6 on the scale), 2706 (29.4%) had "medium" adherence (6 to <8 on the scale) and 2635 (28.6) had "high" adherence (score of 8).'

Open Access This is an Open Access article distributed in accordance with the Creative Commons Attribution Non Commercial (CC BY-NC 4.0) license, which permits others to distribute, remix, adapt, build upon this work non-commercially, and license their derivative works on different terms, provided the original work is properly cited and the use is non-commercial. See: <http://creativecommons.org/licenses/by-nc/4.0/>

© Article author(s) (or their employer(s) unless otherwise stated in the text of the article) 2017. All rights reserved. No commercial use is permitted unless otherwise expressly granted.

BMJ Open 2017;7:e014435corr1. doi:10.1136/bmjopen-2016-014435corr1

CrossMark