Correction: Adherence to guidelines for creatinine and potassium monitoring and discontinuation following renin-angiotensin system blockade: a UK general practice-based cohort study

Schmidt M, Mansfield KE, Bhaskaran K, *et al.* Adherence to guidelines for creatinine and potassium monitoring and discontinuation following renin-angiotensin system blockade: a UK general practice-based cohort study. *BMJ Open* 2017;**7**:e012818. doi: 10.1136/bmjopen-2016-012818

In the 3rd column of Table 5 the heading ‘Serum potassium increase $\geq 30\%^*$ ’ should read ‘Serum potassium increase >6 mmol/L’.

Open Access This is an Open Access article distributed in accordance with the Creative Commons Attribution Non Commercial (CC BY-NC 4.0) license, which permits others to distribute, remix, adapt, build upon this work non-commercially, and license their derivative works on different terms, provided the original work is properly cited and the use is non-commercial. See: <http://creativecommons.org/licenses/by-nc/4.0/>

© Article author(s) (or their employer(s) unless otherwise stated in the text of the article) 2017. All rights reserved. No commercial use is permitted unless otherwise expressly granted.

BMJ Open 2017;**7**:e012818corr1. doi:10.1136/bmjopen-2016-012818corr1

CrossMark