

Supplementary tables

Future trends in the prevalence of severe activity limitations among older adults in Europe: a cross-national population study using EU-SILC

Fixed effects: estimates				
	Estimate	Std. Error	t value	Pr(> t)
Intercept	1.90780	0.0802	23.80	***
Age-group 55-59	-0.1166	0.0203	-5.76	***
Age-group 60-64	-0.2224	0.0203	-10.98	***
Age-group 65-69	-0.3502	0.0203	-17.29	***
Age-group 70-74	-0.5200	0.0203	-25.67	***
Age-group 75-79	-0.7183	0.0203	-35.46	***
Age-group 80-84	-0.9391	0.0203	-46.36	***
Age-group 85+	-1.1419	0.0203	-56.37	***
Country effects: variances				
Intercept	0.1618			
Residuals	0.0370			

Table S1: Random intercept model predicting the ratio of limitation free life expectancy and life expectancy for men in European countries between 2008 and 2014. Age group reference: 50-54; * $p \leq 0.05$; ** $p \leq 0.01$; *** $p \leq 0.001$

Fixed effects: estimates				
	Estimate	Std. Error	t value	Pr(> t)
Intercept	1.6565	0.0838	19.76	***
Age-group 55-59	-0.1120	0.0210	-5.34	***
Age-group 60-64	-0.2286	0.0210	-10.90	***
Age-group 65-69	-0.3761	0.0210	-17.93	***
Age-group 70-74	-0.5591	0.0210	-26.66	***
Age-group 75-79	-0.7610	0.0210	-36.29	***
Age-group 80-89	-0.9724	0.0210	-46.37	***
Age-group 85+	-1.1621	0.0210	-55.41	***
Country effects: variances				
Intercept	0.1770			
Residuals	0.0400			

Table S2: Random intercept model predicting the ratio of limitation free life expectancy and life expectancy for women in European countries between 2008 and 2014. Age group reference: 50-54; * $p \leq 0.05$; ** $p \leq 0.01$; *** $p \leq 0.001$