

# BMJ Open Scoping review of systematic reviews of complementary medicine for musculoskeletal and mental health conditions

Ava Lorenc,<sup>1</sup> Gene Feder,<sup>1</sup> Hugh MacPherson,<sup>2</sup> Paul Little,<sup>3</sup> Stewart W Mercer,<sup>4</sup> Deborah Sharp<sup>1</sup>

**To cite:** Lorenc A, Feder G, MacPherson H, *et al.* Scoping review of systematic reviews of complementary medicine for musculoskeletal and mental health conditions. *BMJ Open* 2018;**8**:e020222. doi:10.1136/bmjopen-2017-020222

► Prepublication history and additional material for this paper are available online. To view please visit the journal (<http://dx.doi.org/10.1136/bmjopen-2017-020222>).

Received 30 October 2017  
Revised 14 June 2018  
Accepted 1 August 2018



© Author(s) (or their employer(s)) 2018. Re-use permitted under CC BY-NC. No commercial re-use. See rights and permissions. Published by BMJ.

<sup>1</sup>Population Health Sciences, Bristol Medical School, Bristol, UK

<sup>2</sup>Department of Health Sciences, University of York, York, UK

<sup>3</sup>Primary Care and Population Science Unit, University of Southampton, Southampton, UK

<sup>4</sup>General Practice and Primary Care, Institute of Health and Wellbeing, University of Glasgow, Glasgow, UK

## Correspondence to

Dr Ava Lorenc;  
ava.lorenc@bristol.ac.uk

## ABSTRACT

**Objective** To identify potentially effective complementary approaches for musculoskeletal (MSK)–mental health (MH) comorbidity, by synthesising evidence on effectiveness, cost-effectiveness and safety from systematic reviews (SRs).

**Design** Scoping review of SRs.

**Methods** We searched literature databases, registries and reference lists, and contacted key authors and professional organisations to identify SRs of randomised controlled trials for complementary medicine for MSK or MH. Inclusion criteria were: published after 2004, studying adults, in English and scoring >50% on Assessing the Methodological Quality of Systematic Reviews (AMSTAR); quality appraisal checklist). SRs were synthesised to identify research priorities, based on moderate/good quality evidence, sample size and indication of cost-effectiveness and safety.

**Results** We included 84 MSK SRs and 27 MH SRs. Only one focused on MSK–MH comorbidity. Meditative approaches and yoga may improve MH outcomes in MSK populations. Yoga and tai chi had moderate/good evidence for MSK and MH conditions. SRs reported moderate/good quality evidence (any comparator) in a moderate/large population for: low back pain (LBP) (yoga, acupuncture, spinal manipulation/mobilisation, osteopathy), osteoarthritis (OA) (acupuncture, tai chi), neck pain (acupuncture, manipulation/manual therapy), myofascial trigger point pain (acupuncture), depression (mindfulness-based stress reduction (MBSR), meditation, tai chi, relaxation), anxiety (meditation/MBSR, moving meditation, yoga), sleep disorders (meditative/mind–body movement) and stress/distress (mindfulness). The majority of these complementary approaches had some evidence of safety—only three had evidence of harm. There was some evidence of cost-effectiveness for spinal manipulation/mobilisation and acupuncture for LBP, and manual therapy/manipulation for neck pain, but few SRs reviewed cost-effectiveness and many found no data.

**Conclusions** Only one SR studied MSK–MH comorbidity. Research priorities for complementary medicine for both MSK and MH (LBP, OA, depression, anxiety and sleep problems) are yoga, mindfulness and tai chi. Despite the large number of SRs and the prevalence of comorbidity, more high-quality, large randomised controlled trials in comorbid populations are needed.

## Strengths and limitations of this study

- A comprehensive scoping review covering a wide range of individual musculoskeletal and mental health conditions, in the notable dearth of systematic reviews (SRs)/randomised controlled trials in comorbid populations.
- This review used systematic searching and screening techniques.
- Results are limited by our pragmatic choice to only include high-quality SRs (high scoring on AMSTAR checklist (Assessing the Methodological Quality of Systematic Reviews)) and definition of complementary medicine as including a practitioner in its delivery.
- Date and language limitations may have excluded some topics.

## INTRODUCTION

Musculoskeletal (MSK) and mental health (MH) disorders are two conditions resulting in some of the highest burden in terms of disability, accounting for 30.5% and 21.1% of years lived with disability, respectively.<sup>1</sup> Both affect one in four UK adults at some point in their life.<sup>2–4</sup> MSK disorders lead to very high healthcare expenditure and loss of work,<sup>5</sup> in the UK accounting for 7.5 million working days lost annually<sup>6</sup> and 60% of occupational sick leave.<sup>2</sup> In England, it has been estimated that MH costs £105 billion, and treatment costs are expected to rise substantially.<sup>7</sup>

Low back pain (LBP) is one of the most common health conditions,<sup>8–10</sup> with a mean global general population prevalence (regardless of prevalence period) of 31%<sup>10</sup> and a UK prevalent population of 17.3 million, with 3.1 million adults suffering during an entire year,<sup>11</sup> costing the National Health Service (NHS) over £500 million/year.<sup>12</sup> In the UK, 8.75 million people sought primary care treatment for osteoarthritis (OA) (over a 7-year

consultation period)<sup>13</sup> which is estimated to have resulted in £14.8 billion indirect costs.<sup>14</sup>

The most common MH conditions are depression and anxiety.<sup>3,4</sup> In England, 19% of the general adult population reported ever being diagnosed with depression and 6% ever being diagnosed with generalised anxiety disorder.<sup>15</sup> Insomnia is another common MH problem, affecting 6.4% of the UK general population.<sup>16</sup>

There is an increasing recognition of the importance of multimorbidity, that is, the coexistence of two or more physical and/or mental long-term conditions in an individual, which is becoming the norm in the UK.<sup>17-19</sup> In a cross-sectional study in Scotland, 8.3% of primary care patients (36% of those with multimorbidity) had both a physical and a MH disorder<sup>20</sup> and around 50% of patients with depression had pain symptoms.<sup>21</sup> Long-term MSK disorders are closely associated with multimorbidity.<sup>22</sup>

Complementary medicine is 'a diverse group of health-related therapies and disciplines which are not considered to be a part of mainstream medical care', including osteopathy, chiropractic, acupuncture, herbal medicine and homeopathy.<sup>23</sup> Estimates of 12-month prevalence of use of any complementary medicine are 0.3%–86% in Europe<sup>24</sup> and 26% in England.<sup>25</sup>

Many MSK and MH conditions, and comorbid MSK and MH, have limited conventional treatment options, 'effectiveness gaps', which are potential key areas for complementary medicine.<sup>26,27</sup> Both MH<sup>28-30</sup> and MSK<sup>24,31</sup> are common reasons for patients to use complementary medicine. We used scoping methodology to identify priority areas where complementary medicine may be useful.<sup>32</sup> The overall aim of scoping studies is 'to map rapidly the key concepts underpinning a research area and the main sources and types of evidence available' in a formal, systematic and transparent way (which is often lacking in research priority setting exercises).<sup>33</sup> Scoping study methodology was used to identify gaps in the research base and priorities for future research. Scoping work is an essential step in the development of a strategy for the integration of complementary medicine with conventional care.<sup>27</sup> Preliminary searches showed that evidence for comorbid populations was very limited, so we chose to include independent evidence on MSK and MH conditions. This review was part of a wider scoping study, which obtained a breadth of perspectives—the results of this review were combined with findings from consultation with conventional and complementary medicine practitioners, a public survey and case studies of NHS provision of complementary medicine, to inform the final choice of conditions and complementary medicine approaches for a randomised controlled trial (RCT).

## METHODS

The scoping study followed Arksey and O'Malley's<sup>32</sup> framework plus the refinements suggested by Levac *et al.*,<sup>34</sup> Daudt *et al.*<sup>35</sup> and Colquhoun *et al.*<sup>36</sup> Scoping reviews are broader and more exploratory than systematic reviews

(SRs). They review a large number of papers to map the evidence—focussing on breadth rather than detail; have few preconceived ideas regarding focus (particularly types of MSK/MH in this context); do not necessarily assess individual study quality; and use a narrative approach to analysis, rather than synthesising or aggregating quantitative data.<sup>32</sup> The reader is advised to refer to the original SRs we have included for more detailed content.

## Aims

This study aimed to identify which practitioner-based complementary approaches have evidence for both MSK disorders and MH conditions in order to identify research priorities in terms of treatment choice and specific patient groups/conditions for a future pragmatic trial of comorbid MSK and MH in UK primary care. The specific aims were to:

- ▶ Collate the evidence on effectiveness, cost-effectiveness and safety from recent SRs of trials of complementary medicine for MSK and MH conditions.
- ▶ Identify areas where there is high-quality evidence of effectiveness but sufficient uncertainty to justify a trial.
- ▶ Identify areas where there is also some evidence of cost-effectiveness and safety.

## Searches

As this was a scoping review, we used a wide range of methods to identify SRs (published, unpublished, in progress). Literature databases were searched (see below for details of searches): AMED, Medline, EMBASE, CINAHL, PsycINFO; Index to theses; Cochrane library; Epistemonikos; SIGLE. Three trial registries were searched (UKCRN, Health service research projects in progress, Prospero). Reference lists of 'overviews of SRs' and reports (from professional/patient organisations) were searched. We emailed key authors in the field. We also asked the project team, steering group and public and patient involvement (PPI) group, and used websites/social media (university websites, Research Gate and Twitter) to access the wider research community. We asked patient and professional organisations to ask their members for any information (by email, newsletters or social media). These included: disease-specific organisations, for example, Arthritis Research UK, Back Care and MIND; complementary medicine organisations, for example, Research Council for Complementary Medicine and Complementary and Natural Healthcare Council; primary care organisations, for example, UK Royal College of General Practitioners. We presented at one conference and distributed a flyer at another. We contacted the authors of eligible conference abstracts and protocols to ascertain if the SR was available, that is, submitted or in press.

For complementary medicine, we used MeSH terms or subject headings, plus additional terms for complementary medicine which were not indexed. To ensure relevance to UK primary care, we only included complementary approaches which involve practitioner-led

treatment. We therefore excluded herbal medicine, as the majority of studies do not involve a practitioner. We included manipulation, manual therapy and mobilisation as techniques commonly practised by some complementary medicine practitioners (as well as conventional practitioners). For MSK conditions, we used ‘musculoskeletal diseases/disorders’ as index terms, adding in any which were missing. For MH, we searched for common, minor MH disorders and symptoms, excluding long-term/severe mental illnesses. Thus, words in titles/abstracts were searched for (rather than subject headings), with reviews needing to include patients with MH symptoms/diagnoses rather than just measuring MH outcomes. See online supplementary appendix 1 for a full list of included and excluded non MSK or MH conditions.

Database searches were conducted in June 2015 and updated in February 2016, in collaboration with a librarian. We used filters for SRs. We excluded reviews published pre-2005, in order to make the final number of papers manageable with minimal impact on our results (given that the majority of SRs were published in 2010 and after (122/158 SRs)). We only included English language reviews. Databases were searched for two topics: complementary medicine+musculoskeletal disorders and complementary medicine+mental health. See online supplementary appendix 1 for details and supplementary appendix 2 for an example search.

We used the inclusion/exclusion criteria (see online supplementary appendix 1) to screen the results from each search. This was a multistage process involving the whole project team. AL screened the titles of all search results, and DS checked the results, with any disagreements discussed and inclusion criteria amended accordingly. AL then screened the abstracts, which were also independently screened by another author. Results were compared, any disagreements discussed and the criteria again amended accordingly. Any remaining disagreements were discussed and resolved with a third author. AL then screened the full text of the agreed papers.

### Data extraction

AL extracted the following data from each SR: author, year, location, complementary medicine/s, health condition/s, for MSK reviews whether MH was included and for MH reviews whether findings applied to MSK populations, methodological features (sources searched, inclusion/exclusion criteria, quality assessment), results and conclusions (effectiveness, safety, cost-effectiveness). Where possible, these data were copied and pasted directly, to avoid any misinterpretation.

### Quality appraisal of SRs

Although not essential,<sup>32</sup> quality assessment is recommended in scoping reviews.<sup>37</sup> We used a limited quality appraisal of SRs to help inform prioritisation, using AMSTAR (Assessing the Methodological Quality of Systematic Reviews), a validated checklist.<sup>38</sup> Due to time

constraints and the number of SRs to be appraised, a simple procedure was followed, using the ‘search’ function in the text of SRs to find particular terms. The emphasis was on obtaining relative scores, allowing ranking of the included reviews, rather than an absolute score of quality. For each of the 11 items on the AMSTAR checklist, a score of 0 was given if they did not meet the criteria or if information was unavailable, a score of 1 if the criteria was met. The scores were summed to give a score for each SR out of a total of 11.

Any reviews which scored 5 or less (half of the possible total) on AMSTAR were not considered in the evidence synthesis.

### Evidence synthesis

In line with Arksey and O’Malley’s<sup>32</sup> framework for conducting scoping reviews, we used a ‘descriptive-analytical’ method, based on the narrative tradition, applying a common analytical framework to the reviews. Evidence synthesis was performed by AL and all authors met regularly to discuss the process and emerging results.

Reviews were placed into a matrix of health condition/symptom versus complementary medicine, for example, LBP and acupuncture (see [tables 1 and 2](#)).

Where there was more than one SR in an area, one (occasionally more than one—where they studied different interventions, eg, types of manipulation) was prioritised, based on recency, breadth, quality (AMSTAR score) and level of analysis (eg, meta-analysis).

There were three steps to the evidence synthesis: step 1: extraction and narrative synthesis of data from prioritised SRs; step 2: ranking of areas (condition vs complementary medicine) using an Excel framework (see below for ranking criteria); and step 3: computation of sample size.

#### Step 1 data extraction

For each area (condition vs complementary medicine) in the matrix, data were synthesised for:

- ▶ Effectiveness: the conclusions on effectiveness were extracted from the prioritised SR/s, including information on comparator and outcomes.
- ▶ Quality of studies: in line with guidance on scoping reviews,<sup>32</sup> we did not assess the quality of the individual RCTs in each SR, but synthesised the SR authors’ conclusions about quality, from the prioritised review/s. We had three categories of quality: poor, moderate or good. Moderate included SRs reporting mixed results.
- ▶ Safety: any information on safety was synthesised across all the included SRs.
- ▶ Cost-effectiveness: any information on cost-effectiveness was synthesised across all the included SRs.

#### Step 2 ranking

Using the information extracted in step 1, we ranked the areas in the matrix (condition vs complementary

**Table 1** Matrix of musculoskeletal reviews (numbers=number of reviews)

	Low back pain		Fibromyalgia		Neck pain/ disorders		Osteoarthritis		Rheumatoid arthritis		MSK pain/ conditions		Myofascial pain		Shoulder pain/ disorder		Other		Total	
	Total number	High quality	Total number	High quality	Total number	High quality	Total number	High quality	Total number	High quality	Total number	High quality	Total number	High quality	Total number	High quality	Total number	High quality		
Acupuncture related																				
Acupressure	2	2			2	1					1	1					1	0	6	
Acupuncture	12	6	9	6	5	2	10	6	4	2	4	2	6	3	2	1	6	5	58	
Moxibustion	1	0	1	0			1	0	2	1							1	0	6	
Manipulation																				
Chiropractic	2	1	3	1			1	0					1	0					7	
Manual therapy/spinal manipulation/mobilisation	12	9	8	4	14	9	2	1			1	0	1	0	5	3	3	2	46	
Osteopathy	3	2	2	1							1	1							6	
Other																				
Integrative therapies/multiple complementary medicine	2	2	2	2															4	
Tai chi	1	1	1	0	1	0	7	5	3	1	1	1							14	
Qigong					1	0	1	0	1	0									3	
Pilates	10	5																	10	
Complementary medicine exercise			2	2															2	
Yoga	8	4	1	1	1	0	3	0	1	0	1	1							15	
Mindfulness/meditation	2	0	5	3					2	0	4	2							13	
Relaxation			3	1			1	0	1	0	1	0							6	
Hypnotherapy	1	0	1	1			1	0	1	0									4	
Spa therapy/balneotherapy	2	1	2	1					1	1									5	
Traditional Chinese Medicine bodywork	2	1			1	0					1	0							4	
Massage/therapeutic touch	4	2			4	2	1	0	1	0	1	0	1	0	1	0	1	1	13	
Homeopathy			2	0															2	
Total*	64	36	42	23	29	14	28	12	17	5	16	8	8	3	8	4	12	8	158 (84)	

\*Note that reviews can occur in more than one row/column.

**Table 2** Matrix of mental health reviews (numbers=number of reviews)

	Depression		Anxiety		Stress		Insomnia/sleep		Distress (psychological)		Total*
	Total number	High quality	Total number	High quality	Total number	High quality	Total number	High quality	Total number	High quality	
<b>Acupuncture-related</b>											
Acupuncture	9	3	2	1			2	1			6
Auricular acupuncture							3	1			4
<b>Multiple</b>											
Complementary medicine (multiple) (did not separate complementary medicine in analysis)	1	0	1	0			2	1			4
<b>Meditation/mindfulness</b>											
Mindfulness	3	2	1	1	1	1			1	1	5
Meditation			3	3	1	1	1	0			9
<b>Exercise</b>											
Meditative movement							3	3			6
Qigong	4	1	2	1	1	1					5
Tai chi	4	2	1	0			3	1			5
Yoga	4	2	3	3	1	0	1	0			8
<b>Other</b>											
Homeopathy							1	1			2
Hypnosis	2	0					2	2			4
Music therapy	3	2	1	1							2
Relaxation	3	1	1	1			1	1			4
Reflexology							1	0			1
Reiki	1	1	1	1							2
Spinal manipulation											0
Massage/therapeutic touch	2	1	2	2							4
Morita therapy			1	1							2
<b>Total*</b>	<b>36</b>	<b>15</b>	<b>19</b>	<b>15</b>	<b>4</b>	<b>3</b>	<b>20</b>	<b>11</b>	<b>1</b>	<b>1</b>	<b>48 (27)</b>

\*Note that reviews can occur in more than one row/column

medicine), based on consideration of the following criteria:

1. Effectiveness: areas were ranked highly where there was enough moderate/good quality evidence to establish proof of concept, but still key gaps. Data on effectiveness came from the conclusions of the prioritised SRs regarding effectiveness and quality.
2. Safety: areas where there was evidence of harm were ranked lower.
3. Cost: areas where there was any indication that the intervention may have potential for cost-effectiveness were ranked highly.
4. Comorbidity: any interventions that showed an indication (of any quality) for comorbid MH and MSK patients were ranked highly.

### Step 3 sample size

Once a ranked list of areas had been compiled, information on total sample size for the prioritised SR/s (ie, total number of participants in the trials included in the SR/s) was added to allow further discrimination, classified as: small (<500 participants), medium (501 to 3000) or large (>3000).

### Patient and public involvement

The wider scoping study included a PPI group of seven people with experience of MSK/MH issues, who advised the research team on a variety of issues as the study took shape and progressed. They met four times over the study period and provided input into the literature review questions, search strategy and selection criteria.

The PPI group also provided ideas on ways to disseminate the findings to the general public and to patient groups.

## RESULTS

### Search results

Figure 1 shows the Preferred Reporting Items for Systematic Reviews and Meta-Analyses flowchart of the searches.<sup>39</sup> Searches resulted in 8393 references—4304 for MH and 4089 for MSK, plus three for MH identified elsewhere (email update sent by journal publisher). After removing duplicates, Cochrane reviews which were withdrawn, Cochrane reviews which had been updated or those not in English, 5230 records remained for title screening. This resulted in 486 MH and 739 MSK records for the abstract screening phase. We then excluded 49 (13 MH and 36 MSK) records, that is, editorials, commentaries, abstracts etc. After abstract screening, there remained 444 studies (180 MH and 264 MSK). At this stage, a further 12 records were added—three from grey literature (Prospero/Sigle/HSRProj), five from conference abstracts and four from the reference lists of overviews of SRs and other key reports. This produced 456 records to be screened at the full-text stage, after which 206 records remained: 48 MH, 158 MSK.

During screening at the title stage (AL and DS), there were 49 disagreements (out of 4411 records; 1%). During abstract screening (AL and each of the six authors), there were a total of 296 initial disagreements (out of 1165 records; 25%). Most were resolved by discussion without requiring a third reviewer.

Of the 158 MSK SRs, 84 were included as high quality (scored six or more on AMSTAR): LBP,<sup>40–69</sup> fibromyalgia,<sup>70–81</sup> neck pain,<sup>40 82–90</sup> OA,<sup>91–101</sup> rheumatoid arthritis,<sup>97 102–105</sup> general MSK pain/conditions,<sup>106–110</sup> shoulder pain/disorders,<sup>111–115</sup> myofascial trigger point pain,<sup>106 116 117</sup> and other musculoskeletal (MSK) conditions.<sup>106 118–123</sup>

The 158 reviews covered 17 different and 15 different disorders. The overall matrix is shown below in table 1. LBP was by far the most common topic, addressed by 64/158 (40%) of reviews. Fibromyalgia was the next most common, followed by OA then neck pain/disorders. Acupuncture was the most common complementary medicine studied (58/158 SRs; 37%), followed by manual therapy (including manipulation and mobilisation). The mean AMSTAR score for methodological quality of MSK reviews was 6/11 (ranging from 0 to 11). Online supplementary appendix 3 shows which items on the AMSTAR checklist were reported by the reviews.

Of the 48 MHSRs, 27 were high quality (scored six or more on AMSTAR): depression,<sup>124–134</sup> anxiety,<sup>124 125 129 133 135–142</sup> sleep,<sup>143–150</sup> stress,<sup>124 125 140</sup> and other.<sup>124</sup> The 48 reviews covered 18 different and five different categories of MH symptom or disorder. The overall matrix is shown in table 2. Many of the reviews included more than one MH condition. Of the 48 reviews, 36 (75%) included

depression. Meditation, yoga, acupuncture and meditative movement were the most common complementary medicine. Online supplementary appendix 3 shows which items on the AMSTAR checklist were reported by the reviews.

The following sections present a summary of the areas where we found moderate/good quality evidence of effectiveness (from the prioritised SRs). For those conditions with evidence for more than one intervention, we have presented the results in tables.

### Evidence for MSK disorders

The areas where there was moderate/good quality evidence were LBP, myofascial trigger point pain, OA, neck pain, fibromyalgia and lateral epicondylitis. It should be noted that the SR quality of evidence is based on the overall assessment of quality of the included RCTs by the SR authors, which is subject to some limitations.

#### Low back pain

Moderate/good quality evidence for LBP (see table 3) was found as follows:

- ▶ Yoga: four high-quality reviews,<sup>51 60–62</sup> one prioritised<sup>62</sup>
- ▶ Osteopathy: two high-quality reviews,<sup>64 65</sup> one prioritised<sup>65</sup>
- ▶ Acupuncture: six high-quality reviews,<sup>40 41 49 50 53 54</sup> two prioritised—as highest quality<sup>40</sup> and most recent<sup>50</sup>
- ▶ Spinal manipulation/mobilisation: nine high-quality reviews,<sup>40–48</sup> three prioritised (due to differences in types of back pain and interventions included)<sup>40 43 44</sup>
- ▶ Spa/balneotherapy: one high-quality review<sup>52</sup>
- ▶ Tai chi: one high-quality review<sup>54</sup>

#### Osteoarthritis

Moderate/good quality evidence for OA (see table 4) was found as follows:

- ▶ Acupuncture: six high-quality reviews,<sup>91–96</sup> two prioritised<sup>92</sup> as a Cochrane review and<sup>94</sup> as most recent.
- ▶ Tai chi: five high-quality reviews,<sup>97–101</sup> one prioritised/<sup>99</sup>

#### Neck pain


Moderate/good quality evidence for neck pain (see table 5) was found as follows:

- ▶ Manual therapy/manipulation: eight high-quality reviews,<sup>40 82–88</sup> three were prioritised due to differing inclusion criteria.<sup>82 87 88</sup>
- ▶ Acupuncture: two high-quality reviews,<sup>40 89</sup> one prioritised.<sup>89</sup>

There was moderate quality evidence in a medium-sized population for a range of other techniques.<sup>82 87 88</sup>

#### Fibromyalgia

Acupuncture for fibromyalgia was reviewed by—five high-quality reviews,<sup>70–73 81</sup> one prioritised.<sup>72</sup> The prioritised review found moderate quality evidence in a small population compared with placebo for stiffness (acupuncture) and for pain and sleep disorders (electroacupuncture). However, there was some evidence that acupuncture is less effective than usual care for



**Figure 1** Preferred Reporting Items for Systematic Reviews and Meta-Analyses flowchart. MH, mental health; MSK, musculoskeletal; SR, systematic review.

**Table 3** Evidence for low back pain

Comparator	Quality	Population size	Complementary medicine	Safety	Cost-effectiveness
Active intervention	Good	Medium	Osteopathy <sup>65</sup>	Some evidence of harm <sup>65*</sup>	NR
	Moderate	Large	Spinal manipulation/mobilisation (chronic only) <sup>40 43 44</sup>	Some evidence of safety <sup>44†</sup>	Mixed results—some evidence of CE <sup>47‡</sup>
		Medium	Yoga <sup>62</sup>	Some evidence of safety <sup>51 60 62§</sup>	No data
	Moderate	Small	Balneotherapy <sup>52</sup>	Some evidence of safety <sup>52</sup>	NR
Usual care	Moderate	Large	Acupuncture <sup>40 50</sup> (for pain, function, well-being, disability, range of movement, quality of life)	Some evidence of safety <sup>53 54¶</sup>	Some evidence of CE <sup>40**</sup>
		Small	Spa therapy <sup>52</sup>	Some evidence of safety <sup>52††</sup>	NR
			Tai chi <sup>54</sup>	NR	NR
Placebo	Moderate	Large	Acupuncture (for pain and quality of life; negative evidence for disability). <sup>40 50</sup>	Some evidence of safety <sup>53 54¶</sup>	Some evidence of CE <sup>40**</sup>

\*Two studies reported minor adverse events.

†One review identified serious adverse events, but this appeared unrelated to treatment. Minor adverse events included muscle soreness, stiffness and/or transient increase in pain.

‡Mixed results regarding cost-effectiveness,<sup>40</sup> with limited evidence that manual therapy is more cost-effective than usual care.<sup>47</sup>

§No serious adverse events associated with the yoga.

¶5% minor adverse events and no serious adverse events.

\*\*Short-term only.

††No adverse events from one study.

CE, cost effectiveness; NR, not reported (by systematic reviews).

MH outcomes in fibromyalgia.<sup>72</sup> There is some evidence of safety (no serious adverse events)<sup>70–73 81</sup> but none of cost-effectiveness.

#### Myofascial trigger point pain

Acupuncture for myofascial trigger point pain was reviewed by—three high-quality reviews.<sup>106 116 117</sup> The prioritised review found good quality evidence in a medium-sized population compared with placebo and other active interventions for pain in the short and medium term but not in the long term.<sup>117</sup> However, safety, cost-effectiveness or impact on MH outcomes were not reported.<sup>106 116 117</sup>

#### Lateral epicondylitis

One high-quality review<sup>120</sup> found moderate quality evidence for manual therapy (mobilisation with

movement) but the comparator and sample size were not reported so we are unable to include it in our synthesis.

#### Evidence for mental health

No reviews provided good quality evidence for complementary medicine for any MH condition nor any evidence of cost-effectiveness, mainly because the latter was not reported.

#### Depression

Moderate quality evidence for depression (see [table 6](#)) was found as follows:

- Mindfulness/meditation: two high-quality reviews,<sup>124 125</sup> both prioritised as included different types of meditation/mindfulness/mindfulness-based stress reduction (MBSR) or populations.

**Table 4** Evidence for osteoarthritis

Comparator	Quality	Population size	Complementary medicine	Safety	Cost-effectiveness
Placebo	Good	Large	Acupuncture <sup>92 94*</sup>	Some evidence of harm <sup>91 92†</sup>	NR
Usual care	Moderate	Small	Tai chi <sup>99‡</sup>	Some evidence of safety <sup>98–100§</sup>	NR

\*Compared to pulsed electromagnetic fields, muscle-strengthening or aerobic exercise, weight loss.

†Up to 45% frequency of minor adverse events.

‡Only in the short term.

§Systematic reviews (SRs) concluded tai chi is safe.

NR, not reported (by SRs).



**Table 5** Evidence for neck pain

Comparator	Quality	Population size	Complementary medicine	Safety	Cost-effectiveness
Active intervention	Good	Medium	Manual therapy <sup>87*</sup>	Some evidence of potential harm <sup>82 83†</sup>	Some evidence of CE <sup>87 88‡</sup>
	Moderate	Large	Manipulation <sup>82 §¶</sup>	Some evidence of potential harm <sup>82 83†</sup>	Some evidence of CE <sup>87 88‡</sup>
	Moderate	Small	Manual therapy <sup>88**</sup>	Some evidence of potential harm <sup>82 83†</sup>	Some evidence of CE <sup>87 88‡</sup>
Placebo	Moderate	Medium	Acupuncture <sup>89</sup>	Some evidence of safety <sup>89</sup>	No data <sup>40</sup>
Usual care	Moderate	Medium	Acupuncture <sup>89</sup>	Some evidence of safety <sup>89</sup>	No data <sup>40</sup>

\*Manual therapy plus exercise compared with exercise alone.

†Reduced cost of therapy.

‡Minor adverse events and rare but devastating adverse effects from manipulation.

§Compared to medication.

¶Also showed effectiveness for pain compared with any control.

\*\*Mobilisation, manipulation and soft tissue techniques.

CE, cost effectiveness.

- ▶ Tai chi: two high quality reviews,<sup>132 134</sup> both prioritised as Chi *et al*<sup>134</sup> only reviewed studies of older people (Liu *et al*<sup>132</sup> found only poor quality evidence).
- ▶ Relaxation: one high-quality review<sup>129</sup> but comparator and sample size were not reported so it is not included here.

### Anxiety

Moderate quality evidence for anxiety (see [table 7](#)) was found as follows:

- ▶ Meditation, including moving meditation (yoga, tai chi and qi gong): three high-quality reviews,<sup>125 137 138</sup> one prioritised.<sup>138</sup>
- ▶ MBSR: one high-quality review.<sup>124</sup>
- ▶ Yoga: three high-quality reviews,<sup>129 135 136</sup> one prioritised.<sup>136</sup>

### Sleep disorders

Three high-quality reviews considered meditative/mind–body movement for sleep.<sup>143–145</sup> The prioritised review found moderate quality evidence in older adults for meditative movement in a medium-sized population, compared with usual care and other active interventions.<sup>143</sup> There was some evidence of safety (no adverse events)<sup>143</sup> and effect on sleep quality in an MSK population.<sup>145</sup> No cost-effectiveness data reported.

### Stress and distress

There was one high-quality review of mindfulness for stress and distress.<sup>124</sup> For stress, they found moderate quality evidence in a medium-sized population compared with usual care and placebo, and some evidence of effectiveness on stress as an outcome in an MSK population.<sup>124</sup> There were no data on cost-effectiveness and no data on safety. For distress, they found moderate quality evidence in a medium-sized population compared with usual care, and some evidence of effectiveness on distress as an outcome in an MSK population.<sup>124</sup> There were no data on cost-effectiveness and safety was not reported.

There was only poor quality evidence for all other combinations of MH condition/complementary medicine/comparator.

### Evidence for effectiveness in comorbidity/MH outcomes in MSK

Only one SR provided data on comorbid MSK and MH conditions (depression and a variety of comorbid MSK conditions).<sup>132</sup> Three SRs provided data on MH outcomes in MSK populations (but did not specify an MH score/diagnosis as inclusion criterion). Tai chi for depression and comorbid conditions (including MSKs) may be less effective than for depression alone<sup>132</sup>; MBSR/meditation

**Table 6** Evidence for depression

Comparator	Quality	Population size	Complementary medicine	Safety	Cost-effectiveness
Active intervention	Moderate	Medium	Meditation <sup>125*</sup>	Some evidence of safety <sup>§</sup>	No data <sup>124</sup>
Usual care	Moderate	Medium	MBSR (mindfulness-based stress reduction) <sup>124*</sup>	Some evidence of safety <sup>§</sup>	No data <sup>124</sup>
	Moderate	Medium	Tai chi <sup>134 †‡</sup>	No data <sup>134</sup>	No data <sup>134</sup>

\*Also some evidence of effectiveness for both in musculoskeletal (MSK) populations.<sup>124 125</sup>

†Older adults only.

‡May be less effective in comorbid populations (including some MSK conditions).<sup>132</sup>

§Nine trials found no adverse events.

**Table 7** Evidence for anxiety

Comparator	Quality	Population size	Complementary medicine	Safety	Cost-effectiveness
Active intervention	Moderate	Medium	Meditation <sup>138*</sup> †	Some evidence of safety <sup>125 137 138</sup>	NR
	Moderate	Medium	Yoga <sup>136</sup> ‡	No data <sup>135</sup>	NR
Placebo	Moderate	Medium	Meditation <sup>138*</sup> †	Some evidence of safety <sup>125 137 138</sup>	NR
Usual care	Moderate	Medium	Meditation <sup>138*</sup> †	Some evidence of safety <sup>125 137 138</sup>	NR
	Moderate	Medium	MBSR <sup>124</sup> †	Some evidence of safety <sup>125 137 138</sup>	NR

\*Moving meditation (tai chi, qi gong or yoga) was more effective than static meditation.

†Mindfulness-based stress reduction (MBSR)/meditation was also effective for anxiety (as an outcome) in a musculoskeletal population.<sup>124 125</sup>

‡For women with breast cancer only.

NR, not reported (by systematic reviews).

was effective for MH outcomes (depression, anxiety, stress/distress) in an MSK population<sup>124 125</sup>; meditative/mind–body movement was effective for sleep outcomes in an MSK population.<sup>145</sup>

### Complementary medicine with evidence for both MSK and MH conditions

From the above synthesis, complementary medicine with moderate/good quality evidence for both MSK and MH conditions (from separate SRs) are yoga and tai chi. Yoga appears to be effective for both LBP and anxiety compared with other active interventions, although the anxiety finding was only from studies of women with breast cancer. Meditative movement, which includes yoga, is effective for sleep disorders, although only for older adults (there is also some evidence in chronic pain and fibromyalgia populations). Tai chi appears to be effective compared with usual care for LBP, knee OA and depression, in older people (although it appears less effective in comorbid depressed patients). Meditative movement, which includes tai chi, is effective for sleep disorders, although only for older adults (there is some evidence in chronic pain and fibromyalgia populations as well).

### Update

Since completing this review, in July 2017, we searched for recent SRs on the highest ranking topics (results are not included in tables 3–7). These either provide additional evidence for MH outcomes in an MSK population, or evidence supporting our original conclusions.

- Yoga: a new Cochrane review for LBP has been published which suggests there is moderate-quality evidence for yoga compared with other active interventions for pain, and for function at 6 months only, although they also found moderate-quality evidence of harm for yoga (primarily exacerbation of back pain).<sup>151</sup> They also found very low-quality MH evidence for yoga improving depression (as an outcome not a diagnosis) in LBP (comparator not stated).<sup>151</sup>
- MBSR/mindfulness/meditation: a new SR provides evidence for mindfulness and MH, including high-quality evidence (sample size not reported) for mindfulness helping with depression (compared with

all controls—usual care, and active interventions (education, stress management); (as an outcome not a diagnosis) in LBP (and fibromyalgia).<sup>152</sup> Another SR of mindfulness for LBP was recently published but cannot be included as there were no conclusions regarding study quality.<sup>153</sup>

- Acupuncture (compared with other active interventions): Two new SRs confirm the effectiveness of acupuncture for OA of the knee, with moderate-quality evidence in a large sample compared with active interventions and placebo.<sup>154 155</sup>

### DISCUSSION

This scoping review has highlighted the large and increasing number of SRs of complementary medicine for MSK and MH disorders, covering 29 different complementary medicine approaches, but identified no high-quality evidence for complementary medicine for MSK–MH comorbid populations. Three reviews showed that MBSR/meditation/meditative movement may improve MH outcomes in MSK populations. Acupuncture, yoga, tai chi and mindfulness/meditation have evidence for both MSK and MH conditions.

We have identified moderate/good-quality evidence (as reported by SR authors) for effectiveness (any comparator) in a moderate/large population for: LBP (yoga, acupuncture, spinal manipulation/mobilisation, osteopathy), OA (acupuncture, tai chi), neck pain (acupuncture, manipulation/manual therapy), myofascial trigger point pain (acupuncture), depression (MBSR, meditation, tai chi, relaxation), anxiety (meditation/MBSR, moving meditation, yoga), sleep disorders (meditative/mind–body movement) and stress/distress (mindfulness). The majority of these complementary approaches had some evidence of safety—only three had evidence of harm—osteopathy for LBP, acupuncture for OA and manipulation/manual therapy for neck pain. There was some evidence of cost-effectiveness for spinal manipulation/mobilisation and acupuncture for LBP, and manual therapy/manipulation for neck pain, but few SRs reviewed cost-effectiveness and many found no data.

Regarding comorbidity, from our original searches, the only evidence for MH outcomes in an MSK population (not necessarily with MH conditions) was for MBSR/meditation/meditative movement/mindfulness. Tai chi and yoga were effective for MSK and MH conditions in separate SRs (not comorbid populations). Our 2017 update found new (but poor quality) evidence for yoga in MSK–MH comorbidity and more evidence for mindfulness in MSK–MH comorbidity.

Some of these studies showed effectiveness compared with an active comparator (spinal manipulation/mobilisation and yoga for LBP; meditation for depression and anxiety), providing some evidence that these interventions have effects over and above non-specific attention.

The quality of the SRs was very variable. We used a quality tool, AMSTAR, which itself has some limitations, as it includes both methodological and reporting items which are not all necessarily equal in importance, and some of which can be difficult to interpret. Future reviews may consider prioritising certain items on AMSTAR, using the new version of AMSTAR,<sup>156</sup> or using an alternative critical appraisal tool such as the Oxford Centre for Evidence-based Medicine checklist (<https://www.cebm.net/2014/06/critical-appraisal/>). An assessment of risk of bias, using, for example, Risk Of Bias In Systematic reviews (ROBIS), may be useful in addition to these quality assessments.

The quality of included trials also varied greatly; it is notable that none of the MH SRs concluded that trial quality was ‘good’ overall. In addition, assessment of individual RCT study quality is likely to have varied between SR authors, a common issue with complementary medicine studies which rarely fit conventional quality criteria (particularly blinding). Very few of the MSK SRs and none of the MH SRs mentioned cost-effectiveness. Safety was rarely adequately reported in SRs or the studies they included. Future SRs in the areas of complementary medicine and MSK/MH need to ensure they include a priori design/protocol registration, a list of excluded studies and conflict of interest statements, and use duplicate study selection and data extraction, assess publication bias and search for grey literature.

Both a strength and limitation of this scoping review was its comprehensive scope which included a wide range of complementary medicine approaches and types of MSK and MH condition, but limited the depth of analysis and discussion.

Other key strengths were the systematic searching techniques, prioritising high-quality SRs using MeSH terms where possible, and search techniques to capture grey and unpublished literature. The whole project team was involved in independently screening the literature search results, although a limitation was that only one author conducted data extraction. We included a quality appraisal of the reviews, although time constraints meant this was an abbreviated assessment. One of the key limitations was the pragmatic decision to only include SRs, which may have led to the exclusion of some complementary

medicines, and means results are subject to the limitations of SR and trial methodology, study designs which are sometimes challenging for complementary medicine.<sup>157–159</sup> Another key limitation is our definition of complementary medicine as including a practitioner in its delivery (which relied on review authors reporting these details), which excluded over-the-counter products or self-care practices, herbal medicine in particular, but was necessary to limit the scope of the review to complementary medicine which may be appropriate for an integrated care model using referral from a general practitioner. We are aware that by excluding reviews published pre-2005, we may have excluded some topics. We were unable to include non-English publications.

Although complementary medicine is commonly used by patients in the UK,<sup>25</sup> it is not widely available via the NHS. Given the high burden of MSK and MH conditions to individuals and society in the UK, and the prevalence of comorbidity, complementary medicine may be worth considering for some patients, where there is evidence of effectiveness, cost-effectiveness and safety.

One of the main aims of this study was to inform the design of a future trial to assess the effectiveness and cost-effectiveness of an integrated medicine approach for patients with comorbid MSK and MH disorders. We will combine the results of this review with the views of conventional and complementary medicine practitioners, the public and NHS providers, to inform the final focus for an RCT. Our proposed trial would be pragmatic and focus on overall effectiveness, so interventions with evidence compared with any comparator are potential candidates.

SRs of complementary medicine consistently conclude that further high-quality trials are needed, with longer follow-up and larger sample sizes, assessing cost-effectiveness and use of appropriate, sensitive, validated outcome measures.

Based on our criteria of study quality, safety, cost-effectiveness and evidence for MSK MH comorbidity, this scoping review suggests that the most promising complementary medicine for future research in the area of MH–MSK comorbidity are: mindfulness (evidence for MH in MSK populations and evidence of safety), yoga (evidence for MSK and MH conditions) and tai chi (evidence for MSK and MH conditions, evidence of safety). The comorbid populations that would benefit most are less clear, as so few reviews focus on comorbidity, but may include LBP, OA, depression, anxiety and sleep.

**Acknowledgements** We are grateful to Catherine Borwick for her help with literature searching and Sandra Hollinghurst for her help with screening the papers. We are also very grateful to our PPI group.

**Contributors** AL conducted all literature searches, data extraction and quality appraisal, with the project team (GF, HM, PL, SWM and DS) all screening papers for inclusion. The team (GF, HM, PL, SWM and DS) all provided input into the inclusion/exclusion criteria. DS advised on the review process. GF, HM, PL, SWM and DS were all involved in the prioritisation of topics from the review results. AL drafted the paper and all authors reviewed and edited it.

**Funding** The report is based on independent research commissioned and funded by the NIHR Policy Research Programme (The Effectiveness And Cost Effectiveness Of Complementary And Alternative Medicine (Cam) For Multimorbid Patients With Mental Health And Musculoskeletal Problems In Primary Care In The UK: A Scoping Study).

**Disclaimer** The views expressed in the publication are those of the author(s) and not necessarily those of the NHS, the NIHR, the Department of Health, 'arms' length bodies or other government departments.

**Competing interests** None declared.

**Patient consent** Not required.

**Provenance and peer review** Not commissioned; externally peer reviewed.

**Data sharing statement** We are able to provide copies of searches and lists of excluded references on request.

**Open access** This is an open access article distributed in accordance with the Creative Commons Attribution Non Commercial (CC BY-NC 4.0) license, which permits others to distribute, remix, adapt, build upon this work non-commercially, and license their derivative works on different terms, provided the original work is properly cited, appropriate credit is given, any changes made indicated, and the use is non-commercial. See: <http://creativecommons.org/licenses/by-nc/4.0/>.

## REFERENCES

- Murray CJL, Richards MA, Newton JN, et al. UK health performance: findings of the Global Burden of Disease Study 2010. *Lancet* 2013;381:997–1020.
- Department of Health, 2006. The Musculoskeletal Services Framework Y1. Secondary The Musculoskeletal Services Framework Y1. [http://webarchive.nationalarchives.gov.uk/20130107105354/http://www.dh.gov.uk/prod\\_consum\\_dh/groups/dh\\_digitalassets/@dh/@en/documents/digitalasset/dh\\_4138412.pdf](http://webarchive.nationalarchives.gov.uk/20130107105354/http://www.dh.gov.uk/prod_consum_dh/groups/dh_digitalassets/@dh/@en/documents/digitalasset/dh_4138412.pdf)
- Centre HSCI, et al. Sally McManus HM, Brugha T, Bebbington P, Jenkins R, eds. *Adult psychiatric morbidity in England, 2007: results of a household survey*, 2009.
- Halliwel E, Main L, Richardson C. The fundamental facts. Secondary the fundamental facts 2007. [http://www.mentalhealth.org.uk/content/assets/PDF/publications/fundamental\\_facts\\_2007.pdf?view=Standard](http://www.mentalhealth.org.uk/content/assets/PDF/publications/fundamental_facts_2007.pdf?view=Standard)
- World Health Organization, 2013. Chronic diseases and health promotion: chronic rheumatic conditions. Secondary Chronic diseases and health promotion: chronic rheumatic conditions. <http://www.who.int/chp/topics/rheumatic/en/>
- Health and Safety Executive, 2012. Annual Statistics Report 2011–12. Secondary Annual Statistics Report 2011–12. <http://www.hse.gov.uk/statistics/overall/hssh1112.pdf>
- HM Government. No health without mental health: a cross-government mental health outcomes strategy for people of all ages. Secondary no health without mental health: a cross-government mental health outcomes strategy for people of all ages 2011. [https://www.gov.uk/government/uploads/system/uploads/attachment\\_data/file/213761/dh\\_124058.pdf](https://www.gov.uk/government/uploads/system/uploads/attachment_data/file/213761/dh_124058.pdf).
- Dionne CE, Dunn KM, Croft PR. Does back pain prevalence really decrease with increasing age? A systematic review. *Age Ageing* 2006;35:229–34.
- Jeffries LJ, Milanese SF, Grimmer-Somers KA. Epidemiology of adolescent spinal pain: a systematic overview of the research literature. *Spine* 2007;32:2630–7.
- Hoy D, Bain C, Williams G, et al. A systematic review of the global prevalence of low back pain. *Arthritis Rheum* 2012;64:2028–37.
- Maniadakis N, Gray A. The economic burden of back pain in the UK. *Pain* 2000;84:95–103.
- National Institute for Health and Clinical Excellence. Low back pain final scope. Secondary low back pain final scope 2007. <https://www.nice.org.uk/guidance/cg88/documents/low-back-pain-final-scope2>
- Arthritis Research UK. Osteoarthritis in general practice. Secondary Osteoarthritis in general practice 2013. [http://www.arthritisresearchuk.org/~media/Files/Policy%20files/Policy%20pages%20files/Keel\\_Data\\_Report\\_digital\\_output.ashx?la=en](http://www.arthritisresearchuk.org/~media/Files/Policy%20files/Policy%20pages%20files/Keel_Data_Report_digital_output.ashx?la=en)
- Oxford Economics. The economic costs of arthritis for the UK economy. Secondary the economic costs of arthritis for the UK economy 2010. <http://www.oxfordeconomics.com/my-oxford/projects/128882>
- Centre HSCI. Health Survey for England 2014: health, social care and lifestyles - Summary of key findings. Secondary health survey for England 2014: health, social care and lifestyles - summary of key findings 2014. <http://www.hscic.gov.uk/catalogue/PUB19295/HSE2014-Sum-bklet.pdf>
- Ohayon MM. Epidemiology of insomnia: what we know and what we still need to learn. *Sleep Med Rev* 2002;6:97–111.
- Salisbury C, Johnson L, Purdy S, et al. Epidemiology and impact of multimorbidity in primary care: a retrospective cohort study. *Br J Gen Pract* 2011;61:e12–e21.
- Violan C, Foguet-Boreu Q, Flores-Mateo G, et al. Prevalence, determinants and patterns of multimorbidity in primary care: a systematic review of observational studies. *PLoS One* 2014;9:e102149.
- Mercer S, Salisbury C, Fortin M. *ABC of Multimorbidity*: John Wiley & Sons, 2014.
- Barnett K, Mercer SW, Norbury M, et al. Epidemiology of multimorbidity and implications for health care, research, and medical education: a cross-sectional study. *Lancet* 2012;380:37–43.
- Katona C, Peveler R, Dowrick C, et al. Pain symptoms in depression: definition and clinical significance. *Clin Med* 2005;5:390–5.
- Fisher P. *Connecting with the Strategic Agenda: integrated medicine, polypharmacy, antimicrobial resistance and multimorbidity CAMSTRAND 2015*.
- The House of Lords. *What is complementary and alternative medicine? Growing use of CAM in the United Kingdom science and technology - sixth report*, 2000.
- Eardley S, Bishop FL, Prescott P, et al. A systematic literature review of complementary and alternative medicine prevalence in EU. *Forsch Complementmed* 2012;19(Suppl 2):18–28.
- Hunt KJ, Coelho HF, Wider B, et al. Complementary and alternative medicine use in England: results from a national survey. *Int J Clin Pract* 2010;64:1496–502.
- Wye L, Shaw A, Sharp D. Designing a 'NHS friendly' complementary therapy service: a qualitative case study. *BMC Health Serv Res* 2008;8:173.
- Fisher P, van Haselen R, Hardy K, et al. Effectiveness gaps: a new concept for evaluating health service and research needs applied to complementary and alternative medicine. *J Altern Complement Med* 2004;10:627–32.
- Unwin J, Peters D. Gatekeepers and the Gateway--a mixed-methods inquiry into practitioners' referral behaviour to the Gateway Clinic. *Acupunct Med* 2009;27:21–5.
- Kessler RC, Soukup J, Davis RB, et al. The use of complementary and alternative therapies to treat anxiety and depression in the United States. *Am J Psychiatry* 2001;158:289–94.
- Mamtani R, Cimino A. A primer of complementary and alternative medicine and its relevance in the treatment of mental health problems. *Psychiatr Q* 2002;73:367–81.
- Artus M, Croft P, Lewis M. The use of CAM and conventional treatments among primary care consultants with chronic musculoskeletal pain. *BMC Fam Pract* 2007;8:26.
- Arksey H, O'Malley L. Scoping studies: towards a methodological framework. *Int J Soc Res Methodol* 2005;8:19–32.
- Mays N, Roberts E, Popay E. Chapter 12: synthesising research evidence. *Methods for studying the delivery and organisation of health services*. London: Routledge, 2001:188–220.
- Levac D, Colquhoun H, O'Brien KK. Scoping studies: advancing the methodology. *Implement Sci* 2010;5:69.
- Daudt HM, van Mossel C, Scott SJ. Enhancing the scoping study methodology: a large, inter-professional team's experience with Arksey and O'Malley's framework. *BMC Med Res Methodol* 2013;13:48.
- Colquhoun HL, Levac D, O'Brien KK, et al. Scoping reviews: time for clarity in definition, methods, and reporting. *J Clin Epidemiol* 2014;67:1291–4.
- Pham MT, Rajić A, Greig JD, et al. A scoping review of scoping reviews: advancing the approach and enhancing the consistency. *Res Synth Methods* 2014;5:371–85.
- Shea BJ, Hamel C, Wells GA, et al. AMSTAR is a reliable and valid measurement tool to assess the methodological quality of systematic reviews. *J Clin Epidemiol* 2009;62:1013–20.
- Moher D, Liberati A, Tetzlaff J, et al. PRISMA Group. Preferred reporting items for systematic reviews and meta-analyses: the PRISMA statement. *BMJ* 2009;339:b2535.
- Furlan AD, Yazdi F, Tsertsvadze A, et al. Complementary and alternative therapies for back pain II. *Evid Rep Technol Assess* 2010;194:1–764.
- Rubinstein SM, van Middelkoop M, Kuijpers T, et al. A systematic review on the effectiveness of complementary and alternative medicine for chronic non-specific low-back pain. *Eur Spine J* 2010;19:1213–28.

42. Dagenais S, Gay RE, Tricco AC, *et al.* NASS Contemporary Concepts in Spine Care: spinal manipulation therapy for acute low back pain. *Spine J* 2010;10:918–40.
43. Rubinstein SM, van Middelkoop M, Assendelft WJ, *et al.* Spinal manipulative therapy for chronic low-back pain: an update of a Cochrane review. *Spine* 2011;36:E825–46.
44. Rubinstein SM, Terwee CB, Assendelft WJ, *et al.* Spinal manipulative therapy for acute low-back pain. *Cochrane Database Syst Rev* 2012;170:CD008880.
45. Merezpeza A. Effects of spinal manipulation versus therapeutic exercise on adults with chronic low back pain: a literature review. *J Can Chiropr Assoc* 2014;58:456–66.
46. Hidalgo B, Detrembleur C, Hall T, *et al.* The efficacy of manual therapy and exercise for different stages of non-specific low back pain: an update of systematic reviews. *J Man Manip Ther* 2014;22:59–74.
47. Tsertsvadze A, Clar C, Court R, *et al.* Cost-effectiveness of manual therapy for the management of musculoskeletal conditions: a systematic review and narrative synthesis of evidence from randomized controlled trials. *J Manipulative Physiol Ther* 2014;37:343–62.
48. Franke H, Fryer G, Ostelo RW, *et al.* Muscle energy technique for non-specific low-back pain. *Cochrane Database Syst Rev* 2015;2:CD009852.
49. Manheimer E, White A, Berman B, *et al.* Meta-analysis: acupuncture for low back pain[correction appears in Ann Intern Med 2005 Jun 7;142(11):950–1; Ann Intern Med 2005;143(9):695]. *Ann Intern Med* 2005;142:651.
50. Xu M, Yan S, Yin X, *et al.* Acupuncture for chronic low back pain in long-term follow-up: a meta-analysis of 13 randomized controlled trials. *Am J Chin Med* 2013;41:1–19.
51. Holtzman S, Beggs RT. Yoga for chronic low back pain: a meta-analysis of randomized controlled trials. *Pain Res Manag* 2013;18:267–72.
52. Pittler MH, Karagülle MZ, Karagülle M, *et al.* Spa therapy and balneotherapy for treating low back pain: meta-analysis of randomized trials. *Rheumatology* 2006;45:880–4.
53. Furlan AD, van Tulder M, Cherkin D, *et al.* Acupuncture and dry-needling for low back pain: an updated systematic review within the framework of the cochrane collaboration. *Spine* 2005;30:944–63.
54. Yuan QL, Guo TM, Liu L, *et al.* Traditional Chinese medicine for neck pain and low back pain: a systematic review and meta-analysis. *PLoS One* 2015;10:e0117146.
55. Posadzki P, Lizi P, Hagner-Derengowska M. Pilates for low back pain: a systematic review. *Complement Ther Clin Pract* 2011;17:85–9.
56. Lim ECW, Poh RLC, Low AY, *et al.* Effects of Pilates-Based Exercises on Pain and Disability in Individuals With Persistent Nonspecific Low Back Pain: A Systematic Review With Meta-analysis. *J Orthop Sports Phys Ther* 2011;41:70–80.
57. Miyamoto GC, Costa LO, Cabral CM. Efficacy of the Pilates method for pain and disability in patients with chronic nonspecific low back pain: a systematic review with meta-analysis. *Braz J Phys Ther* 2013;17:517–32.
58. Aladro-Gonzalvo AR, Araya-Vargas GA, Machado-Díaz M, *et al.* Pilates-based exercise for persistent, non-specific low back pain and associated functional disability: a meta-analysis with meta-regression. *J Bodyw Mov Ther* 2013;17:125–36.
59. Yamato TP, Maher CG, Saragiotto BT, *et al.* Cochrane Back and Neck Group. Pilates for low back pain. *Cochrane Database Syst Rev* 2015;37:CD010265.
60. Posadzki P, Ernst E. Yoga for low back pain: a systematic review of randomized clinical trials. *Clin Rheumatol* 2011;30:1257–62.
61. Ward L, Stebbings S, Cherkin D, *et al.* Yoga for functional ability, pain and psychosocial outcomes in musculoskeletal conditions: a systematic review and meta-analysis. *Musculoskeletal Care* 2013;11:203–17.
62. Cramer H, Lauche R, Haller H, *et al.* A systematic review and meta-analysis of yoga for low back pain. *Clin J Pain* 2013;29:450–60.
63. Furlan AD, Giraldo M, Baskwill A, *et al.* Cochrane Back and Neck Group. Massage for low-back pain. *Cochrane Database Syst Rev* 2015;18. Art. No.: CD001929.
64. Orrock PJ, Myers SP. Osteopathic intervention in chronic non-specific low back pain: a systematic review. *BMC Musculoskelet Disord* 2013;14:129.
65. Franke H, Franke JD, Fryer G. Osteopathic manipulative treatment for nonspecific low back pain: a systematic review and meta-analysis. *BMC Musculoskelet Disord* 2014;15:286.
66. Kim YC, Lee MS, Park E-S, *et al.* Acupressure for the Treatment of Musculoskeletal Pain Conditions: A Systematic Review. *J Musculoskelet Pain* 2012;20:116–21.
67. Hu XY, Chen NN, Chai QY, *et al.* Integrative treatment for low back pain: An exploratory systematic review and meta-analysis of randomized controlled trials. *Chin J Integr Med* 2015. Accepted for publication.
68. Kizhakkevettill A, Rose K, Kadar GE. Integrative therapies for low back pain that include complementary and alternative medicine care: a systematic review. *Glob Adv Health Med* 2014;3:49–64.
69. Walker BF, French SD, Grant W, *et al.* Combined chiropractic interventions for low-back pain. *Cochrane Database Syst Rev* 2010;29:CD005427. N.PAG.
70. Langhorst J, Klose P, Musial F, *et al.* Efficacy of acupuncture in fibromyalgia syndrome--a systematic review with a meta-analysis of controlled clinical trials. *Rheumatology* 2010;49:778–88.
71. Cao H, Liu J, Lewith GT. Traditional Chinese Medicine for treatment of fibromyalgia: a systematic review of randomized controlled trials. *J Altern Complement Med* 2010;16:397–409.
72. Deare JC, Zheng Z, Xue CCL, *et al.* Acupuncture for treating fibromyalgia. *Cochrane Database Syst Rev* 2013;143:CD007070.
73. Yang B, Yi G, Hong W, *et al.* Efficacy of acupuncture on fibromyalgia syndrome: a meta-analysis. *J Tradit Chin Med* 2014;34:381–91.
74. Porter NS, Jason LA, Boulton A, *et al.* Alternative medical interventions used in the treatment and management of myalgic encephalomyelitis/chronic fatigue syndrome and fibromyalgia. *J Altern Complement Med* 2010;16:235–49.
75. Li YH, Wang FY, Feng CQ, *et al.* Massage therapy for fibromyalgia: a systematic review and meta-analysis of randomized controlled trials. *PLoS One* 2014;9:e89304.
76. Yuan SL, Matsutani LA, Marques AP. Effectiveness of different styles of massage therapy in fibromyalgia: a systematic review and meta-analysis. *Man Ther* 2015;20:257–64.
77. Lauche R, Cramer H, Dobos G, *et al.* A systematic review and meta-analysis of mindfulness-based stress reduction for the fibromyalgia syndrome. *J Psychosom Res* 2013;75:500–10.
78. Theadom A, Cropley M, Smith HE, *et al.* Cochrane Musculoskeletal Group. Mind and body therapy for fibromyalgia. *Cochrane Database Syst Rev* 2015;13:CD001980.
79. Bernardy K, Füber N, Klose P, *et al.* Efficacy of hypnosis/guided imagery in fibromyalgia syndrome--a systematic review and meta-analysis of controlled trials. *BMC Musculoskelet Disord* 2011;12:133.
80. Mist SD, Firestone KA, Jones KD. Complementary and alternative exercise for fibromyalgia: a meta-analysis. *J Pain Res* 2013;6:247–60.
81. Cao H, Li X, Han M, *et al.* Acupoint stimulation for fibromyalgia: a systematic review of randomized controlled trials. *Evid Based Complement Alternat Med* 2013;2013:1–15.
82. Gross A, Langevin P, Burnie SJ, *et al.* Cochrane Back and Neck Group. Manipulation and mobilisation for neck pain contrasted against an inactive control or another active treatment. *Cochrane Database Syst Rev* 2015;17.
83. Carlesso LC, Gross AR, Santaguida PL, *et al.* Adverse events associated with the use of cervical manipulation and mobilization for the treatment of neck pain in adults: a systematic review. *Man Ther* 2010;15:434–44.
84. Lin JH, Chiu TT, Hu J. Chinese manipulation for mechanical neck pain: a systematic review. *Clin Rehabil* 2012;26:963–73.
85. Huisman PA, Speksnijder CM, de Wijer A. The effect of thoracic spine manipulation on pain and disability in patients with non-specific neck pain: a systematic review. *Disabil Rehabil* 2013;35:1677–85.
86. Boyles R, Toy P, Mellon J, *et al.* Effectiveness of manual physical therapy in the treatment of cervical radiculopathy: a systematic review. *J Man Manip Ther* 2011;19:135–42.
87. Miller J, Gross A, D'Sylva J, *et al.* Manual therapy and exercise for neck pain: a systematic review. *Man Ther* 2010;15:334–54.
88. D'Sylva J, Miller J, Gross A, *et al.* Cervical Overview Group. Manual therapy with or without physical medicine modalities for neck pain: a systematic review. *Man Ther* 2010;15:415–33.
89. Trinh KV, Graham N, Gross AR, *et al.* Cervical Overview Group. Acupuncture for neck disorders. *Cochrane Database Syst Rev* 2006:CD004870.
90. Patel KC, Gross A, Graham N, *et al.* Massage for mechanical neck disorders. *Cochrane Database Syst Rev* 2012;11:CD004871.
91. White A, Foster NE, Cummings M, *et al.* Acupuncture treatment for chronic knee pain: a systematic review. *Rheumatology* 2007;46:384–90.

92. Manheimer E, Cheng K, Linde K, *et al*. Cochrane Musculoskeletal Group. Acupuncture for peripheral joint osteoarthritis. *Cochrane Database Syst Rev* 2010;38:CD001977.
93. Cao L, Zhang XL, Gao YS, *et al*. Needle acupuncture for osteoarthritis of the knee. A systematic review and updated meta-analysis. *Saudi Med J* 2012;33:526–32.
94. Corbett MS, Rice SJ, Madurasinghe V, *et al*. Acupuncture and other physical treatments for the relief of pain due to osteoarthritis of the knee: network meta-analysis. *Osteoarthritis Cartilage* 2013;21:1290–8.
95. Turner K, Igo S. Efficacy of acupuncture in the treatment of osteoarthritis of the knee joint. *Journal - Acupuncture Association of Chartered Physiotherapists* 2013;2013:21–39.
96. Manyanga T, Froese M, Zarychanski R, *et al*. Pain management with acupuncture in osteoarthritis: a systematic review and meta-analysis. *BMC Complement Altern Med* 2014;14:312.
97. Hall A, Maher C, Latimer J, *et al*. The effectiveness of Tai Chi for chronic musculoskeletal pain conditions: a systematic review and meta-analysis. *Arthritis Rheum* 2009;61:717–24.
98. Kang JW, Lee MS, Posadzki P, *et al*. T'ai chi for the treatment of osteoarthritis: a systematic review and meta-analysis. *BMJ Open* 2011;1:e000035.
99. Lauche R, Langhorst J, Dobos G, *et al*. A systematic review and meta-analysis of Tai Chi for osteoarthritis of the knee. *Complement Ther Med* 2013;21:396–406.
100. Yan JH, Gu WJ, Sun J, *et al*. Efficacy of Tai Chi on pain, stiffness and function in patients with osteoarthritis: a meta-analysis. *PLoS One* 2013;8:e61672.
101. Mat S, Tan MP, Kamaruzzaman SB, *et al*. Physical therapies for improving balance and reducing falls risk in osteoarthritis of the knee: a systematic review. *Age Ageing* 2015;44:16–24.
102. Casimiro L, Barnsley L, Brosseau L, *et al*. Acupuncture and electroacupuncture for the treatment of rheumatoid arthritis. *Cochrane Database Syst Rev* 2005;38:CD003788.
103. Lee MS, Shin BC, Ernst E. Acupuncture for rheumatoid arthritis: a systematic review. *Rheumatology* 2008;47:1747–53.
104. Sun Z-ling, Xu X, Du S-zheng, *et al*. Moxibustion for treating rheumatoid arthritis: A systematic review and meta-analysis of randomized controlled trials. *Eur J Integr Med* 2014;6:621–30.
105. Verhagen AP, Bierma-Zeinstra SMA, Boers M, *et al*. Cochrane Musculoskeletal Group. Balneotherapy (or spa therapy) for rheumatoid arthritis. *Cochrane Database Syst Rev* 2015;25:CD000518.
106. Law D, McDonough S, Bleakley C, *et al*. Laser acupuncture for treating musculoskeletal pain: a systematic review with meta-analysis. *J Acupunct Meridian Stud* 2015;8:2–16.
107. Zhao HJ, Tan JY, Wang T, *et al*. Auricular therapy for chronic pain management in adults: A synthesis of evidence. *Complement Ther Clin Pract* 2015;21:68–78.
108. Wong Lit Wan D, Wang Y, Xue CC, *et al*. Local and distant acupuncture points stimulation for chronic musculoskeletal pain: A systematic review on the comparative effects. *Eur J Pain* 2015;19:1232–47.
109. Song Y, Lu H, Chen H, *et al*. Mindfulness intervention in the management of chronic pain and psychological comorbidity: A meta-analysis. *Int J Nurs Sci* 2014;1:215–23.
110. Bawa FL, Mercer SW, Atherton RJ, *et al*. Does mindfulness improve outcomes in patients with chronic pain? Systematic review and meta-analysis. *Br J Gen Pract* 2015;65:e387–e400.
111. Braun C, Hanchard NCA. Manual therapy and exercise for impingement-related shoulder pain. *Physical Therapy Reviews* 2010;15:62–83.
112. Page Matthew J, Green S, Kramer S, *et al*. Manual therapy and exercise for adhesive capsulitis (frozen shoulder). *Cochrane Database of Systematic Reviews* 2014;8:CD011275.
113. Clar C, Tsertsvadze A, Court R, *et al*. Clinical effectiveness of manual therapy for the management of musculoskeletal and non-musculoskeletal conditions: systematic review and update of UK evidence report. *Chiropr Man Therap* 2014;22:12.
114. Desjardins-Charbonneau A, Roy JS, Dionne CE, *et al*. The efficacy of manual therapy for rotator cuff tendinopathy: a systematic review and meta-analysis. *J Orthop Sports Phys Ther* 2015;45:330–50.
115. Green S, Buchbinder R, Hetrick SE. Cochrane Musculoskeletal Group. Acupuncture for shoulder pain. *Cochrane Database Syst Rev* 2005;7:CD005319.
116. Tough EA, White AR, Cummings TM, *et al*. Acupuncture and dry needling in the management of myofascial trigger point pain: a systematic review and meta-analysis of randomised controlled trials. *Eur J Pain* 2009;13:3–10.
117. Liu L, Huang QM, Liu QG, *et al*. Effectiveness of dry needling for myofascial trigger points associated with neck and shoulder pain: a systematic review and meta-analysis. *Arch Phys Med Rehabil* 2015;96:944–55.
118. Clark RJ, Tighe M. The effectiveness of acupuncture for plantar heel pain: a systematic review. *Acupunct Med* 2012;30:298–306.
119. Qin ZL, Wu Xiaoxu, Zhai Jiani, *et al*; *Acupuncture for sciatica: a systematic review*, 2015.
120. Bisset L, Hing W, Vicenzino B. A systematic review of the efficacy of MWM. In: Bill Vicenzino WH, Rivett D, Hall T, eds. *Mobilisation with Movement: The Art and Science*. Australia: Elsevier, 2011:26–63.
121. Tang H, Fan H, Chen J, *et al*. Acupuncture for Lateral Epicondylitis: A Systematic Review. *Evid Based Complement Alternat Med* 2015;2015:1–13.
122. Southerst D, Marchand AA, Côté P, *et al*. The effectiveness of noninvasive interventions for musculoskeletal thoracic spine and chest wall pain: a systematic review by the Ontario Protocol for Traffic Injury Management (OPTiMa) collaboration. *J Manipulative Physiol Ther* 2015;38:521–31.
123. Piper S, Shearer HM, Côté P, *et al*. The effectiveness of soft-tissue therapy for the management of musculoskeletal disorders and injuries of the upper and lower extremities: A systematic review by the Ontario Protocol for Traffic Injury management (OPTiMa) collaboration. *Man Ther* 2016;21:18–34.
124. de Vibe M, Arlid B, Elizabeth T, *et al*. Mindfulness Based Stress Reduction (MBSR) for Improving Health, Quality of Life. *and Social Functioning in Adults* 2012.
125. Goyal M, Singh S, Sibinga EM, *et al*. Meditation programs for psychological stress and well-being: a systematic review and meta-analysis. *JAMA Intern Med* 2014;174:357–68.
126. Smith CA, Hay PPJ, MacPherson H; Cochrane Common Mental Disorders Group. Acupuncture for depression. *Cochrane Database Syst Rev* 2010;9:CD004046.
127. Zhang ZJ, Chen HY, Yip KC, *et al*. The effectiveness and safety of acupuncture therapy in depressive disorders: systematic review and meta-analysis. *J Affect Disord* 2010;124(1-2):9–21.
128. Wang H, Qi H, Wang BS, *et al*. Is acupuncture beneficial in depression: a meta-analysis of 8 randomized controlled trials? *J Affect Disord* 2008;111(2-3):125–34.
129. Klainin-Yobas P, Oo WN, Suzanne Yew PY, *et al*. Effects of relaxation interventions on depression and anxiety among older adults: a systematic review. *Aging Ment Health* 2015;19:1043–55.
130. Cramer H, Lauche R, Langhorst J, *et al*. Yoga for depression: a systematic review and meta-analysis. *Depress Anxiety* 2013;30:1068–83.
131. Maratos A, Gold C, Wang X, *et al*. Music therapy for depression. *Cochrane Database Syst Rev* 2008;25:CD004517.
132. Liu X, Clark J, Siskind D, *et al*. A systematic review and meta-analysis of the effects of Qigong and Tai Chi for depressive symptoms. *Complement Ther Med* 2015;23:516–34.
133. Joyce J, Herbison GP. Reiki for depression and anxiety. *Cochrane Database Syst Rev* 2015:CD006833 <http://onlinelibrary.wiley.com/doi/10.1002/14651858.CD006833.pub2/abstracthttp://onlinelibrary.wiley.com/store/10.1002/14651858.CD006833.pub2/asset/CD006833.pdf?v=1&t=ib3g7f71&s=fc7f57607f3f9ab6e5f4c9efed82890cbf11111>.
134. Chi I, Jordan-Marsh M, Guo M, *et al*. Tai chi and reduction of depressive symptoms for older adults: a meta-analysis of randomized trials. *Geriatr Gerontol Int* 2013;13:3–12.
135. Kirkwood G, Rampes H, Tuffrey V, *et al*. Yoga for anxiety: a systematic review of the research evidence. including commentary by Ramaratnam S. *British journal of sports medicine* 2005;39:884–91. discussion 91 doi:[published Online First: Epub Date].
136. Chugh-Gupta N, Baldassarre FG, Vrkljan BH. A systematic review of yoga for state anxiety: considerations for occupational therapy. *Can J Occup Ther* 2013;80:150–70.
137. Krisanaprakornkit T, Krisanaprakornkit W, Piyavhatkul N, *et al*. Meditation therapy for anxiety disorders. *Cochrane Database Syst Rev* 2006;1:CD004998.
138. Chen KW, Berger CC, Manheimer E, *et al*. Meditative therapies for reducing anxiety: a systematic review and meta-analysis of randomized controlled trials. *Depress Anxiety* 2012;29:545–62.
139. Pilkington K, Kirkwood G, Rampes H, *et al*. Acupuncture for anxiety and anxiety disorders--a systematic literature review. *Acupunct Med* 2007;25(1-2):1–10.
140. Wang CW, Chan CH, Ho RT, Rt H, *et al*. Managing stress and anxiety through qigong exercise in healthy adults: a systematic review and meta-analysis of randomized controlled trials. *BMC Complement Altern Med* 2014;14:8.
141. Wu H, Yu D, He Y, *et al*. Morita therapy for anxiety disorders in adults. *Cochrane Database Syst Rev* 2015;2:CD008619.

142. Robinson J, Biley FC, Dolk H. Therapeutic touch for anxiety disorders. *Cochrane Database Syst Rev* 2007;4:CD006240.
143. Wu WW, Kwong E, Lan XY, W-w W, X-y L, *et al.* The Effect of a Meditative Movement Intervention on Quality of Sleep in the Elderly: A Systematic Review and Meta-Analysis. *J Altern Complement Med* 2015;21.
144. Neuendorf R, Wahbeh H, Chamine I, *et al.* The Effects of Mind-Body Interventions on Sleep Quality: A Systematic Review. *Evidence-Based Complementary and Alternative Medicine* 2015;2015:1–17.
145. Wang F, Eun-Kyoung Lee O, Feng F, *et al.* The effect of meditative movement on sleep quality: A systematic review. *Sleep Med Rev* 2016;30:43–52.
146. Huang W, Kutner N, Bliwise DL. A systematic review of the effects of acupuncture in treating insomnia. *Sleep Med Rev* 2009;13:73–104.
147. Lan Y, Wu X, Tan HJ, *et al.* Auricular acupuncture with seed or pellet attachments for primary insomnia: a systematic review and meta-analysis. *BMC Complement Altern Med* 2015;15:103.
148. Du S, Dong J, Zhang H, *et al.* Taichi exercise for self-rated sleep quality in older people: a systematic review and meta-analysis. *Int J Nurs Stud* 2015;52:368–79.
149. Cooper KL, Relton C. Homeopathy for insomnia: a systematic review of research evidence. *Sleep Med Rev* 2010;14:329–37.
150. Lam TH, Chung KF, Yeung WF, *et al.* Hypnotherapy for insomnia: a systematic review and meta-analysis of randomized controlled trials. *Complement Ther Med* 2015;23:719–32.
151. Wieland LS, Skoetz N, Pilkington K, *et al.* Yoga treatment for chronic non-specific low back pain. *Cochrane Database Syst Rev* 2017;1:CD010671.
152. Hilton L, Hempel S, Ewing BA, *et al.* Mindfulness Meditation for Chronic Pain: Systematic Review and Meta-analysis. *Ann Behav Med* 2017;51:199–213.
153. Anheyer D, Haller H, Barth J, *et al.* Mindfulness-Based Stress Reduction for Treating Low Back Pain: A Systematic Review and Meta-analysis. *Ann Intern Med* 2017;166:799–807.
154. Shim JW, Jung JY, Kim SS. Effects of Electroacupuncture for Knee Osteoarthritis: A Systematic Review and Meta-Analysis. *Evid Based Complement Alternat Med* 2016;2016:1–18.
155. Lin X, Huang K, Zhu G, *et al.* The Effects of Acupuncture on Chronic Knee Pain Due to Osteoarthritis: A Meta-Analysis. *J Bone Joint Surg Am* 2016;98:1578–85.
156. Shea BJ, Reeves BC, Wells G, *et al.* AMSTAR 2: a critical appraisal tool for systematic reviews that include randomised or non-randomised studies of healthcare interventions, or both. *BMJ* 2017;358:j4008.
157. Linde K. Can you trust systematic reviews of complementary and alternative therapies? *Eur J Integr Med* 2009;1:117–23.
158. Verhoef MJ, Lewith G, Ritenbaugh C, *et al.* Complementary and alternative medicine whole systems research: beyond identification of inadequacies of the RCT. *Complement Ther Med* 2005;13:206–12.
159. Carter B. Methodological issues and complementary therapies: researching intangibles? *Complement Ther Nurs Midwifery* 2003;9:133–9.