

# BMJ Open Molecular diagnosis of inherited peripheral neuropathies by targeted next-generation sequencing: molecular spectrum delineation

Juliette Bacquet,<sup>1</sup> Tanya Stojkovic,<sup>2</sup> Amandine Boyer,<sup>1</sup> Nathalie Martini,<sup>1</sup> Frédérique Audic,<sup>3</sup> Brigitte Chabrol,<sup>3</sup> Emmanuelle Salort-Campana,<sup>4,5</sup> Emilien Delmont,<sup>4</sup> Jean-Pierre Desvignes,<sup>5</sup> Annie Verschuere,<sup>4</sup> Shahram Attarian,<sup>4,5</sup> Annabelle Chausse,<sup>6</sup> Valérie Delague,<sup>5</sup> Nicolas Levy,<sup>1,5</sup> Nathalie Bonello-Palot<sup>1,5</sup>

**To cite:** Bacquet J, Stojkovic T, Boyer A, *et al.* Molecular diagnosis of inherited peripheral neuropathies by targeted next-generation sequencing: molecular spectrum delineation. *BMJ Open* 2018;**8**:e021632. doi:10.1136/bmjopen-2018-021632

► Prepublication history and additional material for this paper are available online. To view these files, please visit the journal online (<http://dx.doi.org/10.1136/bmjopen-2018-021632>).

Received 11 January 2018  
Revised 30 May 2018  
Accepted 12 September 2018



© Author(s) (or their employer(s)) 2018. Re-use permitted under CC BY-NC. No commercial re-use. See rights and permissions. Published by BMJ.

For numbered affiliations see end of article.

## Correspondence to

Dr Juliette Bacquet;  
[juliette.bacquet@ap-hm.fr](mailto:juliette.bacquet@ap-hm.fr)

## ABSTRACT

**Purpose** Inherited peripheral neuropathies (IPN) represent a large heterogeneous group of hereditary diseases with more than 100 causative genes reported to date. In this context, targeted next-generation sequencing (NGS) offers the opportunity to screen all these genes with high efficiency in order to unravel the genetic basis of the disease. Here, we compare the diagnostic yield of targeted NGS with our previous gene by gene Sanger sequencing strategy. We also describe several novel likely pathogenic variants.

**Design and participants** We have completed the targeted NGS of 81 IPN genes in a cohort of 123 unrelated patients affected with diverse forms of IPNs, mostly Charcot-Marie-Tooth disease (CMT): 23% CMT1, 52% CMT2, 9% distal hereditary motor neuropathy, 7% hereditary sensory and autonomic neuropathy and 6.5% intermediate CMT.

**Results** We have solved the molecular diagnosis in 49 of 123 patients (~40%). Among the identified variants, 26 variants were already reported in the literature. In our cohort, the most frequently mutated genes are respectively: *MFN2*, *SH3TC2*, *GDAP1*, *NEFL*, *GAN*, *KIF5A* and *AARS*. Panel-based NGS was more efficient in familial cases than in sporadic cases (diagnostic yield 49% vs 19%, respectively). NGS-based search for copy number variations, allowed the identification of three duplications in three patients and raised the diagnostic yield to 41%. This yield is two times higher than the one obtained previously by gene Sanger sequencing screening. The impact of panel-based NGS screening is particularly important for demyelinating CMT (CMT1) subtypes, for which the success rate reached 87% (36% only for axonal CMT2).

**Conclusion** NGS allowed to identify causal mutations in a shorter and cost-effective time. Actually, targeted NGS is a well-suited strategy for efficient molecular diagnosis of IPNs. However, NGS leads to the identification of numerous variants of unknown significance, which interpretation requires interdisciplinary collaborations between molecular geneticists, clinicians and (neuro)pathologists.

## Strengths and limitations of this study

- First study concerning patients with inherited peripheral neuropathies in South of France.
- New single-nucleotide variation mutations important for the scientific and medical community.
- New copy number variations detected by the analysis of next-generation sequencing data.
- Lack of power (123 patients).
- Lack of functional validation of variants of unknown significance.

## INTRODUCTION

We defined three main categories of inherited peripheral neuropathy (IPN): hereditary motor and sensory neuropathy, more commonly called Charcot-Marie-Tooth disease (CMT), hereditary sensory and autonomic neuropathy (HSAN) and distal hereditary motor neuropathy (dHMN).

With a prevalence of 1/2500,<sup>1</sup> CMT is one of the most frequent cause of neurological disability, characterised by extensive phenotypic and genetic heterogeneity, with all modes of inheritance described. Based on histopathological and electrophysiological criteria, CMT are further categorised into CMT type 1, or demyelinating type (CMT1/HMSN1), and CMT type 2, or axonal type (CMT2/HMSN2). While CMT type 1 is associated with reduced nerve conduction velocities, correlated to decreased myelination of the peripheral nerves, CMT type 2 is characterised by decreased amplitudes of motor and sensory nerve action potentials, related to primarily axonal loss in the peripheral nerve fibres. Patients with both signs of demyelination and axonal degeneration are diagnosed with intermediate CMT, although there is

much controversy about the exact definition of intermediate CMT.<sup>2</sup>

The disease usually starts in childhood or the teenage years and it generally aggravates slowly and progressively. Clinically, most patients present distal motor and sensory weakness associated with feet deformations (most frequently pes cavus) and sometimes other skeletal deformations such as scoliosis. Additional manifestations may exist such as deafness, optic atrophy or pyramidal signs. The disease is characterised by high clinical heterogeneity, both intrafamilial and interfamilial. Concerning the latter, the severity of the disease is highly variable, ranging from almost asymptomatic adult patients to severely disabled children.

Genetically, the disease is also highly heterogeneous, with more than 100 defective genes reported today.<sup>3 4</sup> In our laboratory, before the advent of next-generation sequencing (NGS), the screening strategy was based on Sanger sequencing of candidate genes, except for the *PMP22* duplication, following the recommendations of the French Network of Molecular Diagnosis Laboratories for NeuroMuscular Diseases (<http://www.anpgm.fr/>), who had set up decision trees based on data from the literature. The screening of one specific gene was determined by clinical, genealogical and electrophysiological criteria. For all patients, the 1.5 Mb duplication at chromosome 17p11.2 encompassing the *PMP22* gene, the most frequent cause of CMT, was first achieved by multiplex ligation-dependent probe amplification (MLPA). In negative cases, a subsequent gene by gene sequential screening was carried out, and the screened genes were different whether the patient presented demyelinating or axonal CMT. For patients with autosomal-dominant demyelinating CMT (CMT1), the screening of *GJB1*, the gene encoding connexin 32, was then carried out through Sanger sequencing, followed by the sequential screening of: *MPZ*, *LITAF/SIMPLE*, *NEFL*, *GDAPI* and *EGR2*. In autosomal-dominant axonal CMT (CMT2), the patients were subjected to Sanger sequencing screening of the following genes: *GJB1*, *MFN2* (especially if the patient show pyramidal signs or optic atrophy), then *MPZ* and *NEFL*.

In the last few years, the molecular diagnosis improved with the advent of NGS, which is now the strategy used in routine in our laboratory, in patients negative for the *PMP22* duplication. The list of genes involved in IPNs is constantly rising, and is now above 100.

Our objective is to compare the diagnostic yield of a targeted NGS strategy (panel of 81 IPN/CMT genes) with the previous gene by gene Sanger sequencing strategy, by comparing the results of NGS in a cohort of 123 patients and the results of Sanger sequencing in a group of 56 patients. We compare the molecular diagnostic resolution rates between demyelinating and axonal CMT forms. We also report on new likely pathogenic variants, not yet described in the literature, and we present the most frequently mutated genes in our group. Finally, we describe two candidate copy number variations (CNVs), identified from the NGS data.

## MATERIALS AND METHODS

### Clinical data

For NGS screening, we have prospectively included 123 index cases affected with hereditary motor and sensory neuropathy, dHMN and HSAN, seen in consultation in the Neuromuscular Disease Reference Center, since 2015.

For Sanger sequencing, we have studied a retrospective cohort of 56 patients seen in consultation between 2012 and 2014.

### PATIENT AND PUBLIC INVOLVEMENT

Patients and/or public were involved neither in the design of this study protocol nor in the development of the research question. Patients and/or public will not be involved in the recruitment process. The results of this study will be presented at the next FILNEMUS conference.

### STATISTICAL ANALYSIS

We have compared the targeted NGS strategy to the previous gene analysis by Sanger sequencing on a retrospective cohort of 56 patients seen between 2012 and 2014 using a two-tailed Fisher's exact test.

### GENETIC STUDIES

#### Samples

A written and signed approval has been collected from the patients in accordance with French recommendations as well as in agreement with the local ethics committee rules.

DNA was extracted from peripheral blood using standard procedures. DNAs were prepared and stored at the accredited Biological Resource Centre (CRB TAC component (NF S96-900 and ISO 9001v2015 Certification) Department of Medical Genetics, Timone Hospital of Marseille (APHM). All DNAs belong to a biological sample collection declared to the French Ministry of Health (declaration number DC-2008-429) whose use for research purposes was authorised by the French ministry of Health (authorisation number AC-2011-1312 and AC-2017-2986).

All patients were searched for the *PMP22* duplication before including them in the NGS analysis protocol. To test for the *PMP22* duplication, we used an MLPA protocol (MRC-Holland) following the manufacturer's recommendations.

### NEXT-GENERATION SEQUENCING

#### Library preparation and sequencing

Libraries were prepared using the ClearSeq Inherited Disease Panel from Agilent (Santa Clara, California, USA), which enables the capturing of 2742 genes known to cause inherited disorders. The coding regions and flanking intronic regions of the 2742 genes were

enriched, in solution, using the SureSelect Target Enrichment System from Agilent (Santa Clara, California, USA), following the manufacturer recommendations.

For sequencing, we used the Ion Proton platform (Thermo Fisher Scientific, USA).

### Computational analysis

After capturing enrichment and sequencing, raw data were converted to Fastq files, aligned to the reference sequence of the human genome (University of California Santa Cruz (UCSC) hg19/GRCh37), and annotated, using the Ion Proton platform integrated workflow (Thermo Fisher Scientific, USA). The obtained variant call format and binary alignment map (BAM) files were used for variant search as described below.

In a diagnosis settings, we realised a targeted analysis of the NGS data, focusing on a list of 81 IPN genes (see online supplementary table 1). The data were filtered, using an 'in-house' tool for variant annotation and Filtering Variant Annotation and Filter Tool (<https://varaft.eu/download.php>):<sup>5</sup> (1) variants with allele frequencies <1% in the Exome Aggregation Consortium (ExAC) data set (<http://exac.broadinstitute.org/>) were removed, (2) the remaining variants were filtered based on their type and genomic localisation, thus, synonymous, intronic, variants in intergenic, 3' and 5' Untranslated Region (UTR) regions were discarded. In order to predict the deleterious effect of the identified sequence variations, different bioinformatics tools were applied, such as MutationTaster (<http://www.mutationtaster.org/>),<sup>6</sup> Sorting Intolerant From Tolerant (SIFT) (<http://sift.bii.a-star.edu.sg/>),<sup>7</sup> PolyPhen-2 (<http://genetics.bwh.harvard.edu/pph2/>)<sup>8</sup> and Universal Mutation Database (UMD) predictor (<http://umd-predictor.eu/>).<sup>9</sup>

In order to check whether variants had already been inventoried and classified, we looked up in Online Mendelian Inheritance in Man (OMIM) (<https://www.ncbi.nlm.nih.gov/omim>), ClinVar (<https://www.ncbi.nlm.nih.gov/clinvar/>) and HGMD (<http://www.hgmd.cf.ac.uk>). Finally, thanks to the American College of Medical Genetics and Genomics (ACMG) recommendations (Richards classification, 2015)<sup>10</sup> and to in silico analyses, we classified these variants into five categories: pathogenic, probably pathogenic, variant of unknown significance (VUS), probably benign and benign.

### Copy number variations

In order to identify CNV in our cohort, we used ExomeDepth.<sup>11</sup> This tool uses the BAM files from the NGS sequencing run as well as the bed file containing the target regions to be studied (81 genes, see online supplementary table 1). This tool allows to compare the reading depth of our patients' BAM files, to a set of reference samples, in order to eliminate the capturing and sequencing mistakes. Variations in the depth of sequencing are linearly correlated to the variation of the copy number. Deletions and duplications found by ExomeDepth were confirmed by a quantitative (Q)-PCR

(Applied Biosystems ViiA 7 Real-Time PCR System).<sup>12</sup> The CNV analysis of *TFG* and *FGD4* genes was performed using a commercial kit TaqMan Copy Number Assays (HS00918833-CN) by real-time PCR multiplex. Each analysis was done in triplicate.

### Sanger sequencing

Variants found by NGS were confirmed and segregated by Sanger sequencing. In our previous sequential gene by gene analysis, the complete coding sequence of genes were PCR amplified and sequenced by fluorescent Sanger sequencing.

In both situations, genomic and cDNA sequences of the genes were obtained from the UCSC Genomic Browser, February 2009, human reference sequence (GRCh37). Primers used for PCR amplification were designed using Primer3 software (<http://frodo.wi.mit.edu>) to amplify the region surrounding the candidate DNA variations.

PCR products were purified by mixing with a volume ratio (1/8) (36  $\mu$ L) of AMPure beads (Beckman Coulter, USA) according to the manufacturer's instructions and both strands were sequenced using the Big Dye Terminator V.1.1 Cycle Sequencing Kit (Applied Biosystems). Sequence reactions were purified on Sephadex G50 (Amersham Pharmacia Biotech, Foster City, California, USA) and capillary electrophoresis was performed on Genetic Analyser ABI3500XL (Life Technologies, USA). Electrophoregrams were analysed on the Sequence Analysis Software V.5.2 (Applied Biosystems) and aligned with the reference sequence using Sequencher V.5.4.6.

## RESULTS

Among the 123 patients, 28 (23%) had CMT type 1, 64 (52%) had CMT type 2, 11 (9%) had dHMN, 9 (7%) had HSAN, 8 (6,5%) had intermediary CMT and 3 (2,5%) had CMT without clinical information. **Table 1** sums up the phenotypic features of the patients for whom we identified a mutation or a candidate variant (**table 1A** for patients with CMT1, **table 1B** for patients with CMT2 and **table 1C** for patients with others forms of IPN).

The average depth obtained in our panel of 81 genes was 196X and the average coverage at 20X was 98%, with a weaker coverage at 20X for *SOX10*, *INF2* and *CTDP1* genes (81%, 86% and 86% respectively).

Through targeted NGS of 81 IPN genes (see online supplementary table 1), we found one or several potentially pathogenic variants in 60 patients from our cohort of 123 index cases, thereby defining a success rate of 49%. The average age was 20 for positive cases, while it was significantly higher (45 years old) for negative cases.

More precisely, we found a pathogenic variant for 49 patients (40%) and a potentially pathogenic variant for 11 patients (9%). Thirty-seven per cent of cases showed a recessive transmission and 63% a dominant transmission. Twenty-six variants were reported in the literature, whereas 52 variants were never reported. Among these 52

**Table 1A** Clinical and electrophysiological features from 19 IPN index patients with CMT1, for whom a candidate variant was identified

Patient	Disease	Gene	Gender	Age at onset	Deep tendon reflexes	Foot deformities	Muscular weakness and wasting of distal muscles	Sensory loss	Other clinical signs	Median nerve motor (m/s)	Median nerve MNCV (mV)	Median nerve distal CMAP (mV)	Nerve biopsy
2	CMT1 AR	SH3TC2	M	3 years	-	NA	+	+		NA	NA	NA	-
3	CMT1 SPO	GJB1	M	19 years	-	+	+	-	Scoliosis	36	NR	NR	-
5	CMT1 SPO	INF2	F	6 years	-	+	+	+	Deafness+ diaphragmatic palsy	NA	NA	NA	-
9	CMT1 AD	DNM2	M	22 years	-	+	+	-		35	NA	NA	-
10	CMT1 AD	NEFL	F	60 years	NA	+	+	+	Cerebellar ataxia	NA	NA	NA	NA
12	CMT1 AD	LMNA	F	22 months	-	-	-	-		NR*	NR*	NR*	-
14	CMT1 AR	SH3TC2	M	44 years	-	+	+	+		21	NA	NA	-
15	CMT1 AD	LITAF/SIMPLE	M	23 years	-	+	-	+		23	5.7	5.7	-
16	CMT1 AD	GJB1	F	15 years	-	NA	+	+		36	NA	NA	Onion bulbs
17	CMT1 AR	GDAP1	F	8 months	-	+	+	-	Kyphosis	39	1.02	1.02	-
18	CMT1 AD	AARS	M	43 years	-	+	+	+		25	NA	NA	-
20	CMT1 AR	FIG4	F	12 years	-	+	+	-		26	3.6	3.6	NA
21	CMT1 AD	TRPV4 LRSAM1 KIF1B	M	10 years	-	+	+	+	Scoliosis	25	NA	NA	NA
28	CMT1 AD	LITAF DCTN1	F	20 months	+	+	+	-		33	3.9	3.9	NA
32	CMT1 SPO	NEFL	M	2 years	-	+	+	-	Sensorineural hearing loss	29	4.42	4.42	-
34	CMT SPO	KIF1B	F	2.5 years	-	+	-	-		NA	NA	NA	-
41	CMT1 AR	SH3TC2	F	3 years	-	+	-	-	Scoliosis	33	1.04	1.04	-
45	CMT1 AD	INF2	F	3 years	-	+	-	-		21	5.2	5.2	-
47	CMT1 AR	HK1	M	8 years	-	+	+	+	Scoliosis	22	NA	NA	-

\*ENMG show chronic denervation.

+, presence; -, absence; AD, autosomal-dominant; AR, autosomal recessive; CMAP, compound muscle action potential; CMT1, Charcot-Marie-Tooth disease type 1; ENMG, Electroneuromyogram; F, female; IPN, inherited peripheral neuropathy; MNCV, motor nerve conduction velocity; NA, not available; NR, not recordable; M, male; SPO, sporadic.



**Table 1B** Clinical and electrophysiological features from 27 IPN index patients with CMT2, for whom a candidate variant was identified

Patient	Disease	Gene	Gender	Age at onset	Deep tendon reflexes	Foot deformities	Muscular weakness and wasting of distal muscles	Sensory loss	Other clinical signs	Median nerve motor MNCV (m/s)	Median nerve distal CMAP (mV)	Nerve biopsy
1	CMT2 AR	MFN2 GDAP1	F	6 years	-	+	+	+		42	NA	Denervation-reinnervation
4	CMT2 AD	NEFL	M	9 years	-	+	+	-		NA	NA	-
7	CMT2 AD	GAN	M	68 years	-	+	+	+	Right ptosis and cerebellar ataxia	50	8	-
8	CMT2 AR	MFN2 GDAP1	M	7 years	NA	+	+	-		39	6.7	-
11	CMT2 AR	IGHMBP2	F	7 years	-	+	+	+		39	NA	-
19	CMT2 AR	GAN	M	8 years	NA	+	+	+		-	-	-
22	CMT2 AD	AAPS	M	14 years	NA	NA	+	-		52	7.4	NA
23	CMT2 AR	GDAP1 DCTN1	F	43 years	-	+	+	+		50	NA	-
24	CMT2 AR	DAFS2	M	8 years	-	-	+	+	Surgery of right ureter	42	2.6	NA
26	CMT2 SPO	MFN2	M	3 years	NA	NA	+	NA		NA	NA	Demyelinating aspect with secondary axonal degeneration
27	CMT2 AD	INF2	F	13 years	-	+	+	+		47	1.99	-
29	CMT2 AD	LRSAM1	M	15 years	-	-	+	+	Scoliosis	64	5.5	-
31	CMT2 AD	NEFL	M	Infancy	-	+	+	+	Vocal cords palsy	48	NA	-
33	CMT2 AD	KIF1B	F	30 years	+	+	+	+	Scoliosis, Hashimoto disease, erythema nodosum	60	NA	-
35	CMT2 AD	INF2	F	Infancy	-	-	+	-	Deafness	47	1.43	-
37	CMT2 AD	KIF5A	F	47 years	-	+	+	+		43	NA	-
38	CMT2 SPO	SEPT9 ARHGEF10	M	Adolescence	-	+	+	+	Renal cancer	47	6.5	-
42	CMT2 AD	MFN2	F	9 years	+	+	+	-		62	7.4	-
43	CMT2 AD	BAG3	F	7 years	+	+	+	-	Scoliosis+learning disability+ventricular dilatation+epilepsy	53	8.9	-
44	CMT2 AD	KIF5A	M	Adolescence	-	+	+	-		76	4.5	NA
46	CMT2 AR	MFN2	M	3 years	NA	NA	+	+		48	0.13	-
51	CMT2 AD	HSPB1	M	40 years	+	+	+	+		50	NA	-
53	CMT2 AD	NEFL	F	Infancy	+	+	+	-		58	1.03	NA

Continued

Table 1B Continued

Patient	Disease	Gene	Gender	Age at onset	Deep tendon reflexes	Foot deformities	Muscular weakness and wasting of distal muscles	Sensory loss	Other clinical signs	Median nerve motor MNCV (m/s)	Median nerve distal CMAP (mV)	Nerve biopsy
54	CMT2 AR	GAN	F	2 years	-	+	+	-	Congenital ptosis	NA	NA	-
58	CMT2 AD	BICD2	F	8 years	+	+	+	-		48	11.4	-
59	CMT2 SPO	SPTLC1	F	36 years	-	-	+	+	Chronic inflammatory demyelinating polyneuropathy with 140 antineurofascin Ab+membranous glomerulonephritis	41	5	Demyelinating aspect and axonal degeneration
60	CMT2 AD	AARS	M	20 years	NA	+	+	+		NA	NA	NA

+, presence; -, absence; AD, autosomal dominant; AR, autosomal recessive; CMAP, compound muscle action potential; CMT2, Charcot-Marie-Tooth disease type 2; F, female; IPN, inherited peripheral neuropathy; M, male; MNCV, motor nerve conduction velocity; NA, not available; SPO, sporadic.

new variants, we were able to confirm the pathogenicity of 25, based on: (1) their low frequency in the ExAC database, (2) compatible phenotype, (3) segregation in the family study and (4) in silico pathogenicity prediction. However, we were unable to establish the pathogenicity in 11 patients requiring further explorations at either clinical or genetic levels (see table 2A,B).

Among the 28 patients affected with demyelinating forms, 16 (57%) had a pathogenic variant, 3 (11%) had a potentially pathogenic variant and 9 (32%) remained negative. The most frequently mutated gene was *SH3TC2* responsible for CMT4C with five variants found, of which two were not reported in the literature. The other mutations were found in the following genes: *NEFL* (2), *FIG4* (2), *KIF1B* (2), *HK1* (2), *INF2* (2), *LITAF/SIMPLE* (2), *DNM2* (1), *LMNA* (1), *DCTN1* (1), *GJB1* (2), *GDAP1* (1), *TRPV4* (1), *AARS* (1) and *LRSAM1* (1).

Among the 64 patients with axonal forms, 23 (36%) had a pathogenic variant, 4 (6%) had a potentially pathogenic variant and 37 (58%) remained negative. The most frequently mutated gene was *MFN2* with six variants found, of which two were not reported, and then the following genes: *NEFL* (3), *GDAP1* (3), *AARS* (2), *IGHMBP2* (2), *DCTN1* (1), *DARS2* (2), *KIF5A* (2), *INF2* (2), *LRSAM1* (1), *KIF1B* (1), *SETP9* (1), *ARHGEF10* (1), *HSPB1* (1), *SPTLC1* (1), *BAG3* (1) and *BICD2* (1).

Among the 11 patients who had dHMN, 5 (45%) had a pathogenic variant and 6 (55%) remained negative. The most frequently mutated gene was *SPG11* with two variants found not described in the literature, and then *DYNC1H1* (1), *TRPV4* (1), *REEP1* (1) and *MYH14* (1).

Among the nine patients who had HSAN, three (33%) had a pathogenic variant, two (22%) had a potentially pathogenic variant and four (45%) remained negative. *SPTLC2* was the most frequently mutated gene with two variants found, of which one was not reported. Only one variant was reported in the following genes: *FAM134B*, *HSPB1* and *NEFL*.

Among the eight patients who had intermediate CMT, two (25%) had a pathogenic variant, two (25%) had a potentially pathogenic variant and four (50%) remained negative. A potentially pathogenic variant was found in genes: *DNM2*, *KIF5A*, *YARS* and *INF2*.

Overall, the most frequently mutated genes were (by decreasing order) (see figure 1): *MFN2* (7.7%), *SH3TC2* (6.4%), *NEFL* (5.1%), *GDAP1* (5.1%) and *GAN* (5.1%). Then come the following genes: *AARS* (3.8%), *KIF5A* (3.8%), *KIF1B* (2.6%), *INF2* (2.6%), *DARS2* (2.6%), *DCTN1* (2.6%), *DNM2* (2.6%), *FIG4* (2.6%), *HK1* (2.6%), *IGHMBP2* (2.6%), *LITAF* (2.6%), *SPG11* (2.6%), *SPTLC2* (2.6%), *MYH14* (2.6%), *GJB1* (1.3%), *HSPB1* (1.3%), *TRPV4* (1.3%), *LRSAM1* (1.3%), *BICD2* (1.3%), *SPTLC1* (1.3%), *REEP1* (1.3%), *HSPB3* (1.3%), *FAM134B* (1.3%), *DYNC1H1* (1.3%) and *BAG3* (1.3%).

In our laboratory, before the advent of targeted NGS, 20 genes responsible for IPN were explored by Sanger

**Table 1C** Clinical and electrophysiological features from 14 IPN index patients with hereditary sensory and autonomic neuropathy (HSAN), distal hereditary motor neuropathy (dHMN) and intermediate CMT, for whom a candidate variant was identified

Patient	Disease	Gene	Gender	Age at onset	Deep tendon reflexes	Foot deformities	Muscular weakness and wasting of distal muscles	Sensory loss	Other clinical signs	Median nerve		Nerve biopsy
										motor MNCV (m/s)	distal CMAP (mV)	
6	dHMN AD	TRPV4	M	2 years	+	+	+	-	Scoliosis	Normal	NA	NA
30	dHMN AR	SPG11	M	19 years	-	NA	+	-		NA	NA	NA
39	dHMN AD	REEP1	F	33 years	+	+	+	-	Dysphagia	NA	NA	NA
55	dHMN SPO	DYNC1H1	F	12 years	NA	-	+	-		NA	NA	NA
57	dHMN SPO	MYH14	M	50 years	-	+	+	-	Scoliosis	67	NA	NA
13	HSAN AD	SPTLC2 HSPB3	M	24 years	-	+	+	+		NA	NA	NA
25	HSAN AD	NEFL	F	20 years	-	-	-	+		46	9.79	38
48	HSAN AR	FAM134B	F	12 years	-	+	+	+	Ulceromutilating neuropathy	NA	NA	NA
49	HSAN AD	HSPB1	M	49 years	NA	-	-	+	Sensorineural hearing loss	49	9.2	31
56	HSAN AD	SPTLC2 MYH14	M	10 years	NA	-	-	+	Neurological pain	NA	NA	NA
36	CMT inter AD	DNM2	M	41 years	-	+	NA	NA		54	3.6	NA
40	CMT inter AD	KIF5A	F	53 years	+	+	+	-		48,9	NA	NA
50	CMT inter AD	YARS	M	14 years	+	+	+	+		34	2.2	NA
52	CMT inter AD	INF2	F	NR	NA	NA	NA	NA		NA	NA	NA

+, presence; -, absence; AD, autosomal dominant; AR, autosomal recessive; CMAP, compound muscle action potential; CMT, Charcot-Marie-Tooth disease; F, female; M, male; MNCV, motor nerve conduction velocity; NA, not available; SPO, sporadic.

**Table 2A** Known mutations identified in 81 IPN genes by next-generation sequencing

Patient	Disease classification	Inheritance	Gene	Haplotype	cDNA change	Amino acid change	ACMG classification	Report
2	CMT1	AR	SH3TC2	comp het	c.[211C>T];[2860C>T]	p.[Gln71*];[Arg954*]	P/P	Senderek et al <sup>25</sup>
3	CMT1	SPO	GJB1	het	c.223C>T	p.Arg75Trp	P	Silander et al <sup>26</sup>
6	HMN	AD	TRPV4	het	c.806G>A	p.Arg269His	P	Landouré et al <sup>27</sup>
8	CMT2	AR	MFN2 GDAP1	het	c.707C>T;c.473C>T	p.Thr236Met p.Thr158Ile	PP PP	Kijima et al <sup>28</sup>
9	CMT1	AD	DNM2	het	c.1597G>A	p.Gly533Ser	PP	Fabrizi et al <sup>29</sup>
10	CMT1	AD	NEFL	het	c.1319 C>T	p.Pro440Leu	PP	Benedetti et al <sup>30</sup>
11	CMT2	AR	IGHMBP2	comp het	c.[2368C>T];[2911_2912delAG]	p.[Arg790*];[Arg971Glufs4*]	P/P	Maystact et al <sup>31</sup>
15	CMT1	AD	LITAF	het	c.334G>A	p.Gly112Ser	P	Streeter et al <sup>32</sup>
17	CMT1	AR	GDAP1	hoz	c.786_786delG	p.Phe263Leufs*22	P	Nellis et al <sup>33</sup>
18	CMT1	AD	AARS	het	c.986G>A	p.Arg329His	P	Latour et al <sup>34</sup>
20	CMT1	AR	FIG4	comp het	c.[122T>C];[830_837delGTAAATTT]	p.[Ile41 Thr];[Lys278Trpfs*6]	PP/P	Chow et al <sup>35</sup>
23	CMT2	AR	GDAP1 DCTN1	het	c.358C>T;c.2384G>A	p.Arg120Trp p.Arg795His	P PP	Pedrola et al, <sup>36</sup> Sivera et al <sup>37</sup>
28	CMT1	AD	LITAF DCTN1	het	c.334G>A;c.875G>A	p.Gly112Ser p.Arg292His	PP VUS	Street et al <sup>32</sup>
31	CMT2	AD	NEFL	het	c.803T>G	p.Leu268Arg	PP	Fabrizi et al <sup>38</sup>
32	CMT1	SPO	NEFL	het	c.293A>G	p.Asn98Ser	PP	Yoshihara et al <sup>39</sup>
41	CMT1	AR	SH3TC2	hoz	c.3325 C>T	p.Arg1109*	P	Gooding et al <sup>40</sup>
42	CMT2	AD	MFN2	het	c.311G>T	p.Arg104Leu	P	Sitarz et al <sup>41</sup>
43	CMT2	AD	BAG3	het	c.343C>T	p.Pro115Ser	PP	Villard et al, <sup>42</sup> Selcen et al <sup>43</sup>
47	CMT1	AR	HK1	comp het	c.(1--22124G>C);[c.1--20809 G>A]		PP/PP	Hantke et al <sup>44</sup>
51	CMT2	AD	HSPB1	het	c.523C>T	p.Gln175*	P	Rosser et al <sup>45</sup>
53	CMT2	AD	NEFL	het	c.998T>C	p.Leu333Pro	PP	Choi et al <sup>46</sup>
54	CMT2	AR	GAN	comp het	c.[1429C>T];[1724T>C]	p.[Arg477*];[Ile575Thr]	P/VUS	Bomont et al <sup>47</sup>
56	HSAN	AD	SPTLC2 MYH14	het	c.547 C>T;c.71C>G	p.Arg183Trp p.Ala24Gly	PP VUS	Suriyanarayanan et al <sup>48</sup>
60	CMT2	AD	AARS	het	c.986G>A	p.Arg329His	P	Latour et al <sup>34</sup>

AD, autosomal dominant; AR, autosomal recessive; CMT1, Charcot-Marie-Tooth disease type 1; CMT2, Charcot-Marie-Tooth disease type 2; comp het, compound heterozygous; het, heterozygous; HMN, hereditary motor neuropathy; hoz, homozygous; HSAN, hereditary sensory and autonomic neuropathy; IPN, inherited peripheral neuropathy; P, pathogenic; PP, probably pathogenic; SPO, sporadic; VUS, variant of unknown significance.



**Table 2B** Novel pathogenic sequence variations identified in IPN genes by next-generation sequencing


Patient	Disease classification	Inheritance	Gene	Haplotype	cDNA change	Amino acid change	ACMG	ExAC	Mutation Taster	Polyphen-2	UMD-Predictor	PhyloP
1	CMT2	AR	<i>MFN2</i>	GDAP1	het	c.311G>A c.400delG	PP P	NA NA	D1 NA	1 PD NA	100 P NA	5,37 HC NA
5	CMT1	SPO	<i>INF2</i>		het	c.311G>A	PP	NA	D1	NA	NA	5,69 HC
7	CMT2	AD	<i>GAN</i>		het	c.862C>G	PP	NA	D1	0,001 B	NA	2,95 MC
13	HSAN	AD	<i>SPTLC2</i> ( <i>HSPB3</i> )		het	c.1304G>T (c.246_246delT)	PP VUS	NA 5*10-5	D1 NA	1 PD NA	78 P NA	6,18 HC NA
14	CMT1	AR	<i>SH3TC2</i>	comp.het	c.[224G>C]; [3377T>C]	p.[Arg75Pro]; [Leu1126Pro]	PP/PP	NA/NA	D1/D1	1 PD/0999PD	81 P/ 84 P	3,43 MC 4,81 HC
19	CMT2	AR	<i>GAN</i>	hoz	c.890C>T	p.Pro297Leu	PP	NA	D1	0,906 Possibdam	NA	5,86 HC
22	CMT2	AD	<i>AARS</i>	het	c.525C>G	p.Phe175Leu	PP	1/121242	D1	1 PD	96 P	2,22 MC
24	CMT2	AR	<i>DARS2</i>	comp.het	c.[74_74delT(;)713C>T]	p.[Ile25fs*38(]; Ser238Phe]	P/VUS	2/ 121358/NA	D1/NA	0,959 PD/NA	93 P/ NA	0,85 WC NA
26	CMT2	SPO	<i>MFN2</i>	het	c.1127T>G	p.Met376Arg	PP	NA	D1	0,984 PD	96 P	4,48 HC
29	CMT2	AD	<i>LFSAM1</i>	het	c.2138_2139delT	p.Ile713Thrfs*22	P	NA	NA	NA	NA	NA
30	HMN	AR	<i>SPG11</i>	comp.het	c.[31G>C]; [6167A>T]	p.[Ala11Pro]; [Glu2056Val]	PP/VUS	6/121316 5/57750	D1/Poly	0,808 Possibdam	84 P/ 44 Poly	2,14 MC 0,04 WC
33	CMT2	AD	<i>KIF1B</i>	het	c.2086C>G	p.His696Asp	PP	NA	D1	0,978 PD	81 P	5,61 HC
34	CMT	SPO	<i>KIF1B</i>	het	c.2030G>A	p.Ser677Asn	PP	NA	D1	1 PD	78 P	1,044 WC
36	CMT inter	AD	<i>DNM2</i>	het	c.1352G>T	p.Arg451Leu	PP	NA	D1	NA	93 P	3,68 MC
37	CMT2	AD	<i>KIF5A</i>	het	c.395A>G	p.Lys132Arg	PP	NA	D1	1 PD	78 P	1,088 WC
39	HMN	AD	<i>REEP1</i>	het	c.568G>A	p.Gly190Ser	PP	3/119252	0,999 Poly	0,148 B	63ProbPoly	0,61 WC
40	CMT inter	AD	<i>KIF5A</i>	het	c.854C>T	p.Thr285Ile	PP	NA	D1	1 PD	93 P	5,29 HC
44	CMT2	AD	<i>KIF5A</i>	het	c.332G>C	p.Arg111Pro	PP	NA	D1	1 PD	96 P	5,61 HC
45	CMT1	AD	<i>INF2</i>	het	c.314 T>G	p.Val105Gly	PP	NA	D1	1 PD	90 P	3,11 MC
46	CMT2	AR	<i>MFN2</i>	comp.het	c.[1714C>T]; [1928T>C]	p.[Gln572Glu]; [Leu643Pro]	PP/PP	1/246228 1/246238	D1/D1	0,073 B/ 0,983 PD	100 P 100 P	5,13 HC 4,08 MC
48	HSAN	AR	<i>FAM134B</i>	hoz	c.896-897delAA	p.Lys299Argfs*6	P	1/1 20334	NA	NA	NA	NA
55	HMN	SPO	<i>DYNC1H1</i>	het	c.5578C>A	p.Gln1860Lys	PP	NA	D1	0,001 B	75 P	6,34 HC
57	HMN	SPO	<i>MYH14</i>	het	c.4517G>T	p.Arg1506Leu	PP	NA	0,929 D	0,987 PD	81 P	2,95 MC
58	CMT2	AD	<i>BICD2</i>	het	c.2042C>T	p.Ser681Leu	PP	NA	D1	1 PD	90 P	5,61 HC
59	CMT2	SPO	<i>SPTLC1</i>	het	c.451C>T	p.Arg151Cys	PP	2/118640	D1	0,264 B	100 P	4,00 MC
4	CMT2	AD	<i>NEFL</i>	het	c.67C>G	p.Arg23Gly	VUS	NA	NA	NA	NA	2,09 WC
12	CMT1	AD	<i>LMNA</i>	het	c.1694A>T	p.His565Leu	PP	NA	NA	NA	84 P	2,30 MC
16	CMT1	AD	<i>GJB1</i>	het	c.830G>A	p.Ser277Asn	PB	2/26732	0,876 Poly	0,002 B	63 ProbPoly	2,38 MC

Continued

**Table 2B** Continued

Patient	Disease classification	Inheritance	Gene	Haplotype	cDNA change	Amino acid change	ACMG	ExAC	Mutation Taster	Polyphen-2	UMD-Predictor	PhyloP
21	CMT1	AD	TRPV4 KIF1B LRSAM1	het	c.812G>A c.3350A>G c.1658A>C	p.Arg271His p.Tyr117Cys p.Gln553Pro	VUS VUS VUS	1/120838 1/120270	D1 Poly D1	0,764 Possibdam 0 B 0,994 PD	78 P NA 93 P	5,13 HC - 1,57 NC 3,60 MC
25	HSAN	AD	NEFL	het	c.995_997delAAGC	p.Gln332del	P	NA	NA	NA	NA	NA
27	CMT2	AD	INF2	het	c.2709G>C	p.Gln903His	VUS	1/119080	D 0,991	0,998 PD	57 ProbPoly	2,47 MC
35	CMT2	AD	INF2	het	c.3637C>T	p.Arg1213Trp	VUS	2/61282	Poly	0,998 PD	72 ProbPatho	1,66 WC
38	CMT2	SPO	SEPT9 ARHGEF10	het	c.53G>A.c.3289A>G	p.Arg18Thr p.Ile1097Val	VUS VUS	1/118806 1/120124	Poly Poly	0,015 B 0,023 B	69 ProbPatho 66 ProbPatho	0,28 NC 0,04 WC
49	HSAN	AD	HSPB1	het	c.224G>A	p.Arg75His	VUS	NA	D1	0,539 Possibdam	54 ProbPoly	2,38 MC
50	CMT inter	AD	YARS	het	c.710G>A	p.Arg237Gln	VUS	3/121190	0,983 D	0,018 B	78 P	2,47 MC
52	CMT inter	AD	INF2	het	c.2459G>A	p.Arg820Gln	VUS	NA	Poly	0,001 B	39 Poly	-0,68 NC
61	CMT1	SPO	FGD4		Duplication of exons 3 to 14	Not described in conrad.hg19						
62	CMT2	AR	TFG		Duplication of exons 2 and 3	Not described in conrad.hg19						
28	CMT2	AD	TFG		Sequence variation in LITAF and DCTN1 +duplication of exons 2 and 3	Not described in conrad.hg19						

AD, autosomal dominant; AR, autosomal recessive; B, benign; CMT1, Charcot-Marie-Tooth disease type 1; CMT2, Charcot-Marie-Tooth disease type 2; comp het, compound heterozygous; comp het, composite heterozygous; D1, disease causing; HC, highly conserved; het, heterozygous; hoz, homozygous; HSAN, hereditary sensory and autonomic neuropathy; IPN, inherited peripheral neuropathy; MC, moderately conserved; NA, not available; NT, not conserved; P, pathogenic; Poly, polymorphism; Possibdam, possibly damaging; PB, probably benign; PD, probably damaging; PP, probably pathogenic; ProbPoly, probably polymorphism; ProbPatho, probably pathogenic; SPO, sporadic; VUS, variant of unknown significance; WC, weakly conserved.



**Figure 1** Frequency of mutations in each gene identified in our cohort of 123 patients.

sequencing : *MPZ*, *PMP22*, *PRX*, *EGR2*, *MTMR2*, *NDRG1*, *HSPB1*, *HSPB8*, *BSCL2*, *FGD4*, *LMNA*, *LITAF*, *TRPV4*, *GJB1*, *SH3TC2*, *HK1*, *GDAP1*, *SPTLC1*, *SPTLC2* and *MFN2*. The duplication of *PMP22* responsible for CMT1A was tested by MLPA.

Between 2012 and 2014, 102 patients with IPN were analysed through Sanger sequencing and MLPA. Ninety patients had CMT1 and 12 patients had CMT2. 46/102 (45%) patients had a duplication of *PMP22*. The remaining 56 patients were analysed by Sanger sequencing of candidate genes. A mutation was found in 13 patients (23% of positive cases): four mutations in *MPZ*, four mutations in *GJB1*, two mutations in *GDAP1*, two mutations in *MFN2* and one mutation in *HK1*. Out of the 44 patients with CMT1, we achieved the molecular diagnosis in 25% of them, while only 2 out of 12 patients with CMT2 (17%) could be ascertained. Among the remaining 43 negative cases, 17 were analysed by targeted analysis of 81 genes from NGS data and for 10 of these 17 patients, we identified the molecular defect in a gene unexplored by Sanger sequencing.

Between 2015 and 2017, a pathogenic variant was found in 49 patients (40%) by targeted analysis of 81 genes from NGS data. During the same period, the *PMP22* duplication was identified by MLPA in 67 patients.

In conclusion, in our laboratory, we were able to reach a diagnosis in 23% of patients using Sanger sequencing and in 40% of patients when using targeted NGS of 81 IPN genes. When including *PMP22* duplication, 63% of patients with CMT1 were able to get a diagnosis with Sanger analysis against 87% with NGS. In parallel, we were able to make a diagnosis for 17% of CMT2 patients with Sanger analysis against 36% with NGS.

Our comparative statistical analysis allows us to conclude that the molecular diagnosis yield by targeted

NGS was two times higher than molecular diagnosis by Sanger sequencing (OR=2.18,  $p=0.04$ , (1.0224; 4.8954)).

We compared the solve rate between sporadic cases and family cases. Within our cohort of 123 patients, 42 were sporadic cases; we found a potentially pathogenic variant in 8 of them (19%). Conversely, 51 were familial cases following an autosomal inheritance mode, 25 patients (49%) were able to have a molecular diagnosis.

One of the other objectives of this study was to search for CNVs in our cohort. We found one triplication and one duplication in *FGD4* and *TFG* genes in two patients with CMT1 and CMT2, respectively, as well as a duplication in one of the 81 genes in patient 28 for whom we also found two heterozygous variants in *LITAF* and *DCTN1* using targeted NGS (see table 2B). Overall, ExomeDepth and Q-PCR enabled us to make a potential molecular diagnosis for two additional patients and consequently raise our percentage of diagnosis to 41%.

## DISCUSSION

Before targeted NGS, a strategy of sequential molecular diagnosis through Sanger sequencing was implemented. Genes to be screened were chosen according to phenotype, inheritance and electrophysiological criteria. Therefore, the strategy relied mainly on genotype–phenotype correlation. Targeted NGS allows for a more comprehensive analysis with broader panels of genes, faster and cost-effective outcomes.<sup>13 14</sup>

In our laboratory, we chose targeted NGS in order to avoid the interpretation of numerous variants generated with other NGS strategy, such as whole exome sequencing (WES) or whole genome sequencing, in particular when only one individual is available in the same family.

The use of NGS allowed us to raise our rate of molecular diagnosis to 87% for CMT1 and 36% for CMT2. The high success rate obtained for CMT1 with NGS is due to the high prevalence of *PMP22* duplication in this disease subgroup. The *PMP22* duplication is responsible for CMT1A, the most frequent CMT subtype, accounting for 48.8% to 63.2% of all subtypes.<sup>15–17</sup> Several publications have shown that mutations in *GJB1*, *MFN2*, *MPZ* and *PMP22* account for 80 to 94.9% of CMT, and recommended to first complete targeted Sanger sequencing based on clinical phenotype. *MFN2* was the most frequently mutated gene found among our cohort of 64 patients with CMT2. NGS enabled us to find a high frequency of pathogenic variants in *NEFL*, *GAN*, *AARS* and *KIF5A*. These were less frequent and we would not have explored them as first line. Other variants found in *BAG3*, *BICD2*, *DYNC1H1*, *REEP1* and *FAM134B* are much more rare. A survey of 17 880 patients with CMT compared the diagnosis outcomes of Sanger-MLPA and NGS-MLPA<sup>17</sup> analyses. This study suggested that the frequency of positive results for 14 CMT genes was not significantly different ( $p<0.05$ ) in spite of differences in testing strategies between Sanger and NGS. But the bias of this survey is that it compared the same genes in the two groups.

In our cohort of 123 patients, NGS enabled us to make a positive molecular diagnosis of point mutation (single-nucleotide variation (SNV)) for 49 patients (40%) while saving considerable time and cost. This result is consistent with those found in previous studies, including Hartley *et al.*,<sup>18</sup> who were able to make a molecular diagnosis by WES in 12 patients (24%) in a cohort of 50 families with IPN. Similarly, Gonzaga-Jauregui *et al.*<sup>4</sup> was able to make a molecular diagnosis in 17 patients (46%) by WES in a cohort of 37 families with CMT.

Recently, Dorhn and collaborators<sup>19</sup> describe a cohort of 612 subjects who came from Germany affected by IPN and found a majority of point mutation in *MPZ*, *MFN2*, *GJB1* and *SH3TC2*. Our study describes patients who came from the south of France and found a majority of mutation in *MFN2*, *SH3TC2*, *NEFL*, *GAN*, *GDAP1*, *AARS* and *KIF5A*. In fact, our cohort is composed of people who came from the Mediterranean region which probably explains the different spectrum delineation of our study, notably the relatively high frequency of *GAN* variants.

Moreover, 11 patients in our cohort were undiagnosed but we found a potentially pathogenic variant without confirmation of pathogenicity. In the following, we will detail some cases of particular interest.

Boyer and collaborators<sup>20</sup> have described nine mutations in *INF2* in 12 patients with CMT with focal segmental glomerulosclerosis. In our study, patients 5, 27, 35, 45 and 52 have a sequence variation in this gene and, to date, no nephrological disease is known in these patients. Patients 5 and 45 have a probably pathogenic variant according to Richards classification. Those mutations are located in the same codon that patient described by Boyer. While patient 27, 35 and 52 have a VUS according to Richards classification. At that time, without nephrological examination and familial study, we were not able to exclude those variants.

We have two patients in this cohort affected by CMT1 (patient 21) and CMT2 (patient 38) carrying numerous variants in potential genes. We found three VUS in *TRPV4* (CMT2C), *LRSAM1* (CMT2P) and *KIF1B* (CMT2A1) genes for patient 21. Mutations in these three genes were reported to be in relation with CMT type 2 or dHMN. In that case, we have too much variant which could explain patient's phenotype at isolated state or in association. A familial study is indicated. Patient 38 showed two VUS in *SEPT9* (hereditary neuralgic amyotrophy)<sup>21</sup> and *ARHGEF10* (slowed nerve conduction velocity).<sup>22</sup> These mutations could thus separately or in association explain these patient's phenotype. Unfortunately, the family study could not be achieved and we could not make any conclusions related to the contribution of these variants in the disease.

Only one variant was classified as probably benign in Richards classification, and namely variant c.830G>A in *GJB1* which was identified in patient 16 suffering from autosomal-dominant CMT1 with first clinical manifestations at the age of 15. She presents a severe phenotype with standard clinical signs of peripheral motor and

sensory neuropathy. Motor nerve conduction velocity was 36 m/s in the electromyogram and the nerve biopsy showed Schwann cell proliferation in the form of an onion bulb. The segregation analysis found this variant in the asymptomatic mother. Her mother could have an inactivation of the mutant allele on one of the X-chromosomes, thus only expresses wild-type Cx32 from the normal allele. NGS analysis did not enable us to identify another potential variant responsible for her neuropathy. Patient 4, affected with autosomal-dominant CMT2, showed severe clinical signs at the age of 9. We found a VUS c.67C>G, (p.Arg23Gly) in *NEFL* (CMT1F, CMT2E). The segregation analysis found this one in the asymptomatic mother. This case can maybe illustrate incomplete penetrance for this mutation in this family. Alternatively, the disease, in this patient, is due to another mutation in a gene not explored in this targeted NGS.

Patients 13 and 56 are affected with HSN and carry a mutation in *SPTLC2*, known to be responsible for HSN1C (MIM 613640). They also have a VUS (according to Richards classification) in another gene. One in *HSPB3* (HMN2C, MIM 613376) for patient 13 and the second one in *MYH14* (peripheral neuropathy, myopathy, hoarseness and hearing loss, MIM 614369) for patient 56. These two patients have perforating ulcers of the foot, hypoaesthesia at sock level but their motor picture also included a severe progressive motor deficiency, severe wasting, contractures and neuropathic pain. In these cases, *SPTLC2* could alone be responsible for the patient's phenotype and we can consider *HSPB3* and *MYH14* as modifiers or modulators factors likely to account for the motor phenotype of these two patients. In comparison, Sinkiewicz-Darol's team<sup>23</sup> showed that the presence of a variant p.Ile92Val in gene *LITAF/SIMPLE* of patients who presented with a duplication or deletion of *PMP22* was linked with an earlier onset of CMT1A or Hereditary Neuropathy by Hypersensitivity to Pressure Palsie (HNPP) and could be considered as a modifier. These additional variants could contribute to the variability of the clinical phenotype expression.

Even if the majority of IPNs are explained by SNV, CNV may equally be a genetic cause of IPN and, thanks to NGS strategy, we can now detect these two types of variations. We found three CNVs in three patients, that is, 2.4% in our cohort. In comparison, Pfundt's team<sup>24</sup> looked for CNVs with the coNIFER software using reading depth data with exome high-throughput sequencing in a cohort of 2603 patients who had different genetic pathologies. In the group with neurodevelopment disorders, muscular disorders and abnormal coordination, they respectively found 1.3%, 0.6% and 0.9% of CNV. We found a higher percentage of CNVs in our IPN group (none significant Fisher's exact test). This study of CNV enabled us to raise our rate of molecular diagnosis to 51 patients (41%) out of a cohort of 123 patients.

Targeted NGS allows us to improve our molecular diagnosis in IPN and allows an accurate genetic counselling in



families. Moreover, a positive molecular diagnosis is the first step to participate in clinical trials.

## CONCLUSION

In our laboratory, NGS improved the molecular diagnostic rate, allowing for 40% of the patients suffering from IPN. Eighty-seven per cent of patients with CMT1 can now get a precise molecular diagnosis. On the other hand, 64% of CMT2 cases remain unsolved. Indeed, numerous aetiologies of neuropathies are found in elderly patients, such as diabetic, inflammatory, alcoholic, idiopathic and autoimmune neuropathies and may be confused with hereditary neuropathies. Most importantly, NGS analysis allows to describe novel mutations, not yet reported in the literature, that are of significant importance for the scientific and medical community. Moreover, the analysis of NGS data enabled us to detect possible duplications and deletions of genes not investigated routinely. Targeted NGS with a panel of 81 genes is therefore well adapted to IPN molecular diagnosis. However, the generation of many variants of unknown significance requires a collegial interpretation by biologists, geneticists and neurologists.

## Author affiliations

<sup>1</sup>Département de génétique médicale, Hôpital Timone enfants, Assistance Publique Hôpitaux de Marseille, Marseille, France

<sup>2</sup>Centre de référence des maladies neuromusculaires, Hôpital Pitié-Salpêtrière, Assistance-Publique Hôpitaux de Paris, Paris, France

<sup>3</sup>Centre de référence des maladies neuromusculaires, Hôpital Timone enfants, Assistance Publique Hôpitaux de Marseille, Marseille, France

<sup>4</sup>Centre de référence des maladies neuromusculaires, Hôpital Timone Adultes, Assistance Publique Hôpitaux de Marseille, Marseille, France

<sup>5</sup>INSERM, MMG, UMR 1251, Aix Marseille Univ, Marseille, France

<sup>6</sup>Département de génétique médicale, Hôpital Archet 2, CHU de Nice, Nice, France

**Acknowledgements** The authors would like to thank the patients and their families for their kind cooperation. They also thank Caroline Lacoste and Christophe Pécheux for supplying the NGS platform and Catherine Badens for helping.

**Contributors** JB, NB-P, VD and NL contributed to the design of the study. NB-P contributed to the supervision and mentorship. JB, AB, J-PD and NM participated in data analysis and interpretation. JB, TS and NB-P contributed to manuscript drafting. TS, FA, BC, ES-C, ED, AV, SA, AC and NL contributed to the collection of clinical data during consultations. All authors participated in critical revisions of the manuscript and can take responsibility for its integrity and the accuracy of the data analysis.

**Funding** The authors have not declared a specific grant for this research from any funding agency in the public, commercial or not-for-profit sectors.

**Competing interests** None declared.

**Patient consent** obtained.

**Ethics approval** Aix-Marseille University Ethics Committee.

**Provenance and peer review** Not commissioned; externally peer reviewed.

**Data sharing statement** The data are not available freely. Enquiries and requests for further information should be made to corresponding author.

**Open access** This is an open access article distributed in accordance with the Creative Commons Attribution Non Commercial (CC BY-NC 4.0) license, which permits others to distribute, remix, adapt, build upon this work non-commercially, and license their derivative works on different terms, provided the original work is properly cited, appropriate credit is given, any changes made indicated, and the use is non-commercial. See: <http://creativecommons.org/licenses/by-nc/4.0/>.

## REFERENCES

- Braathen GJ. Genetic epidemiology of Charcot-Marie-Tooth disease. *Acta Neurol Scand* 2012;126:iv–22.
- Berciano J, Garcia A, Gallardo E, *et al*. Intermediate Charcot-Marie-Tooth disease: an electrophysiological reappraisal and systematic review. *J Neurol* 2017;264:1655–77.
- Timmerman V, Strickland AV, Züchner S, *et al*. Genetics of Charcot-Marie-Tooth (CMT) Disease within the Frame of the Human Genome Project Success. *Genes* 2014;5:13–32.
- Gonzaga-Jauregui C, Harel T, Gambin T, *et al*. Exome sequence analysis suggests that genetic burden contributes to phenotypic variability and complex neuropathy. *Cell Rep* 2015;12:1169–83.
- Desvignes JP, Bartoli M, Delague V, *et al*. VarAFT: a variant annotation and filtration system for human next generation sequencing data. *Nucleic Acids Res*. In Press. 2018;46:W545–W553.
- Schwarz JM, Rödelsparger C, Schuelke M, *et al*. MutationTaster evaluates disease-causing potential of sequence alterations. *Nat Methods* 2010;7:575–6.
- Kumar P, Henikoff S, Ng PC. Predicting the effects of coding non-synonymous variants on protein function using the SIFT algorithm. *Nat Protoc* 2009;4:1073–81.
- Adzhubei IA, Schmidt S, Peshkin L, *et al*. A method and server for predicting damaging missense mutations. *Nat Methods* 2010;7:248–9.
- Salgado D, Desvignes JP, Rai G, *et al*. UMD-Predictor: A High-throughput sequencing compliant system for pathogenicity prediction of any human cDNA substitution. *Hum Mutat* 2016;37:439–46.
- Richards S, Aziz N, Bale S, *et al*. Standards and guidelines for the interpretation of sequence variants: A joint consensus recommendation of the American college of medical genetics and genomics and the association for molecular pathology. *Genet Med* 2015;17:405–23.
- Ellingford JM, Campbell C, Barton S, *et al*. Validation of copy number variation analysis for next-generation sequencing diagnostics. *Eur J Hum Genet* 2017;25:719–24.
- Wieme MH, Monia Ben H, Yosr B, *et al*. Confirmation of the spinal motor neuron gene 2 (SMN2) copy numbers by real-time PCR. *Diagn Mol Pathol* 2012;21:172–5.
- Vissers L, van Nimwegen KJM, Schieving JH, *et al*. A clinical utility study of exome sequencing versus conventional genetic testing in pediatric neurology. *Genet Med* 2017;19:1055–63.
- Monroe GR, Frederix GW, Savelberg SM, *et al*. Effectiveness of whole-exome sequencing and costs of the traditional diagnostic trajectory in children with intellectual disability. *Genet Med* 2016;18:949–56.
- Tazir M, Hamadouche T, Nouioua S, *et al*. Hereditary motor and sensory neuropathies or Charcot-Marie-Tooth diseases: an update. *J Neurol Sci* 2014;347:14–22.
- Saporta AS, Sottile SL, Miller LJ, *et al*. Charcot-Marie-Tooth disease subtypes and genetic testing strategies. *Ann Neurol* 2011;69:22–33.
- DiVincenzo C, Elzinga CD, Medeiros AC, *et al*. The allelic spectrum of Charcot-Marie-Tooth disease in over 17,000 individuals with neuropathy. *Mol Genet Genomic Med* 2014;2:522–9.
- Hartley T, Wagner JD, Warman-Chardon J, *et al*. Whole-exome sequencing is a valuable diagnostic tool for inherited peripheral neuropathies: Outcomes from a cohort of 50 families. *Clin Genet* 2018;93.
- Dohrn MF, Glöckle N, Mulahasanovic L, *et al*. Frequent genes in rare diseases: panel-based next generation sequencing to disclose causal mutations in hereditary neuropathies. *J Neurochem* 2017;143:507–22.
- Boyer O, Nevo F, Plaisier E, *et al*. INF2 mutations in Charcot-Marie-Tooth disease with glomerulopathy. *N Engl J Med* 2011;365:2377–88.
- Kuhlenbäumer G, Hannibal MC, Nelis E, *et al*. Mutations in SEPT9 cause hereditary neuralgic amyotrophy. *Nat Genet* 2005;37:1044–6.
- Verhoeven K, De Jonghe P, Van de Putte T, *et al*. Slowed conduction and thin myelination of peripheral nerves associated with mutant rho Guanine-nucleotide exchange factor 10. *Am J Hum Genet* 2003;73:926–32.
- Sinkiewicz-Darol E, Lacerda AF, Kostera-Pruszczyk A, *et al*. The LITAF/SIMPLE I92V sequence variant results in an earlier age of onset of CMT1A/HNPP diseases. *Neurogenetics* 2015;16:27–32.
- Pfundt R, Del Rosario M, Vissers L, *et al*. Detection of clinically relevant copy-number variants by exome sequencing in a large cohort of genetic disorders. *Genet Med* 2017;19:667–75.
- Senderek J, Bergmann C, Stendel C, *et al*. Mutations in a gene encoding a novel SH3/TPR domain protein cause autosomal recessive Charcot-Marie-Tooth type 4C neuropathy. *Am J Hum Genet* 2003;73:1106–19.



26. Silander K, Meretoja P, Pihko H, *et al.* Screening for connexin 32 mutations in Charcot-Marie-Tooth disease families with possible X-linked inheritance. *Hum Genet* 1997;100:391–7.
27. Landouré G, Zdebik AA, Martinez TL, *et al.* Mutations in TRPV4 cause Charcot-Marie-Tooth disease type 2C. *Nat Genet* 2010;42:170–4.
28. Kijima K, Numakura C, Izumino H, *et al.* Mitochondrial GTPase mitofusin 2 mutation in Charcot-Marie-Tooth neuropathy type 2A. *Hum Genet* 2005;116:23–7.
29. Fabrizi GM, Ferrarini M, Cavallaro T, *et al.* Two novel mutations in dynamin-2 cause axonal Charcot-Marie-Tooth disease. *Neurology* 2007;69:291–5.
30. Benedetti S, Previtali SC, Coviello S, *et al.* Analyzing histopathological features of rare charcot-marie-tooth neuropathies to unravel their pathogenesis. *Arch Neurol* 2010;67:1498–505.
31. Maystadt I, Zarhrate M, Landrieu P, *et al.* Allelic heterogeneity of SMARD1 at the IGHMBP2 locus. *Hum Mutat* 2004;23:525–6.
32. Street VA, Bennett CL, Goldy JD, *et al.* Mutation of a putative protein degradation gene LITAF/SIMPLE in Charcot-Marie-Tooth disease 1C. *Neurology* 2003;60:22–6.
33. Nelis E, Erdem S, Van Den Bergh PY, *et al.* Mutations in GDAP1: autosomal recessive CMT with demyelination and axonopathy. *Neurology* 2002;59:1865–72.
34. Latour P, Thauvin-Robinet C, Baudalet-Méry C, *et al.* A major determinant for binding and aminoacylation of tRNA(Ala) in cytoplasmic Alanyl-tRNA synthetase is mutated in dominant axonal Charcot-Marie-Tooth disease. *Am J Hum Genet* 2010;86:77–82.
35. Chow CY, Zhang Y, Dowling JJ, *et al.* Mutation of FIG4 causes neurodegeneration in the pale tremor mouse and patients with CMT4J. *Nature* 2007;448:68–72.
36. Pedrola L, Espert A, Wu X, *et al.* GDAP1, the protein causing Charcot-Marie-Tooth disease type 4A, is expressed in neurons and is associated with mitochondria. *Hum Mol Genet* 2005;14:1087–94.
37. Sivera R, Espinós C, Vilchez JJ, *et al.* Phenotypical features of the p.R120W mutation in the GDAP1 gene causing autosomal dominant Charcot-Marie-Tooth disease. *J Peripher Nerv Syst* 2010;15:334–44.
38. Fabrizi GM, Cavallaro T, Angiari C, *et al.* Charcot-Marie-Tooth disease type 2E, a disorder of the cytoskeleton. *Brain* 2007;130:394–403.
39. Yoshihara T, Yamamoto M, Hattori N, *et al.* Identification of novel sequence variants in the neurofilament-light gene in a Japanese population: analysis of Charcot-Marie-Tooth disease patients and normal individuals. *J Peripher Nerv Syst* 2002;7:221–4.
40. Gooding R, Colomer J, King R, *et al.* A novel Gypsy founder mutation, p.Arg1109X in the CMT4C gene, causes variable peripheral neuropathy phenotypes. *J Med Genet* 2005;42:e69.
41. Sitarz KS, Yu-Wai-Man P, Pyle A, *et al.* MFN2 mutations cause compensatory mitochondrial DNA proliferation. *Brain* 2012;135:e219–3.
42. Villard E, Perret C, Gary F, *et al.* A genome-wide association study identifies two loci associated with heart failure due to dilated cardiomyopathy. *Eur Heart J* 2011;32:1065–76.
43. Selcen D, Muntoni F, Burton BK, *et al.* Mutation in BAG3 causes severe dominant childhood muscular dystrophy. *Ann Neurol* 2009;65:83–9.
44. Hantke J, Chandler D, King R, *et al.* A mutation in an alternative untranslated exon of hexokinase 1 associated with hereditary motor and sensory neuropathy - Russe (HMSNR). *Eur J Hum Genet* 2009;17:1606–14.
45. Rossor AM, Davidson GL, Blake J, *et al.* A novel p.Gln175X [corrected] premature stop mutation in the C-terminal end of HSP27 is a cause of CMT2. *J Peripher Nerv Syst* 2012;17:201–5.
46. Choi BO, Lee MS, Shin SH, *et al.* Mutational analysis of PMP22, MPZ, GJB1, EGR2 and NEFL in Korean Charcot-Marie-Tooth neuropathy patients. *Hum Mutat* 2004;24:185–6.
47. Bomont P, loos C, Yalcinkaya C, *et al.* Identification of seven novel mutations in the GAN gene. *Hum Mutat* 2003;21:446.
48. Suriyanarayanan S, Auranen M, Toppila J, *et al.* The Variant p.(Arg183Trp) in SPTLC2 Causes Late-Onset Hereditary Sensory Neuropathy. *Neuromolecular Med* 2016;18:81–90.