The use of herbal medicines for induction of labour and onset of labour (by study design)

Study ID		RR (95% CI)	Events, Herbs	Events, No-herbs	% Weight
Experimental Azhari (2006) Pirdadeh (2007) Saberi (2008) Gilad (2018) Subtotal (I-squared = 86.6%, p = 0.000)		12.46 (1.77, 87.73 12.46 (1.77, 87.73 5.83 (3.38, 10.07) 1.21 (0.69, 2.10) 4.72 (1.33, 16.72))13/24 70/100 16/38	1/23 1/23 12/100 15/43 29/189	8.08 8.08 15.10 15.07 46.33
Non-experimental Davis (1984) Garry (2000) Boel (2009) Neri (2017) Subtotal (I-squared = 88.4%, p = 0.000)	+ 	1.28 (1.04, 1.57) 13.85 (3.50, 54.85 0.93 (0.48, 1.81) 6.33 (2.03, 19.72) 2.70 (0.96, 7.63))30/52 8/43	52/89 2/48 68/341 3/40 125/518	16.12 10.83 14.60 12.12 53.67
Overall (I-squared = 90.2%, p = 0.000) <u>NOTE: Weights are from random effects analysis</u> .05 Favours control	1 10 2 Favours treatment	3.46 (1.58, 7.55) 20	249/428	154/707	100.00