Corrections: Maternal seafood intake and the risk of small for gestational age newborns: a case-control study in Spanish women

Amezcu-Prieto C, Martínez-Galiano JM, Salcedo-Bellido I, *et al.* Maternal seafood intake and the risk of small for gestational age newborns: a case-control study in Spanish women *BMJ Open* 2018;8:e020424. doi: 10.1136/bmjopen-2017-020424.

This article was previously published with some errors.

In Table 3 in the quantity of quintiles of seafood intake (in grams), where it says: Q1 (≤ 8); Q2 (8-14); Q3 (14-21.2); Q4 (21.2-29); Q5 (> 29)

It should be:

Q1 (≤ 56); Q2 (56.1-74.0); Q3 (74.1-92.6); Q4 (92.7-121); Q5 (> 121)

Considering this correction, some sections changes from the original published manuscript:

In the Abstract, in the result section, the sentence:

The risk of an SGA newborn was lower among women who consumed > 29 g/day fish compared with women who consumed ≤ 8 g (adjusted OR 0.63, 95% CI 0.41 to 0.98; $p=0.025$ for a trend).

Should be:

The risk of an SGA newborn was lower among women who consumed > 121 g/day fish compared with women who consumed ≤ 56 g (adjusted OR 0.63, 95% CI 0.41 to 0.98; $p=0.025$ for a trend).

In Abstract, in the conclusion, the sentence:

An average seafood intake of at least 29 g/ day during pregnancy, equivalent to 2–3 servings/week, reduced the risk of an SGA newborn, compared with an average seafood intake of less than 8 g/day.

Should be:

An average seafood intake of at least 121 g/day during pregnancy, equivalent to 3–4 servings/week, reduced the risk of an SGA newborn, compared with an average seafood intake of less than 56 g/day.

In the result section of the manuscript, in page 4, the sentence:

The OR of delivering an SGA newborn was lower with seafood intakes of > 29 g/day (Q5; aOR 0.63, 95% CI 0.41 to 0.98) compared with intakes of ≤ 8 g/day (Q1; $p=0.025$ for this trend).

Should be:

The OR of delivering an SGA newborn was lower with seafood intakes of > 121 g/day (Q5; aOR 0.63, 95% CI 0.41 to 0.98) compared with intakes of ≤ 56 g/day (Q1; $p=0.025$ for this trend).

In the conclusion:

In summary, we found that an intake of at least 29 g/day of seafood, or at least 1 g/day of n3-fatty acids, was observed nearly two times as frequently in controls than in women with SGA newborns.

Should be:

In summary, we found that an intake of at least 121 g/day of seafood, or at least 1 g/day of n3-fatty acids, was observed nearly two times as frequently in controls than in women with SGA newborns.

Open access This is an open access article distributed in accordance with the Creative Commons Attribution Non Commercial (CC BY-NC 4.0) license, which permits others to distribute, remix, adapt, build upon this work non-commercially, and license their derivative works on different terms, provided the original work is properly cited, appropriate credit is given, any changes made indicated, and the use is non-commercial. See: <http://creativecommons.org/licenses/by-nc/4.0/>.

© Author(s) (or their employer(s)) 2018. Re-use permitted under CC BY-NC. No commercial re-use. See rights and permissions. Published by BMJ.

BMJ Open 2018;**8**:e020424corr1. doi:10.1136/bmjopen-2017-020424corr1