

S7 Table. Most commonly prescribed anti-infectives by cohort.

Measles patients	N	Percent	Matched controls	N	Percent
Amoxicillin 125mg/5ml oral suspension sugar free	2,246	0.2194	Amoxicillin 125mg/5ml oral suspension sugar free	14,871	0.2184
Amoxicillin 125mg/5ml oral suspension	1,238	0.1209	Amoxicillin 125mg/5ml oral suspension	7,500	0.1101
Chloramphenicol 0.5% eye drops	606	0.0592	Chloramphenicol 0.5% eye drops	4,111	0.0604
Trimethoprim 50mg/5ml oral suspension sugar free	567	0.0554	Phenoxyethylpenicillin 125mg/5ml oral solution	3,527	0.0518
Phenoxyethylpenicillin 125mg/5ml oral solution	458	0.0447	Fusidic acid 2% cream	2,921	0.0429
Erythromycin ethyl succinate 125mg/5ml oral suspension	440	0.0430	Trimethoprim 50mg/5ml oral suspension sugar free	2,791	0.0410
Amoxicillin 250mg/5ml oral suspension sugar free	322	0.0314	Erythromycin ethyl succinate 125mg/5ml oral suspension	2,476	0.0364
Flucloxacillin 125mg/5ml oral solution	235	0.0230	Amoxicillin 250mg/5ml oral suspension sugar free	2,041	0.0300
Co-amoxiclav 125mg/31mg/5ml oral suspension sugar free	234	0.0229	Flucloxacillin 125mg/5ml oral solution	1,874	0.0275
Fusidic acid 2% cream	196	0.0191	Chloramphenicol 1% eye ointment	1,292	0.0190