BMJ Open Prevalence and patterns of multimorbidity in Amazon Region of Brazil and associated determinants: a cross-sectional study

Maria Elizete A Araujo,^{1,2} Marcus T Silva,^{3,4} Tais F Galvao,⁵ Bruno P Nunes,⁶ Mauricio G Pereira⁷

To cite: Araujo MEA, Silva MT, Galvao TF, *et al.* Prevalence and patterns of multimorbidity in Amazon Region of Brazil and associated determinants: a cross-sectional study. *BMJ Open* 2018;**8**:e023398. doi:10.1136/bmjopen-2018-023398

► Prepublication history and additional material for this paper are available online. To view, please visit the journal (<http://dx.doi.org/10.1136/bmjopen-2018-023398>).

Received 4 April 2018

Revised 21 August 2018

Accepted 26 September 2018

ABSTRACT

Objectives To estimate the prevalence of multimorbidity and to identify factors associated with it in the adult population from the metropolitan region of Manaus.

Design Cross-sectional population-based study.

Setting Interviews conducted between May and August of 2015 in eight cities that compose the metropolitan region of Manaus, Amazonas, Brazil.

Participants 4001 adults aged ≥ 18 years.

Primary outcome measures Multimorbidity, measured by the occurrence of ≥ 2 and ≥ 3 chronic diseases, was the primary outcome. The associated factors were investigated by calculating the prevalence ratio (PR) obtained by Poisson regression, with robust adjustment of the variance in a hierarchical model. A factor analysis was conducted to investigate multimorbidity clusters.

Results Half of the interviewees were women. The presence of a chronic disease was reported by 57.2% (95% CI 56.6% to 59.7%) of the interviewees, and the mean morbidity was 1.2 (1.1–1.2); 29.0% (95% CI 27.6% to 30.5%) reported ≥ 2 morbidities and 15.2% (95% CI 14.1% to 16.4%) reported ≥ 3 chronic conditions. Back pain was reported by one-third of the interviewees. Multimorbidity was highest in women, PR=1.66 (95% CI 1.50 to 1.83); the elderly, PR=5.68 (95% CI 4.51 to 7.15) and individuals with worse health perception, PR=3.70 (95% CI 2.73 to 5.00). Associated factors also included undergoing medical consultations, hospitalisation in the last year, suffering from dengue in the last year and seeking the same healthcare service. Factor analysis revealed a pattern of multimorbidity in women. The factor loading the most strength of association in women was heart disease. In men, an association was identified in two groups, and lung disease was the disease with the highest factorial loading.

Conclusion Multimorbidity was frequent in the metropolitan region of Manaus. It occurred most often in women, in the elderly and in those with worse health perception.

BACKGROUND

Multimorbidity is the occurrence of different chronic clinical conditions in an individual, without a single condition being considered

Strengths and limitations of this study

- This is the first study on the prevalence of multimorbidity in adults from the metropolitan region of Manaus, Amazonas, Brazil, using data from a population-based survey.
- We used probabilistic complex sampling in three stages, census track, household and individual, to include 4001 adults living in one of the eight cities of the metropolitan region.
- This research increases knowledge about the epidemiological factors associated with multimorbidity.
- The method used to measure outcomes, self-report, is subject to errors and influenced by memory bias.

the main cause.^{1 2} Multimorbidity is operationally defined as the occurrence of two or more chronic diseases.^{3–5} In recent decades, population ageing, lifestyle changes, improved socioeconomic conditions and increased diagnostic ability of health services have contributed to a significant rise in the population that survives serious diseases, causing an accumulation of health problems in specific population groups. This situation has contributed to the increased prevalence of multimorbidity.^{6–8}

The frequency of multimorbidity varies according to the evaluated diseases, the age of the population, the individual's socioeconomic and demographic level and the individual's health condition. The rising prevalence of multimorbidity has resulted in higher costs of health services.^{9–12} The costs associated with multimorbidity can reach 75% of total health expenditures, which includes physician consultation, hospitalisation, odontological care, medication and rehabilitation.¹³

In Brazil, multimorbidity ranged from 26% to 29% in adults living in the southern—and more developed—region and from 14% to 19% in the northern region.¹⁴ Studies in

© Author(s) (or their employer(s)) 2018. Re-use permitted under CC BY-NC. No commercial re-use. See rights and permissions. Published by BMJ.

For numbered affiliations see end of article.

Correspondence to

Dr Maria Elizete A Araujo; elizetemanaus@gmail.com

specific populations conducted in the south and south-east Brazilian regions identified higher prevalence of multimorbidity in women and the elderly than in other groups.^{15 16} Differences detected suggest heterogeneity due to socioeconomic development.¹⁷ In northern Brazil, there is a lack of studies identifying more susceptible groups and studies that expand our knowledge about multimorbidity at the local level.

To obtain evidence of the health status and usage of health services, a large survey was performed in 2015 in the Manaus metropolitan region,^{18 19} the most populated region and largest economic cluster in northern Brazil. This region comprises >60% of the 3.5 million people of Amazonas, which has the largest land area, the lowest population density and the highest population of indigenous people (4.7%) in Brazil.²⁰ Health coverage is mainly public (Unified Health System), and this region had the lowest coverage of health insurance in the country in 2013 (13.0%).²¹ The present research estimated the prevalence of and factors associated with multimorbidity in the adult population of the metropolitan region of Manaus.

METHODS

Study design

This is a cross-sectional population-based study on the urban population of the metropolitan region of Manaus, consisting of the capital of Amazonas, Manaus and seven surrounding cities. Multimorbidity was considered a primary outcome, which was categorised as ≥ 2 or ≥ 3 chronic diseases. The present analysis is part of a larger study aimed to examine the use of health services and inputs in the region from May to August 2015. Details of the study design and the representativeness of the sample are available elsewhere.¹⁸

Participants and study size

We calculated the sample size as 4000 adults aged ≥ 18 years to be interviewed, who were selected by probabilistic complex sampling—by cluster and stratified by sex and age—in three stages (census track, household and individual).¹⁸ We assume an estimated 50% prevalence of use of health services, considering a CI of 95%, absolute precision of 2% and a design effect of 1.5.²² We added 10% to compensate for possible losses and refusals.

Variables and data collection

The primary outcome was self-reported multimorbidity, defined as two or more affirmative answers to any of the following questions: "Have any doctors ever diagnosed you with [...]?" (1) hypertension, (2) diabetes, (3) high cholesterol, (4) heart disease (heart attack, angina, heart failure or other), (5) stroke, (6) asthma or asthmatic bronchitis, (7) arthritis or rheumatism, (8) depression, (9) pulmonary disease (pulmonary emphysema, chronic bronchitis or chronic obstructive pulmonary disease), (10) cancer or (11) chronic kidney disease and (12) "Do you have any chronic spinal problems, such as chronic

back or neck pain, low back pain, sciatic pain, vertebral or disc problems?" These questions were previously used in the National Health Survey.²³

The independent variables were sex; age (18–24; 25–34; 35–44; 45–59 and ≥ 60 years)²⁴; marital status; self-reported skin colour; education; social class²⁵; occupation; private health insurance (yes, no); self-perception of health status; place of attendance (capital, countryside); seeking the same healthcare service when in need of attendance (health reference; yes, no); physician visit in the last 12 months (yes, no); hospitalisation in the last year (yes, no); malaria in the last 12 months (yes, no); dengue in the last 12 months (yes, no) and types of services one usually seeks when in need of medical care (primary, secondary or tertiary).

Interviewers with experience in conducting home interviews collected the data on a mobile electronic device (Samsung Galaxy Tab3 SM-T110). Interview records were transmitted over the internet and stored using Survey To Go software (Dooblo, Israel).

Statistical analyses

Statistical analyses were carried out in Stata V.14.2. In all calculations, the complex sampling design was weighted by incorporating sample weights (*svy* command).

Descriptive statistics were initially obtained through prevalence calculation. The respective CIs and P values of difference were calculated by Pearson's X^2 between sociodemographic characteristics and multimorbidity. The prevalence of the most common diseases stratified by sex, age group and multimorbidity was also calculated. At this stage, morbidities with a prevalence of $< 5\%$ were excluded.

Bivariate analyses were performed between all independent and dependent variables to calculate prevalence ratios (PRs) and 95% CIs. To identify the factors associated with multimorbidity, PRs were adjusted using Poisson regression with robust variance adjustment.^{26–28}

A hierarchical model consisting of three blocks was constructed of the most distal to the most proximal determinants of multimorbidity: (1) demographic variables (sex, age, race, marital status); (2) socioeconomic variables (economic, education classification, occupation) and (3) health variables (private health insurance, health status, demand for the same health service, physician visit, hospitalisation, dengue, malaria and type of service usually used). The variables from the first block were retained for the next stage if they presented a p value ≤ 0.05 . Multicollinearity among the independent variables was discarded by assessing the variance inflation factors.²⁹

Exploratory factor analysis stratified by sex was performed to identify multimorbidity patterns, that is, to identify associations, selecting variables with potentially common causal factors, such as interaction between diseases and/or common risk factors.^{30 31} The tetrachoric correlation coefficient was used in the analysis because it is better than Pearson's correlation coefficient for dichotomous outcomes.³² The suitability of the sampling was

evaluated by the Kaiser-Meyer-Olkin test (KMO), which was considered adequate if the index was ≥ 0.70 , and the Bartlett sphericity test, which was considered adequate if its p value was ≤ 0.05 .^{30 33} To establish the number of factors to be maintained, Cattell's scree plot was used, which represents the eigenvalues of the correlation matrix in descending order. The factor number extracted corresponds to the eigenvalue that produces the inflection point in the curve (eigenvalue >1) and explains the minimum variance ($>10\%$ for each component). Variables were defined as associated with a factor if they presented loads ≥ 0.30 ³⁰ (the closer to 1, the greater the association). Oblique rotation (promax) was performed to allow for better interpretation of the factor analysis.³⁰

Patient and public involvement

Patients and public were involved in neither the design of the research question nor in developing plans for the design or implementation of the study. The study had no patient advisers. Outcomes were self-reported by patients based on predefined questions. Due to the cross-sectional nature of the study, feedback regarding the results was not planned for those involved.

RESULTS

Participant characteristics

Table 1 shows the characteristics of the participants, prevalence of any chronic disease and multimorbidity. The sample was composed of 4001 adults and had a response rate of 76%. Women constituted over half of the sample. About one-half of the interviewees were aged between 25 and 44 years, and 81% were black, brown or indigenous. The predominant social stratum was the lower middle class (57%), and approximately one-third of the participants were students or housewives. More than half reported good health status (54%), and the majority had a physician visit in the last year (76%). In the last 12 months, 7% reported dengue and 6% reported malaria. One-half of respondents reported seeking a tertiary health service when they needed care (47%). The prevalence of any chronic disease was 57.2% (95% CI 56.6% to 58.7%), with a mean $\pm$ SD of 1.2 $\pm$ 1.5 chronic disease per person. This average increased with age (0.5 $\pm$ 0.8 in the group aged 18–24 years and 2.5 $\pm$ 1.9 in those 60 years or above).

Prevalence of multimorbidity

The prevalence of ≥ 2 chronic conditions was 29% (95% CI 27.6% to 30.5%), and that of ≥ 3 chronic diseases was 15.2% (95% CI 14.1% to 16.4%). Higher prevalence was observed in women, in widowers, in individuals with lower education, in retired individuals, in individuals who had the worst perceptions of health and in those who visited a doctor and were hospitalised in the last year than in others (table 1). In the previous year, dengue was reported by 44% of those who had two or more chronic conditions.

Approximately half of women aged 35–59 years reported ≥ 2 morbidities (table 2). Back pain was the most frequently reported health problem in both women and men, followed by hypertension. Women ≥ 60 years with two or more morbidities reported more hypertension (92.0%) than men did in the same age group (79.5%).

Table 3 shows the results obtained for factors associated with multimorbidity. After adjustment, multimorbidity (≥ 2 diseases) was associated with female sex (PR=1.66, 95% CI 1.50 to 1.83), age between 45 and 59 years (PR=4.36, 95% CI 3.48 to 5.46) and age ≥ 60 years (PR=5.68, 95% CI 4.51 to 7.15). The presence of ≥ 3 diseases was associated with female sex (PR=2.19, 95% CI 1.88 to 2.56), age 45–59 years (PR=7.62, 95% CI 5.22 to 11.10), age ≥ 60 years (PR=12.03, 95% CI 8.20 to 17.66), dengue in the last 12 months (PR=1.36, 95% CI 1.13 to 1.64) and very poor health status (PR=7.89, 95% CI 4.71 to 13.23). Having ≥ 3 chronic conditions increased the demand for physician visits, hospitalisation in the last year and demand for the same health service. Education, income, occupation and malaria in the last 12 months did not show associations with multimorbidity.

The factor analysis is presented in table 4. The KMO coefficient was 0.82 for women and 0.78 for men, and the Bartlett sphericity test presented a p value ≤ 0.001 for both, suggesting an adequate factor analysis. In women, one multimorbidity pattern (see online supplemental figure 1) explained 81% of the total variance, including the 12 chronic diseases analysed. In men, two factors were identified (see online supplemental figure 2). In the first factor, heart diseases, chronic kidney disease, stroke, arthritis or rheumatism, chronic spinal problems, depressive disorders, asthma or bronchitis and lung diseases were the associated chronic diseases, which explained a total of 62% of the variance. The second factor was essentially cardiometabolic, which explained 56% of the variance.

DISCUSSION

More than half of the adults had some chronic disease. The occurrence of two or more morbidities was reported by more than a quarter of the adults. Four out of 25 individuals reported a multimorbidity of three or greater. Female sex, elderly age, dengue in the last year, poor health status, seeking the same healthcare service when in need of attendance, physician visits and hospitalisation presented associations with multimorbidity. Chronic spinal problems were the most commonly reported diseases.

We used a list of 12 self-reported chronic conditions—some of which were very broad—to assess the primary outcome of this study. A systematic review summarised 39 observational studies from 1993 to 2013 and identified a range of 5–335 diseases for the study of multimorbidity.³ In previous studies, the fewer diseases included in the research was, the lower the prevalence observed became.⁵³⁴ Regardless of the number of chronic conditions reported

Table 1 Characteristics of participants and prevalence of multimorbidity (%), Manaus metropolitan region, Brazil, 2015 (n=4 001)*

Variable	n (%)	Any chronic disease % (95% CI)	P values	Mean number of chronic health problems (95% CI)	Multimorbidity % (95% CI)		P values
					≥2	≥3	
Overall		57.2 (55.6 to 58.7)		1.17 (1.1 to 1.2)	29.0 (27.6 to 30.5)	15.2 (14.1 to 16.4)	
Sex			<0.001				<0.001
Male	1888 (47.2)	52.1 (49.8 to 54.0)		0.9 (0.8 to 0.9)	21.5 (19.7 to 23.4)	9.3 (8.1 to 10.7)	
Female	2113 (52.7)	61.8 (59.7 to 63.8)		1.39 (1.3 to 1.4)	35.8 (33.8 to 37.8)	20.5 (18.8 to 22.3)	
Age (years)			<0.001				<0.001
18–24	838 (20.8)	37.0 (33.8 to 40.3)		0.5 (0.4 to 0.5)	9.8 (7.9 to 12.0)	3.5 (2.5 to 5.0)	
25–34	1152 (28.7)	49.6 (46.7 to 52.5)		0.8 (0.7 to 0.8)	18.6 (16.4 to 20.9)	6.9 (5.6 to 8.5)	
35–44	843 (21.1)	61.0 (57.6 to 64.2)		1.1 (1.0 to 1.2)	30.0 (27.0 to 33.2)	12.7 (10.6 to 15.1)	
45–59	772 (19.3)	71.4 (68.0 to 74.4)		1.7 (1.6 to 1.8)	46.6 (43.1 to 50.2)	28.1 (25.1 to 31.4)	
≥60	396 (9.9)	86.1 (82.3 to 89.2)		2.5 (2.3 to 2.7)	63.5 (58.6 to 68.1)	44.4 (39.6 to 49.3)	
Marital status			<0.001				<0.001
Single	2173 (54.2)	51.5 (49.3 to 53.5)		0.9 (0.8 to 0.9)	20.8 (19.1 to 22.6)	10.3 (8.8 to 11.3)	
Married	1409 (35.2)	62.0 (59.4 to 64.5)		1.3 (1.2 to 1.4)	35.3 (32.9 to 37.9)	19.6 (17.6 to 21.8)	
Separated/divorced	260 (6.5)	65.6 (59.6 to 71.2)		1.5 (1.3 to 1.7)	42.6 (36.8 to 48.7)	19.9 (15.5 to 25.2)	
Widower	159 (4.0)	79.3 (72.3 to 84.9)		2.2 (1.9 to 2.5)	63.1 (55.3 to 70.2)	39.9 (32.5 to 47.7)	
Skin colour			<0.001				0.0016
White/yellow	774 (19.3)	51.7 (48.2 to 55.2)		1.1 (1.0 to 1.2)	27.1 (24.1 to 30.4)	15.7 (13.3 to 18.5)	
Black/brown/indigenous	3227 (80.5)	58.5 (56.8 to 60.2)		1.2 (1.1 to 1.2)	29.5 (27.9 to 31.1)	15.1 (13.9 to 16.4)	
Schooling level			<0.001				<0.001
High education or above†	158 (3.9)	57.1 (49.3 to 64.6)		.2 (1.0 to 1.5)	32.6 (25.7 to 40.4)	16.1 (11.7 to 23.7)	
High school	1903 (47.5)	51.0 (48.8 to 53.3)		0.9 (0.8 to 1.0)	22.5 (20.7 to 24.5)	9.0 (7.8 to 10.4)	
Middle school	649 (16.2)	50.0 (46.2 to 53.8)		0.9 (0.8 to 1.0)	21.1 (18.2 to 24.4)	10.7 (0.8 to 13.3)	
Elementary school or less	1291 (32.2)	69.9 (67.4 to 72.4)	<0.001	1.7 (1.6 to 1.8)	42.1 (39.5 to 44.8)	26.5 (24.1 to 28.9)	
Economic classification‡							<0.001
A–B	629 (15.7)	51.2 (47.2 to 55.1)		0.9 (0.8 to 1.1)	24.6 (21.4 to 28.2)	11.1 (8.8 to 13.8)	
C	2285 (57.1)	55.5 (53.4 to 57.5)		1.0 (1.0 to 1.1)	26.1 (24.3 to 27.9)	13.2 (11.9 to 14.7)	
D–E	1087 (27.2)	64.3 (61.4 to 67.2)		1.5 (1.4 to 1.6)	37.7 (34.9 to 40.7)	21.8 (19.4 to 24.3)	
Occupation			<0.001				<0.001
Formal job	761 (19.0)	52.7 (49.1 to 56.2)		0.9 (0.8 to 1.0)	22.7 (19.8 to 25.8)	9.2 (7.3 to 11.5)	
Informal job	1149 (28.7)	55.9 (53.0 to 58.7)		1.0 (1.0 to 1.1)	27.6 (25.1 to 30.3)	12.9 (11.1 to 15.0)	
Retired	315 (7.9)	79.2 (74.3 to 83.3)		2.4 (2.2 to 2.6)	60.5 (55.4 to 65.8)	43.4 (38.0 to 49.0)	

Continued

Table 1 Continued

Variable	n (%)	Any chronic disease % (95% CI)	P values	Mean number of chronic health problems (95% CI)	Multimorbidity % (95% CI)		P values
					≥2	≥3	
Student/housewife	1199 (29.8)	55.0 (52.2 to 57.8)		1.1 (1.1 to 1.2)	29.0 (25.5 to 31.6)	15.8 (13.8 to 18.0)	
Unemployed	577 (14.4)	58.3 (54.2 to 62.3)		1.0 (0.9 to 1.1)	23.1 (19.8 to 26.7)	11.2 (8.9 to 14.1)	
Private health insurance			0.348			0.697	0.616
Yes	523 (13.0)	55.3 (51.0 to 59.5)		1.0 (0.9 to 1.29)	28.0 (24.6 to 32.3)	14.5 (11.7 to 17.8)	
No\$	3478 (87.0)	57.5 (55.8 to 59.1)		1.2 (1.1 to 1.2)	29.0 (27.6 to 30.7)	15.3 (14.2 to 16.6)	
Health status			<0.001		<0.001	<0.001	<0.001
Very good	471 (11.9)	30.8 (26.8 to 35.2)		0.4 (0.47 to 0.5)	9.7 (7.3 to 12.7)	3.1 (1.9 to 5.1)	
Good	2175 (54.3)	50.4 (48.3 to 52.5)		0.8 (0.8 to 0.9)	20.3 (18.7 to 22.0)	7.9 (6.8 to 9.1)	
Fair	1108 (27.7)	75.1 (72.5 to 77.5)		1.7 (1.6 to 1.8)	45.2 (42.3 to 48.1)	26.0 (23.5 to 28.6)	
Bad	193 (4.8)	88.6 (83.2 to 93.3)		2.6 (2.4 to 2.9)	68.4 (61.6 to 74.6)	50.3 (43.3 to 57.3)	
Very bad	54 (1.3)	81.5 (68.9 to 89.7)		3.6 (3.0 to 4.3)	77.7 (64.8 to 86.9)	70.3 (57.0 to 81.0)	
City			<0.001		<0.001	<0.001	<0.001
Capital	3479 (86.8)	58.4 (56.8 to 60.0)		1.2 (1.1 to 1.2)	30.2 (28.7 to 31.7)	15.8 (14.6 to 17.1)	
Countryside	522 (13.1)	49.2 (44.9 to 53.5)		0.9 (0.8 to 1.0)	21.5 (18.2 to 25.3)	11.3 (0.8 to 14.3)	
Health reference¶			<0.001		<0.001	<0.001	<0.001
Yes	2434 (60.7)	62.3 (60.3 to 64.2)		1.3 (1.2 to 1.4)	34.4 (31.5 to 36.3)	18.7 (17.1 to 20.3)	
No	1567 (39.2)	49.4 (46.9 to 51.9)		0.9 (0.8 to 0.9)	20.8 (18.9 to 22.9)	9.8 (17.2 to 20.3)	
Physician visit			<0.001		<0.001	<0.001	<0.001
Yes	3066 (76.5)	60.8 (59.0 to 62.5)		1.2 (1.2 to 1.3)	32.1 (30.5 to 33.8)	17.4 (16.1 to 18.8)	
No	935 (23.4)	45.5 (42.3 to 48.7)		0.8 (0.7 to 0.9)	18.9 (16.5 to 21.5)	8.2 (6.6 to 10.2)	
Hospitalisation			<0.001		<0.001	<0.001	<0.001
Yes	273 (7.0)	72.9 (67.3 to 77.8)		1.2 (1.2 to 1.3)	47.2 (41.4 to 53.2)	28.1 (23.1 to 33.8)	
No	3728 (93.0)	56.1 (67.3 to 77.8)		0.8 (0.7 to 0.9)	27.7 (26.3 to 29.2)	14.3 (13.2 to 15.4)	
Dengue			0.002		<0.001	<0.001	<0.001
Yes	281 (7.0)	65.9 (60.1 to 71.2)		1.6 (1.4 to 1.8)	44.2 (38.5 to 50.1)	27.6 (22.7 to 33.1)	
No	3720 (93.0)	56.5 (54.9 to 58.1)		1.1 (1.0 to 1.1)	27.9 (26.5 to 29.3)	14.3 (13.2 to 15.5)	
Malaria			0.420		0.122	0.014	
Yes	234 (5.9)	54.6 (48.2 to 60.9)		1.3 (1.1 to 1.5)	33.5 (27.7 to 39.8)	20.8 (16.1 to 26.5)	
No	3767 (94.0)	57.3 (0.55 to 0.58)		1.1 (1.1 to 1.2)	28.8 (27.3 to 30.2)	14.9 (13.8 to 16.0)	
Medical attention			<0.001		0.048	0.531	
Primary	1208 (30.2)	60.8 (58.0 to 63.5)		1.2 (1.1 to 1.3)	29.9 (27.3 to 32.5)	15.2 (13.3 to 17.3)	

Continued

Table 1 Continued

Variable	n (%)	Any chronic disease % (95% CI)	P values	Mean number of chronic health problems (95% CI)	Multimorbidity % (95% CI)	
					≥2	≥3
Secondary	598 (14.9)	62.3 (58.3 to 66.1)		1.3 (1.2 to 1.5)	33.0 (29.4 to 36.9)	18.9 (15.9 to 22.2)
Tertiary	1886 (47.2)	54.0 (51.8 to 56.3)		1.1 (1.0 to 1.2)	27.7 (25.7 to 29.8)	14.3 (12.8 to 15.9)
Others	309 (7.7)	52.7 (47.2 to 58.3)		1.0 (0.9 to 1.2)	26.2 (21.6 to 31.4)	14.2 (10.7 to 18.5)

Physician visit, hospitalisation, dengue and malaria in last 12 months.

*Descriptive statistics using simple frequency and Pearson's χ^2 test.

†Tertiary education or higher.

‡Average household income in 2015: A–B, US\$6500–US\$1419; C, US\$463–US\$772; D–E, US\$205.

§People who use the public health service.

¶Seeking the same healthcare service when in need of attendance.

and how they were defined, multimorbidity estimates are influenced by self-report. Although widely applied,^{3 5 35} such assessments are more likely to suffer classification bias or have no validated instrument for confirmation. In the present research, over-reporting or under-reporting may have occurred,³⁶ as well as recall bias, which is more common in elderly individuals of lower socioeconomic and educational levels than among other individuals.^{7 37} In addition, we did not investigate disease severity. Previous studies recommend inquiring about the degree of disease intensity and diseases' interference with routine activities and disabilities.^{5 35} More reliable estimates of multimorbidity, using medical records, for example, are not available in the region.

The response rate was 76%, which may constitute a source of selection bias. Efforts were made to improve representativeness by using predefined sex and age quotas and interviewing one individual per family, according to official estimates.³⁸ Survival bias may have also influenced the results, as patients who died prematurely from those causes, who were hospitalised or who had more serious diseases were not available in the household to participate in the survey. The cross-sectional nature of the study does not allow for investigation of temporal associations.

This is the first local study to estimate the prevalence of multimorbidity in adults in the state of Amazonas. We used a cut-off point of ≥ 2 and ≥ 3 chronic diseases, as performed in previous studies.^{3 5} We identified the most vulnerable multimorbidity groups to be women and the elderly. Multimorbidity was higher in older people and increased with age; this finding has been observed in previous studies.^{5 9 23} The National Health Survey conducted in Brazil in 2013 reported that women are most affected among all socioeconomic groups, especially the elderly.³⁹

Our results showed similarities to a cross-sectional study conducted in 2012 in Pelotas city in the southern region of Brazil with 2927 subjects, in which 29.1% of the interviewees had more than two chronic diseases and 14% had three or more.⁴⁰ The 2013 National Health Survey also confirmed these findings: 22% of Brazilians reported two or more chronic diseases, and 10% were affected by more than three.¹⁴ The highest prevalence was observed in the south (26%–29%),¹⁴ which is more economically developed and has greater access to health services than the north does.^{17 41} Any chronic disease occurred in 45% of Brazilians, with a lower prevalence in the north region.⁴²

In other contexts, lower prevalence of multimorbidity was found. A survey conducted in 2012 in Italy, with 3 759 836 adults, detected that 15% of individuals presented two or more chronic diseases.¹⁰ In Ireland, a representative sample of the population (11.3% of subjects ≥ 50 years) presented multiple diseases.¹² Furthermore, an electronic medical data analysis conducted in 2007 with 1 751 841 users of the Scottish Health Service found that 23% had multimorbidity.⁹ In an economic context comparable to Brazil's, a population-based Indian study conducted in 2007 with 10 973 interviewees identified

Table 2 Prevalence (%) of most common diseases stratified by sex, age and multimorbidity group, Manaus metropolitan region, Brazil, 2015*

Morbidities	n (%)	Multimorbidity $\geq$ 2			Multimorbidity $\geq$ 3		
		18–34 n (%)	35–59 n (%)	$\geq$ 60 n (%)	18–34 n (%)	35–59 n (%)	$\geq$ 60 n (%)
Women (2113)		195 (18.4)	394 (47.0)	168 (76.9)	78 (7.0)	236 (28.1)	120 (55.0)
Chronic spinal problem†	747 (35.3)	135 (42.6)	248 (75.6)	99 (96.1)	64 (20.1)	176 (53.6)	78 (75.7)
Hypertension	516 (24.4)	78 (62.3)	211 (83.0)	126 (92.0)	94 (32.7)	149 (58.7)	98 (71.6)
Arthritis or rheumatism	414 (19.5)	66 (79.5)	197 (91.6)	110 (94.8)	33 (39.7)	146 (67.7)	92 (79.3)
Hypercholesterolaemia	425 (20.1)	66 (79.5)	217 (90.8)	100 (97.0)	46 (55.4)	149 (62.4)	85 (82.5)
Diabetes	157 (7.4)	15 (93.7)	82 (90.2)	48 (96.0)	14 (87.4)	70 (76.6)	42 (84.5)
Asthma or asthmatic bronchitis	155 (7.3)	50 (68.5)	54 (85.6)	19 (100.0)	28 (38.3)	46 (73.0)	19 (100.0)
Depressive disorder	158 (7.4)	44 (71.0)	69 (91.9)	21 (100.0)	19 (30.6)	55 (73.2)	20 (95.2)
Heart disease ‡	119 (5.6)	23 (71.8)	52 (98.1)	34 (100.0)	17 (53.0)	46 (86.7)	32 (94.0)
Men (1888)		102 (10.9)	220 (28.2)	84 (47.1)	32 (3.4)	89 (11.4)	56 (31.4)
Chronic spinal problem†	662 (35.0)	77 (27.5)	149 (49.7)	54 (65.8)	20 (7.1)	63 (21.0)	42 (51.4)
Hypertension	271 (14.4)	37 (77.1)	112 (76.7)	61 (79.5)	17 (35.3)	60 (41.1)	45 (58.6)
Arthritis or rheumatism	179 (9.5)	25 (89.3)	80 (86.8)	50 (84.9)	13 (46.1)	45 (48.9)	38 (64.5)
Hypercholesterolaemia	171 (9.0)	24 (57.1)	85 (87.6)	28 (87.6)	14 (33.3)	48 (49.4)	25 (78.2)

P values of all variables were ≤ 0.002 .

*Multimorbidity with prevalence $\geq 5\%$.

†Chronic back pain or neck, low back pain, sciatica, vertebral or disc pain.

‡Heart disease, or heart attack, angina, cardiac insufficiency.

smaller proportions (28% had any chronic disease and approximately 9% had multimorbidity).⁴³

Two findings of our research are rarely described in previous studies: the higher frequency of multimorbidity at younger ages and the lack of association with economic status. One-half of adults aged 25–34 years and nearly two-thirds of interviewees aged 35–44 years reported any chronic condition, and nearly one-third had multimorbidity. The development of multimorbidity in young adults is in agreement with previous data from Brazil.^{23 39 40} It is important to emphasise that half of the population of the Manaus metropolitan region is concentrated in this age range (49% aged 25–44 years). A systematic review of 24 cross-sectional studies on multimorbidity found income as a conflicting factor across studies, associated either with richer or poorer individuals, while lower educational attainment was associated with a 64% higher chance of multimorbidity.⁴⁴ No association was found between income and multimorbidity after adjusting for socioeconomic variables.

In low-income countries such as Brazil, which also faces economic austerity policies, rising unemployment and unstable social and health policies,⁴⁵ it is possible to predict a reduction in access to health services, with a consequent increase in morbidity. This effect has been observed in other austerity scenarios, in which this type of policy reduced jobs, education and use of health services, resulting in an increase of chronic diseases.^{46 47} An analysis of high-income countries found that 3.6 more years

of education reduces the risk of cardiovascular disease by one-third.⁴⁸

Back pain was the most frequent disease, reported by one-third of the sample. In our study, this morbidity was assessed using several questions, which may have increased the sensitivity of the assessment. An even higher proportion (49%) of vertebral spine/back issues was observed in a representative cross-sectional survey of Brazilian adults.³⁹ In other contexts, similar prevalence values were estimated.⁴⁹

The presence of dengue was higher in individuals with multimorbidity, possibly due to the lower immunological response observed in chronic diseases such as diabetes, rheumatoid arthritis and asthma.^{50 51} A systematic review of 16 cohort and case-control studies from 2007 to 2013 showed chronic diseases as risk factors for severe dengue.⁵² In another meta-analysis of 10 studies conducted between 2006 and 2014,⁵³ diabetes was significantly associated with haemorrhagic dengue: regardless of demographic and socioeconomic characteristics, the association was 5% higher than that for individuals who did not have diabetes.

A single multimorbidity pattern with all investigated diseases was identified in women. Heart disease presented the highest factor loading, but disease patterns are poorly explained due to the wide range of diseases included in one factor. This finding may be due to our measurement and analytical approach, including sex stratification, broad categories of diseases and the number of chronic

Table 3 Adjusted prevalence ratio (PR) and 95% CI for any chronic disease and multimorbidity ≥ 2 and ≥ 3 , according to sociodemographic and health variables based on hierarchical Poisson regression; Manaus metropolitan region, Brazil, 2015 (n=4 001)

Variable	Any chronic disease PR (95% CI)	P values	Multimorbidity PR (95% CI)			
			≥2	P values	≥3	P values
Demographic block*						
Sex						
Male	1.00		1.00		1.00	
Female	1.19 (1.12 to 1.25)	<0.001	1.66 (1.50 to 1.83)	<0.001	2.19 (1.88 to 2.56)	<0.001
Age group (years)						
18–24	1.00				1.00	
25–34	1.33 (1.19 to 1.48)	<0.001	1.81 (1.42 to 2.30)	<0.001	1.88 (1.24 to 2.84)	0.003
35–44	1.63 (1.47 to 1.81)	<0.001	2.85 (2.26 to 3.60)	<0.001	3.40 (2.28 to 5.06)	<0.001
45–59	1.91 (1.72 to 2.12)	<0.001	4.36 (3.48 to 5.46)	<0.001	7.62 (5.22 to 11.10)	<0.001
≥60	2.32 (2.08 to 2.57)	<0.001	5.68 (4.51 to 7.15)	<0.001	12.03 (8.20 to 17.66)	<0.001
Marital status						
Single	1.00		1.00		1.00	
Married	1.09 (0.96 to 1.07)	0.521	1.20 (1.07 to 1.33)	0.001	1.21 (1.03 to 1.42)	0.017
Separated/divorced	1.00 (0.91 to 1.10)	0.961	1.24 (1.06 to 1.45)	0.006	0.93 (0.72 to 1.21)	0.620
Widower	0.96 (0.87 to 1.06)	0.475	1.18 (1.01 to 1.38)	0.032	0.99 (0.77 to 1.26)	0.962
Skin colour						
White/yellow	1.00		1.00		1.00	
Black/brown/Indigenous	1.10 (1.02 to 1.18)	0.006	1.04 (0.93 to 1.16)	0.474	0.89 (0.75 to 1.05)	0.175
Socioeconomic block†						
Education						
High education or above	1.00		1.00		1.00	
High school	1.00 (0.87 to 1.14)	0.976	0.89 (0.71 to 1.12)	0.357	0.73 (0.50 to 1.05)	0.091
Middle school	0.98 (0.84 to 1.13)	0.754	0.81 (0.63 to 1.06)	0.135	0.80 (0.53 to 1.20)	0.289
Elementary school or less	1.12 (0.98 to 1.29)	0.106	1.01 (0.80 to 1.28)	0.877	1.08 (0.75 to 1.55)	0.667
Economic classification						
A–B	1.00		1.00		1.00	
C	1.04 (0.96 to 1.13)	0.361	1.01 (0.87 to 1.17)	0.881	1.05 (0.83 to 1.33)	0.656
D–E	1.06 (0.97 to 1.16)	0.207	1.15 (0.98 to 1.36)	0.075	1.15 (0.89 to 1.49)	0.254
Occupation						
Formal job	1.00		1.00		1.00	
Informal job	0.95 (0.87 to 1.03)	0.223	0.97 (0.83 to 1.13)	0.728	0.98 (0.76 to 1.28)	0.929
Retired	0.95 (0.86 to 1.06)	0.363	1.11 (0.92 to 1.32)	0.258	1.39 (1.02 to 1.88)	0.033
Student/housewife	0.96 (0.87 to 1.05)	0.331	0.97 (0.83 to 1.14)	0.767	1.05 (0.81 to 1.36)	0.671
Unemployed	1.09 (0.99 to 1.20)	0.081	0.99 (0.81 to 1.20)	0.949	1.13 (0.83 to 1.54)	0.404
Health block‡						
Private health insurance						
No	1.00		1.00		1.00	
Yes	0.99 (0.91 to 1.07)	0.913	1.03 (0.90 to 1.18)	0.593	1.04 (0.84 to 1.28)	0.771
Health status						
Very good	1.00		1.00		1.00	
Good	1.43 (1.24 to 1.64)	<0.001	1.63 (1.23 to 2.15)	<0.001	1.81 (1.09 to 2.99)	0.020
Fair	1.94 (1.69 to 2.23)	<0.001	2.84 (2.15 to 3.76)	<0.001	4.21 (2.56 to 6.93)	<0.001
Bad	2.01 (1.80 to 2.41)	<0.001	3.53 (2.64 to 4.71)	<0.001	6.25 (3.74 to 10.47)	<0.001

Continued

Table 3 Continued

Variable	Any chronic disease PR (95% CI)	P values	Multimorbidity PR (95% CI)			
			≥2	P values	≥3	P values
Very bad	1.91 (1.60 to 2.27)	<0.001	3.70 (2.73 to 5.00)	<0.001	7.89 (4.71 to 13.23)	<0.001
Health reference§						
No	1.00		1.00		1.00	
Yes	1.12 (1.06 to 1.19)	<0.001	1.33 (1.19 to 1.47)	<0.001	1.40 (1.20 to 1.63)	<0.001
Physician visit						
No	1.00		1.00		1.00	
Yes	1.13 (1.05 to 1.22)	<0.001	1.22 (1.07 to 1.40)	0.002	1.33 (1.09 to 1.64)	0.005
Hospitalisation						
No	1.00		1.00		1.00	
Yes	1.18 (1.09 to 1.27)	<0.001	1.36 (1.20 to 1.54)	<0.001	1.43 (1.17 to 1.74)	<0.001
Dengue						
No	1.00		1.00		1.00	
Yes	1.07 (0.99 to 1.16)	0.079	1.23 (1.08 to 1.41)	0.001	1.36 (1.13 to 1.64)	0.001
Malaria						
No	1.00		1.00		1.00	
Yes	0.98 (0.82 to 1.01)	0.109	0.96 (0.81 to 1.13)	0.653	0.99 (0.78 to 1.26)	0.947
Type of service						
Primary	1.00		1.00		1.00	
Secondary	1.02 (0.95 to 1.09)	0.518	1.07 (0.95 to 1.22)	0.265	1.15 (0.95 to 1.40)	0.129
Tertiary	0.95 (0.89 to 1.00)	0.086	1.01 (0.92 to 1.12)	0.780	1.05 (0.90 to 1.25)	0.489
Outros	0.92 (0.82 to 1.03)	0.180	0.97 (0.80 to 1.17)	0.723	1.13 (0.85 to 1.51)	0.374

Significant variables kept in each block of analysis:

Any chronic disease: *sex, age, marital status and race; †sex, age, marital status, education, income, occupation; ‡sex, age, marital status, income, occupation, private health insurance, health status, health reference, medical consultation in last 12 months, hospital admission in last 12 months, dengue, malaria and type of health service that usually comes.

≥2 morbidities: *sex, age, marital status and race; †sex, age, marital status, education, income, occupation; ‡sex, age, marital status, income, occupation, private health insurance, health status, health reference, medical consultation in last 12 months, hospital admission in last 12 months, dengue, malaria and type of health service that usually comes.

≥3 morbidities: *sex, age, marital status and race; †sex, age, marital status, education, income, occupation; ‡sex, age, marital status, income, occupation, private health insurance, health status, health reference, medical consultation in last 12 months, hospital admission in last 12 months, dengue, malaria and type of health service that usually comes.

§Seeking the same healthcare service when in need of attendance.

conditions investigated. In previous studies conducted in Brazil involving similar questions, up to three multimorbidity patterns have been identified: cardiometabolic, musculoskeletal-mental and respiratory.^{14 23 40} Such studies did not stratify by sex when investigating the multimorbidity pattern. An Australian cohort with 13 715 women born between 1946 and 1951 identified five multimorbidity patterns (psychosomatic, musculoskeletal, cardiometabolic, cancer and respiratory) after investigating 18 chronic diseases and 13 symptoms.⁵⁴ The greater number of diseases and symptoms may explain the number of clustering factors in women relative to our analysis (31 vs 12).

In men, lung disease was the disease with the highest factorial loading on factor 1, but no clear pattern of diseases was found in the clustering of this factor. Factor 2 included cardiometabolic diseases, which could be explained by similar risk factors such as sedentary lifestyle

and obesity. An analysis of 2008 electronic medical records from the Spanish National Health System identified cardiometabolic patterns in both men and women in different age ranges.⁵⁵

According to the results, it is estimated that over 1 300 000 residents of the metropolitan region of Manaus have a chronic condition, and nearly 700 000 have multimorbidity. Early diagnosis and treatment of chronic diseases, centred on primary care services, is a priority for enabling sustainability of the health system and a healthier society.⁵⁶

CONCLUSION

Multimorbidity was common in residents of the metropolitan region of Manaus and was associated with female sex, elderly people and poorer health perception. Prevention and control strategies should prioritise these groups. Future analyses should investigate the relationship

Table 4 Factor score of each chronic disease in women and men, Manaus metropolitan region, Brazil, 2015

Morbidities	Women (n=2113)	Men (n=1888)	
	Factor 1	Factor 1	Factor 2
Hypercholesterolaemia	0.70		0.62
Hypertension	0.64		0.78
Heart disease	0.72	0.40	0.47
Diabetes	0.63		0.85
Chronic kidney disease	0.46	0.73	
Stroke	0.66	0.61	
Arthritis	0.65	0.40	0.35
Chronic spinal problem	0.46	0.44	
Depressive disorder	0.49	0.58	
Asthma or asthmatic bronchitis	0.44	0.48	
Lung diseases*	0.53	0.76	
Cancer†	0.52		
Proportion of variance (%)	81.0	62.0	56.0
Kaiser-Meyer-Olkin	0.82	0.78	

Kept factors were those with scores ≥ 0.30 after oblique rotation.

*Pulmonary emphysema, bronchitis, chronic obstructive lung disease.

†Cancer showed negative values in men and was excluded.

between multimorbidity and the use and costs of health services in the region.

Author affiliations

¹Post-Graduate Program Health Sciences, University of Brasília, Brasília, Brazil

²Getúlio Vargas University Hospital, Federal University of Amazonas, Manaus, Brazil

³Faculty Medicine, Federal University of Amazonas, Manaus, Brazil

⁴Post-Graduate Program of Pharmaceutical Sciences, University of Sorocaba, Sorocaba, Brazil

⁵Faculty of Pharmaceutical Sciences, State University of Campinas, Campinas, Brazil

⁶Department of Nursing in Public Health, Federal University of Pelotas, Pelotas, Brazil

⁷Faculty of Medicine, University of Brasília, Brasília, Brazil

Contributors MTS, TFG, and MeAA designed the study. MeAA, MTS and BNP performed the statistical analyses. MeAA, MTS, TFG, BNP and MGP interpreted the results and drafted the manuscript. All authors made critical revisions and provided intellectual content to the manuscript, approved the final version and agreed to be accountable for all aspects of this work.

Funding This work was funded by the National Council for Scientific and Technological Development (Conselho Nacional para o Desenvolvimento Científico e Tecnológico (CNPq)) (404990/2013-4 and 448093/2014-6 to MTS) and the Foundation for Research Support of the State of Amazonas (Fundação de Amparo à Pesquisa do Estado do Amazonas (FAPEAM) to MeAA).

Competing interests None declared.

Patient consent Not required.

Ethics approval The research project was approved by the Research Ethics Committee of the Federal University of Amazonas.

Provenance and peer review Not commissioned; externally peer reviewed.

Data sharing statement Data are available and can be accessed by contacting MTS.

Open access This is an open access article distributed in accordance with the Creative Commons Attribution Non Commercial (CC BY-NC 4.0) license, which permits others to distribute, remix, adapt, build upon this work non-commercially, and license their derivative works on different terms, provided the original work is properly cited, appropriate credit is given, any changes made indicated, and the use is non-commercial. See: <http://creativecommons.org/licenses/by-nc/4.0/>.

REFERENCES

- van den Akker M, Buntinx F, Knottnerus JA. Comorbidity or multimorbidity: what's in a name? A review of literature. *The European Journal of General Practice* 1996;2:65–70.
- Le Reste JY, Nabbe P, Manceau B, et al. The European General Practice Research Network presents a comprehensive definition of multimorbidity in family medicine and long term care, following a systematic review of relevant literature. *J Am Med Dir Assoc* 2013;14:319–25.
- Violan C, Foguet-Boreu Q, Flores-Mateo G, et al. Prevalence, determinants and patterns of multimorbidity in primary care: a systematic review of observational studies. *PLoS One* 2014;9:e102149.
- Diederichs C, Berger K, Bartels DB. The measurement of multiple chronic diseases—a systematic review on existing multimorbidity indices. *J Gerontol A Biol Sci Med Sci* 2011;66:301–11.
- Fortin M, Stewart M, Poitras ME, et al. A systematic review of prevalence studies on multimorbidity: toward a more uniform methodology. *Ann Fam Med* 2012;10:142–51.
- GBD 2015 Mortality and Causes of Death Collaborators. Global, regional, and national life expectancy, all-cause mortality, and cause-specific mortality for 249 causes of death, 1980–2015: a systematic analysis for the Global Burden of Disease Study 2015. *Lancet* 2016;388:1459–544.
- Hussain MA, Huxley RR, Al Mamun A. Multimorbidity prevalence and pattern in Indonesian adults: an exploratory study using national survey data. *BMJ Open* 2015;5:e009810.
- Ryan A, Wallace E, O'Hara P, et al. Multimorbidity and functional decline in community-dwelling adults: a systematic review. *Health Qual Life Outcomes* 2015;13:168.
- Barnett K, Mercer SW, Norbury M, et al. Epidemiology of multimorbidity and implications for health care, research, and medical education: a cross-sectional study. *Lancet* 2012;380:37–43.
- Lenzi J, Avaldi VM, Rucci P, et al. Burden of multimorbidity in relation to age, gender and immigrant status: a cross-sectional study based on administrative data. *BMJ Open* 2016;6:e012812.
- Lima-Costa MF, De Oliveira C, Macinko J, et al. Socioeconomic inequalities in health in older adults in Brazil and England. *Am J Public Health* 2012;102:1535–41.
- Glynn LG, Valderas JM, Healy P, et al. The prevalence of multimorbidity in primary care and its effect on health care utilization and cost. *Fam Pract* 2011;28:516–23.
- Nagl A, Witte J, Hodek JM, et al. Relationship between multimorbidity and direct healthcare costs in an advanced elderly population. Results of the PRISCUS trial. *Z Gerontol Geriatr* 2012;45:146–54.
- Nunes BP, Chiavegatto Filho ADP, Pati S, et al. Contextual and individual inequalities of multimorbidity in Brazilian adults: a cross-sectional national-based study. *BMJ Open* 2017;7:e015885.
- de Souza Santos Machado V, Valadares AL, da Costa-Paiva LS, et al. Multimorbidity and associated factors in Brazilian women aged 40 to 65 years: a population-based study. *Menopause* 2012;19:569–75.
- Nunes BP, Thumé E, Facchini LA. Multimorbidity in older adults: magnitude and challenges for the Brazilian health system. *BMC Public Health* 2015;15:1172.
- Pinto DGC, Costa MAC, MLDAC M. *Índice de desenvolvimento humano municipal brasileiro*. 2013. Brasília: Programa das Nações Unidas para o Desenvolvimento, Instituto de Pesquisa Econômica Aplicada, 2017.
- Silva MT, Galvão TF. Use of health services among adults living in Manaus Metropolitan Region, Brazil: population-based survey, 2015. *Epidemiol Serv Saude* 2017;26:725–34.
- Araujo MEA, Silva MT, Galvão TF, et al. Prevalence of health services usage and associated factors in the Amazon region of Brazil: a population-based cross-sectional study. *BMJ Open* 2017;7:e017966.
- Brasil. Instituto Brasileiro de Geografia e Estatística (IBGE). Censo demográfico de 2010. Os indígenas no censo demográfico de 2010. https://ww2.ibge.gov.br/indigenas/indigena_censo2010.pdf (Accessed 28 Jul 2018).
- Viacava F, Bellido JG. Condições de saúde, acesso a serviços e fontes de pagamento, segundo inquéritos domiciliares. *Ciência & Saúde Coletiva* 2016;21:351–70.

22. Kneipp SM, Yarandi HN. Complex sampling designs and statistical issues in secondary analysis. *West J Nurs Res* 2002;24:552–66.
23. Rzewuska M, de Azevedo-Marques JM, Coxon D, et al. Epidemiology of multimorbidity within the Brazilian adult general population: Evidence from the 2013 National Health Survey (PNS 2013). *PLoS One* 2017;12:e0171813.
24. Macinko J, Mullachery P, Silver D, et al. Patterns of Alcohol Consumption and Related Behaviors in Brazil: Evidence from the 2013 National Health Survey (PNS 2013). *PLoS One* 2015;10:e0134153.
25. Associação Brasileira de Empresas de Pesquisa. Critério de Classificação Econômica Brasil. 2015. <http://www.abep.org/criterio-brasil> (accessed 10 Aug 2017).
26. Barros AJ, Hirakata VN. Alternatives for logistic regression in cross-sectional studies: an empirical comparison of models that directly estimate the prevalence ratio. *BMC Med Res Methodol* 2003;3:1.
27. Espelt A, Mari-Dell'Olmo M, Penelo E, et al. Applied prevalence ratio estimation with different regression models: an example from a cross-national study on substance use research. *Adicciones* 2016;29:105–112.
28. Victora CG, Huttly SR, Fuchs SC, et al. The role of conceptual frameworks in epidemiological analysis: a hierarchical approach. *Int J Epidemiol* 1997;26:224–7.
29. Fotouhi AR. Modelling overdispersion in longitudinal count data in clinical trials with application to epileptic data. *Contemp Clin Trials* 2008;29:547–54.
30. Norman G, Streiner D. *Principal Components and factor analysis. fooling around with factors. biostatistics: the bare essentials*. Hamilton: BMJ Publishing Group, 2008.
31. Kim JO, Mueller CW. *Factor analysis: statistical methods and practical issues*: Sage, 1978.
32. Kubinger KD. On artificial results due to using factor analysis for dichotomous variables. *Psychology Science* 2003;45:106–10.
33. Schäfer I, von Leitner EC, Schön G, et al. Multimorbidity patterns in the elderly: a new approach of disease clustering identifies complex interrelations between chronic conditions. *PLoS One* 2010;5:e15941.
34. Pati S, Swain S, Hussain MA, et al. Prevalence and outcomes of multimorbidity in South Asia: a systematic review. *BMJ Open* 2015;5:e007235.
35. Vancampfort D, Koyanagi A, Ward PB, et al. Chronic physical conditions, multimorbidity and physical activity across 46 low- and middle-income countries. *Int J Behav Nutr Phys Act* 2017;14:6.
36. Gross R, Bentur N, Elhayany A, et al. The validity of self-reports on chronic disease: characteristics of underreporters and implications for the planning of services. *Public Health Rev* 1996;24:167–82.
37. Palladino R, Tayu Lee J, Ashworth M, et al. Associations between multimorbidity, healthcare utilisation and health status: evidence from 16 European countries. *Age Ageing* 2016;45:431–5.
38. Instituto Brasileiro de Geografia e Estatística (IBGE). Censo demográfico de. 2010. <http://www.censo2010.ibge.gov.br/sinopse/index.php?uf=13&dados=4> (accessed 20 Jan 2016).
39. Carvalho JN, Roncalli ÂG, Cancela MC, de Camargo Cancela M, et al. Prevalence of multimorbidity in the Brazilian adult population according to socioeconomic and demographic characteristics. *PLoS One* 2017;12:e0174322.
40. Nunes BP, Camargo-Figuera FA, Guttier M, et al. Multimorbidity in adults from a southern Brazilian city: occurrence and patterns. *Int J Public Health* 2016;61:1013–20.
41. Araújo MEA, Silva MT, Andrade KRC, et al. Prevalence of health services utilization in Brazil: a systematic review and meta-analysis. *Epidemiol Serv Saude* 2017;26:587–602.
42. Malta DC, Bernal RT, de Souza MF, MdFM deS, et al. Social inequalities in the prevalence of self-reported chronic non-communicable diseases in Brazil: national health survey 2013. *Int J Equity Health* 2016;15:153.
43. Pati S, Agrawal S, Swain S, et al. Non communicable disease multimorbidity and associated health care utilization and expenditures in India: cross-sectional study. *BMC Health Serv Res* 2014;14:451.
44. Pathirana TI, Jackson CA. Socioeconomic status and multimorbidity: a systematic review and meta-analysis. *Aust N Z J Public Health* 2018;42:186–94.
45. Machado CV, Lima LD, Baptista TWF. Health policies in Brazil in times of contradiction: paths and pitfalls in the construction of a universal system. *Cad Saude Publica* 2017;33(Suppl 2):e00129616.
46. Stuckler D, Reeves A, Loopstra R, et al. Austerity and health: the impact in the UK and Europe. *Eur J Public Health* 2017;27(suppl_4):18–21.
47. Kyriopoulos II, Zavras D, Skroumpelos A, et al. Barriers in access to healthcare services for chronic patients in times of austerity: an empirical approach in Greece. *Int J Equity Health* 2014;13:54.
48. Tillmann T, Vaucher J, Okbay A, et al. Education and coronary heart disease: mendelian randomisation study. *BMJ* 2017;358:j3542.
49. Harrison C, Britt H, Miller G, et al. Examining different measures of multimorbidity, using a large prospective cross-sectional study in Australian general practice. *BMJ Open* 2014;4:e004694.
50. Pang J, Hsu JP, Yeo TW, et al. Diabetes, cardiac disorders and asthma as risk factors for severe organ involvement among adult dengue patients: A matched case-control study. *Sci Rep* 2017;7:39872.
51. Baig Mirza AM, Fida M, Murtaza G, et al. Association of metabolic factors with dengue viral infection on admission triage which predict its clinical course during Lahore dengue epidemic. *J Pak Med Assoc* 2016;66:1102–1106.
52. Toledo J, George L, Martinez E, et al. Relevance of non-communicable comorbidities for the development of the severe forms of dengue: a systematic literature review. *PLoS Negl Trop Dis* 2016;10:e0004284.
53. Htun NS, Odermatt P, Eze IC, et al. Is diabetes a risk factor for a severe clinical presentation of dengue?—review and meta-analysis. *PLoS Negl Trop Dis* 2015;9:e0003741.
54. Jackson CA, Dobson AJ, Tooth LR, et al. Lifestyle and Socioeconomic determinants of multimorbidity patterns among mid-aged women: a longitudinal study. *PLoS One* 2016;11:e0156804.
55. Prados-Torres A, Poblador-Plou B, Calderón-Larrañaga A, et al. Multimorbidity patterns in primary care: interactions among chronic diseases using factor analysis. *PLoS One* 2012;7:e32190.
56. Starfield B, Shi L, Macinko J. Contribution of primary care to health systems and health. *Milbank Q* 2005;83:457–502.