

Appendix

Table 1 *Comparisons of Psychosocial Adjustment among Different Categorical Study*

Variables (n = 77).

Variables	Psychosocial adjustment				
	Mean	SD	t/F	p	Post Hoc #
Gender			0.543	0.588	
Male	412.61	32.43			
Female	423.00	33.96			
Education level			0.083	0.920	
Primary school and below	415.11	33.51			
Middle to high school	412.76	32.35			
College and above	409.63	33.21			
Marital status			1.75	0.083	
Single	424.56	38.65			
Married	409.49	29.61			
Employment status			3.50	0.001**	
No	420.38	31.44			
Yes	393.38	26.36			
Monthly income (NTD)			3.53	0.034*	①>②
①< 20,000	419.39	31.79			
②20,000 - 39,999	393.40	32.84			
③≥ 40,000	405.25	23.97			

Table 1 Comparisons of Psychosocial Adjustment among Different Categorical Study Variables (continued).

Variables	Psychosocial adjustment				
	Mean	SD	t/F	p	Post Hoc #
Financial hardship			9.95	<0.001***	①>②; ①>③
①Less than enough	427.97	31.79			
②Just enough	398.81	32.84			
③More than enough	396.71	23.9			
Hospital			2.36	0.101	
A	394.29	39.58			
B	407.13	25.51			
C	418.68	33.27			
Tumor site			-2.80	0.007**	
Buccal mucosa	407.26	30.49			
Others	429.18	31.91			
Cancer stage			1.64	0.188	
Stage I	405.46	38.55			
Stage II	412.86	23.53			
Stage III	398.82	37.69			
Stage IV	421.66	32.95			
Adjuvant therapy			0.62	0.537	
No	415.81	27.71			
Yes	411.13	35.26			

Note. t, value of independent t-test, F, value of one-way analyses of variance, ** p-value < .01, *** p-value < .001, # LSD Post Hoc test.

Table 2 *Person Product Moment Correlation Coefficients among the Continuous Study Variables (n = 77)*

Variables	Age	PD	FD	SS	PA
Age	1				
Post-surgical period (PD)	0.30**	1			
Facial disfigurement (FD)	0.04	0.07	1		
Social support (SS)	-0.17	-0.14	-0.21	1	
Psychosocial adjustment (PA)	0.06	-0.22*	0.33**	-0.40***	1

Note. * $p < 0.05$, ** p -value < 0.01 , *** p -value < 0.001