

# BMJ Open Compliance with pathology testing guidelines in Australian general practice: protocol for a secondary analysis of electronic health record data

Gorkem Sezgin,<sup>1</sup> Andrew Georgiou,<sup>1</sup> Rae-Anne Hardie,<sup>1</sup> Ling Li,<sup>1</sup> Lisa G Pont,<sup>1,2</sup> Tony Badrick,<sup>3</sup> Guilherme S Franco,<sup>1</sup> Johanna I Westbrook,<sup>1</sup> Natalie Rinehart,<sup>4</sup> Adam McLeod,<sup>4</sup> Christopher Pearce,<sup>4</sup> Marianne Shearer,<sup>5</sup> Robin Whyte,<sup>6</sup> Elizabeth Deveny<sup>7</sup>

**To cite:** Sezgin G, Georgiou A, Hardie R-A, *et al.* Compliance with pathology testing guidelines in Australian general practice: protocol for a secondary analysis of electronic health record data. *BMJ Open* 2018;**8**:e024223. doi:10.1136/bmjopen-2018-024223

► Prepublication history for this paper is available online. To view these files, please visit the journal online (<http://dx.doi.org/10.1136/bmjopen-2018-024223>).

Received 15 May 2018

Revised 26 July 2018

Accepted 20 September 2018



© Author(s) (or their employer(s)) 2018. Re-use permitted under CC BY-NC. No commercial re-use. See rights and permissions. Published by BMJ.

For numbered affiliations see end of article.

## Correspondence to

Gorkem Sezgin;  
gorkem.sezgin@mq.edu.au

## ABSTRACT

**Introduction** In Australia, general practitioners usually are the first point of contact for patients with non-urgent medical conditions. Appropriate and efficient utilisation of pathology tests by general practitioners forms a key part of diagnosis and monitoring. However overutilisation and underutilisation of pathology tests have been reported across several tests and conditions, despite evidence-based guidelines outlining best practice in pathology testing. There are a limited number of studies evaluating the impact of these guidelines on pathology testing in general practice. The aim of our quantitative observational study is to define how pathology tests are used in general practice and investigate how test ordering practices align with evidence-based pathology guidelines.

**Methods and analysis** Access to non-identifiable patient data will be obtained through electronic health records from general practices across three primary health networks in Victoria, Australia. Numbers and characteristics of patients, general practices, encounters, pathology tests and problems managed over time will be described. Overall rates of encounters and tests, alongside more detailed investigation between subcategories (encounter year, patient's age, gender, and location and general practice size), will also be undertaken. To evaluate how general practitioner test ordering coincides with evidence-based guidelines, five key candidate indicators will be investigated: full blood counts for patients on clozapine medication; international normalised ratio measurements for patients on warfarin medication; glycated haemoglobin testing for monitoring patients with diabetes; vitamin D testing; and thyroid function testing.

**Ethics and dissemination** Ethics clearance to collect data from general practice facilities has been obtained by the data provider from the RACGP National Research and Evaluation Ethics Committee (NREEC 17-008). Approval for the research group to use these data has been obtained from Macquarie University (5201700872). This study is funded by the Australian Government Department of Health Quality Use of Pathology Program (Agreement ID: 4-2QFVW4M). Findings will be reported to the Department of Health and disseminated in peer-reviewed academic journals and presentations (national and international conferences, industry forums).

## Strengths and limitations of this study

- The study population will be drawn from a large region in Victoria, Australia, and is expected to contain a large sample population (data from approximately 350 general practices), along with a large number of demographic variables (eg, region, gender, age).
- Electronic health records contain a vast amount of information on patients, allowing us to control for potential interacting or predictive variables on patient outcomes using statistical modelling.
- Some electronic health record fields may not be completely standardised across practices and could contain inconsistencies and missing information, which may limit the volume of data that can be extracted and analysed.
- Some medications being investigated in this study may be prescribed and/or monitored by specialists, which may result in cases being missed.

## INTRODUCTION

In Australia, general practitioners (GPs) are usually the first point of contact for patients with non-emergency health problems. They play an important role in the early detection, prevention (including shared-care arrangements) and treatment of disease.<sup>1</sup> Pathology tests are a key part of general practice assisting GPs in diagnosing, screening, treating and monitoring diseases. The past decade has seen an increase in both visits to and problems managed by GPs in Australia, resulting in an estimated 24.2 million additional pathology tests being ordered in 2015–2016 compared with 2006–2007.<sup>2</sup>

Appropriate and timely utilisation of pathology tests can improve the quality and outcome of patient care. However, both over-utilisation and underutilisation of pathology tests have been frequently observed in a

range of clinical scenarios.<sup>3,4</sup> Several guidelines have been established to encourage better utilisation of pathology tests among GPs across Australia, USA, Canada and the UK. These include initiatives such as *Choosing Wisely*<sup>5</sup>, *National Institute for Health and Clinical Excellence (NICE)*<sup>6</sup> and *Royal Australian College of General Practitioners (RACGP) Guidelines for preventative activities in general practice*.<sup>7</sup> Currently, there is little evidence on how test ordering practices by GPs align with these recommendations and guidelines in Australia, and elsewhere. A survey of 600 US doctors revealed only 21% of doctors were familiar with the *Choosing Wisely* campaign. However, doctors aware of the campaign had a lower proportion of unnecessary pathology testing.<sup>8</sup> In Australia, increased awareness of best-practice in vitamin D test ordering is reported to have contributed to a reduction in healthcare costs and potentially unnecessary tests.<sup>9</sup> Considering the importance of pathology testing for managing diseases, a better understanding of how pathology testing by GPs coincides with evidence-based guidelines will be invaluable for the success of management and disease-prevention strategies.

Until recently, the survey-based Bettering the Evaluation and Care of Health (BEACH) study provided the most comprehensive data on Australian GP activity.<sup>2</sup> However, its cross-sectional design prevented it from reporting longitudinal patient level changes and outcomes. The BEACH study was discontinued in 2016 and has since left a gap in our understanding of GP activity, particularly in relation to pathology test ordering. The extensive use of computers by Australian GPs has prompted interest in the use of electronic health record (EHR) data as a research source for monitoring the quality of GP services, and has led to research and publications based on EHR data.<sup>10</sup> The Australian Department of Health funded NPS Medicinewise MedicinesInsight data set contains a national collection of EHRs from 650 practices covering over 3300 GPs and nearly 3.6 million active patients.<sup>11</sup> This data set has been used in several population health research projects and has demonstrated the value of EHR data in research.<sup>12</sup> This study will use EHR data from the POLAR Data Space, containing de-identified data from consenting general practices collected on behalf of Australian primary health networks (PHNs) from approximately 350 practices. POLAR Data Space has added measures to ensure robust and accurate data, and implements standardised terminologies to make the data more approachable for research use. This study, undertaken in collaboration with POLAR Data Space Research Consortia and its associated PHNs, will facilitate a comprehensive analysis of general practice activity and its relationship to pathology testing in Australia through EHR data.

## Objectives

The objectives of this study are to:

1. Describe general practice activity and characteristics of pathology test ordering based on electronic health record data.

2. Investigate compliance with evidence-based guidelines to determine the appropriateness and quality use of pathology in general practice.

## METHODS AND ANALYSES

### Study design

A retrospective observational study of Australian general practice health records and pathology testing data. The study will run for a period of approximately 2 years spanning from early 2018 to late 2019.

### Data source

Data to be used in this study will be provided by the POLAR Data Space.<sup>10</sup> POLAR Data Space collects de-identified data from consenting general practices on behalf of Australian PHNs including Gippsland, Eastern Melbourne and South Eastern Melbourne PHNs. Data are extracted from approximately 350 practices in urban and rural regions in Victoria, Australia. The primary purpose of the data collection is to provide information to improve patient care at the practice level and population health initiatives at the PHN level. POLAR Data Space has ethics approval for the collection, storage and de-identification of the data, which it makes available for approved research governed by the involved PHNs.

The data source will include pooled general practice patient data extracted from Best Practice, Medical Director and Zedmed EHRs. Data will include de-identified demographic information about patients and general practices, as well as visit records (diagnosis, past history, medications) and pathology test records (test name and result). Both historical and current information about patients will be acquired, providing a longitudinal record. It is expected that the data will span a period of over a decade, from early 2000s to early 2018.

### Study population

The study population will consist of all patients who visited any of the general practices included in the study and had their visit recorded in the practices' EHR software.

### Variables

The following criteria will be adopted to accurately describe general practice activity and ensure data quality: (i) a patient will be distinguished by a unique (non-identifiable) patient code recorded within a general practice and included if determined to be an active patient (ie, has visited the GP three or more times in the past 2 years, not deceased); (ii) an encounter (ie, consultation, visit) will be defined as a patient visit recorded by a doctor or nurse during which an action (eg, consultation, prescription) is performed, and will be identified through a variable indicating the visit type (eg, surgery, administrative, phone call); (iii) a pathology test will be defined as either a panel of interrelated tests (eg, full blood count) or an individual test (eg, troponin test); (iv) where possible, standardised records such as Logical Observation Identifiers Names and Codes (LOINC) and Systematised

Nomenclature of Medicine (SNOMED) terminologies will be prioritised to identify variables; otherwise, free-text data will be searched for terms of interest (eg, 'diabetes' or its abbreviated forms in the diagnosis field, excluding 'not diabetes').

### Analyses

Data examination and analysis will be performed using Stata/MP V.15.1 (StataCorp). The methods outlined in this protocol are structured according to the Strengthening the Reporting of Observational Studies in Epidemiology checklist of items to be included in observational studies.<sup>13</sup>

### Characteristics of general practice activity and pathology testing

In line with objective 1 of this study, we will analyse and describe Australian general practice characteristics and activity. These characteristics can subsequently be compared with reports on national demographics and healthcare statistics, such as Australian Institute of Health and Welfare's annual reports.

Sample population characteristics will be reported. This will include describing the characteristics of patients, general practices, encounters, tests and problems managed across time. Subsequent analyses will describe overall median rates of encounters and tests, alongside more detailed investigation between subcategories (encounter year, patient's age, gender and postcode and number of active general practitioners in practice) using the following indicators: encounters per patient per year and tests per encounter.

The differences between subcategories will be further described as incidence rate ratios by generalised linear modelling (Poisson or negative binomial, whichever is appropriate).

### Best practice guidelines-based analyses

In line with objective 2 of this study, we will describe the extent to which pathology ordering practices among Australian GPs aligns with evidence-based pathology testing guidelines. Five initial candidate indicators have been identified for analyses.

### Monitoring patients on clozapine medication

#### Rationale

Clozapine is a highly effective antipsychotic drug that is used for managing chronic schizophrenia. However, clozapine use can result in neutropenia in nearly 2% of patients, and agranulocytosis in 1%<sup>14</sup>, warranting close monitoring of patients taking clozapine. Although clozapine is prescribed by specialists, monitoring is more frequently managed by GPs through shared-care arrangements.<sup>15</sup> After 18 weeks of initiation and monitoring under a specialist, GPs can also prescribe clozapine. Australian guidelines recommend patients on clozapine medication have blood tests for white blood cell and neutrophil counts weekly for the first 18 weeks of initiation, and monthly thereafter.<sup>16 17</sup> Furthermore, a patient

cannot obtain clozapine from the pharmacist without a recent blood test. Currently, the state of monitoring for patients on clozapine medication is not known.

#### Analysis

The study population will be patients who are being prescribed clozapine by the GP (and therefore also need to be monitored). The time frame will include records after the first entry of the prescription into the EHR software, and before medication is discontinued (or patient is deceased or is no longer an active patient). The demographic characteristics of patients on clozapine medication as well as number of full blood counts for these patients will be described overall, by patient gender, age and location, general practice size and year of test. The number of full blood counts per patient per year will also be described overall and by the demographic characteristics, by counting the number of full blood count tests conducted for each patient on clozapine medication for each available year. Subsequently, the median number of full blood count tests per patient per year will be calculated, with the inter-quartile range. For patients who undergo more than one test, the time between tests will be determined. Subsequently, median time between tests will be calculated, with the inter-quartile range. As clozapine may also be prescribed by specialists, it may not be possible to determine when the medication was initiated through the EHR software. Consequently, it may not be possible to differentiate patients who require weekly tests from patients who require monthly tests. Nonetheless, it is expected patients on clozapine medication will have at least one full blood count test within approximately 4 to 6 weeks of a prior test; which will indicate compliance with guidelines.

### Measuring international normalised ratio levels for patients on warfarin medication

#### Rationale

Warfarin is a highly effective and widely used anticoagulant in Australia. However, it is also one of the most common causes of prescribed medication-related mortality, due to its risk of causing bleeding.<sup>18</sup> Best practice guidelines recommend that the initiation of warfarin medication should be accompanied by frequent International Normalised Ratio (INR) measurements until a stable therapeutic range is reached. After INR levels are stable, testing frequency should be once every 4 to 6 weeks unless a change (eg, initiation of another medication) that can affect INR levels occurs.<sup>19 20</sup> Failure to correct INR levels is associated with increased mortality.<sup>21</sup>

#### Analysis

The sample population will be patients with a warfarin prescription. Only data entered after the first instance the prescription is recorded in the GP's computer and before the patient permanently stops the medication (or death) will be considered for analysis. As the pathology test frequency and repeat interval requirements for

patients treated with warfarin are similar to the requirements for clozapine, similar reporting standards will be used. As with clozapine medication, it may not be possible to differentiate patients who are initiating warfarin medication from those who have reached stable INR levels. Despite this, based on best practice guidelines, it is expected that patients on warfarin medication will have at least one INR measurement approximately within 4 weeks of a prior measurement, which will be the criteria used to determine compliance with guidelines.

### *Glycated haemoglobin testing for management of patients with diabetes*

#### Rationale

It is estimated that over one million Australians are diagnosed with diabetes, 85% of whom have type 2 diabetes.<sup>22</sup> Poor management of diabetes can lead to a range of complications, including cardiovascular and renal diseases and retinopathy.<sup>23</sup> Best practice guidelines recommend recurrent glycated haemoglobin (HbA1c) testing in at least half-yearly intervals for patients with diabetes.<sup>24</sup> HbA1c levels are a good indicator of long-term blood glucose control over the previous 8–12 weeks. Uncontrolled glucose leading to high HbA1c levels may indicate increased risk of diabetes-related complications. As such, undertesting may be associated with failure to identify complications.

#### Analysis

The sample population will be patients diagnosed with type 2 diabetes. Only data after the diagnosis being recorded and before death will be included. The analyses will be conducted and reported in a similar structure as outlined previously for clozapine and warfarin medication pathology testing guidelines. Based on the best practice guidelines, it is expected that a minimum of two HbA1c tests will be conducted in a year for patients diagnosed with diabetes.

### *Frequency of vitamin D testing*

#### Rationale

A study on vitamin D testing found a considerable number of potentially unnecessary vitamin D tests ordered by Australian GPs.<sup>25</sup> This has led to changes in funding programmes and the establishment of guidelines, suggesting only patients at risk of complications that may arise due to low vitamin D levels (eg, pregnant women, older patients at risk of falls, patients with osteoporosis) should be tested.<sup>26</sup> Ultimately, an overall reduction of vitamin D tests was observed<sup>9</sup>; however, studies on whether the tests are being ordered according to the guidelines are limited.<sup>27</sup>

#### Analysis

To understand if vitamin D tests are being ordered according to guidelines, population and demographic characteristics associated with higher vitamin D testing

will be investigated. The sample population will be all patients from which patients who had a vitamin D test will be identified and flagged (as the outcome variable). Univariate and multivariate logistic regression models will be fitted to identify any variation in vitamin D testing by the demographic characteristics of patients, diagnoses and general practices. Differences will be reported as ORs and their 95% confidence intervals. Descriptive characteristics (patient's gender, age, and location, general practice's size and year of test) of the sample population will also be described. To obtain an understanding of the reasons for vitamin D testing, other tests ordered simultaneously will be identified and examined, as well as preceding medication prescriptions and diagnoses.

### *Thyroid function tests*

#### Rationale

Thyroid dysfunction can occur due to overutilisation or underfunction of the thyroid gland, and can lead to cardiovascular diseases or subclinical hypothyroidism (remains asymptomatic).<sup>28</sup> Guidelines recommend assessing thyroid dysfunction by initial thyroid-stimulating hormone (TSH) tests, which may be followed up by free triiodothyronine (fT3) and free thyroxine (fT4) tests to assist with diagnosis if an abnormal TSH result is observed.<sup>29</sup> Otherwise, fT3 and fT4 tests are not recommended without a prior TSH test. In Australia, there are currently no screening guidelines for when and how frequently TSH tests should be conducted among adults. The benefit of screening for thyroid disease, or even treating subclinical hypothyroidism, remains uncertain.<sup>30</sup> It would be valuable to understand the demographics for and frequency of TSH test ordering by Australian GPs.

#### Analysis

To describe thyroid function test use by Australian GPs, patients with TSH, fT3 and fT4 tests will be identified from the data. The associations with population characteristics and diagnoses in TSH, fT3 and fT4 testing will be investigated and reported by univariate and multivariate logistic regression models, similar to the reporting of vitamin D testing outlined previously. In addition to the population and general practice demographics, the odds of fT3 and fT4 testing by prior TSH test result and normal/abnormal TSH test will also be analysed.

The descriptive characteristics for TSH, fT3 and fT4 tests will be described overall, and by patient's gender, age and location, general practice size and year of test. The number of fT3 and fT4 tests will be further described by TSH testing status: without a prior TSH test, simultaneously ordered with a TSH test and following a reported TSH test. Where TSH test results are available, fT3 and fT4 tests following TSH testing will be further described by whether the initial TSH test was normal or abnormal.

### *Sample size considerations*

The study will be based on a dynamic cohort, with the number of practices, GPs and patients expected to rise.

Current estimates suggest that the study will have data from 350 general practices. Therefore, it is expected that there will be ample scope to detect significant variation in practices across patient and general practice demographic domains.

### Patient and public involvement

There was no involvement of patients or the public in this study.

### DISCUSSION

Pathology tests play an important role in general practice. There are guidelines outlining best practice for utilising tests, although the role of these guidelines in decision making is not well established.

A limitation of using EHRs is that recording of clinical data is not always well-standardised, resulting in variation and inconsistencies in the information available.<sup>31</sup> Issues may arise due to free-text data with no standard formatting and absence of recorded comorbidities and diagnoses or missing data,<sup>32</sup> which are addressed by prioritising standardised terms and adopting stringent criteria to define variables. The POLAR program already codes and organises significant amounts of extracted data, and LOINC and SNOMED, both of which are available, provide standardised pathology tests and diagnoses. Another limitation of this study is that the study population will be drawn from only one region of Australia (Victoria), and the results may not be nationally representative. One other limitation is related to the indicators being measured. Medications, such as clozapine and warfarin, are generally prescribed by specialists who might also continually monitor the patient's status. In such cases, observed compliance with guidelines in general practice may be low. Furthermore, a patient monitored by a general practitioner may have occasional visits to their specialist, who may order the tests, rather than the GP.

This research will help define the extent to which evidence-based best practice guidelines influence decision making in general practices. To date, difficulties in obtaining patient data from EHR software have hindered studying pathology test ordering in general practice. This study will be one of the first in Australia to extensively investigate the impact of best practice guidelines on GP testing patterns. The study can ultimately lead to better efficiency in pathology testing and improvements in patient outcomes by providing much needed information on the adherence of GPs to pathology testing guidelines.

### Ethics and dissemination

The data will be de-identified and reported at an aggregate level, and the results will neither identify GPs nor patients.

The results of this study will be reported to the Australian Government's Department of Health, disseminated in peer-reviewed academic journals and presented in national and international conferences and industry

forums. The involvement of the PHNs in the research process also allows for the research findings to inform their activities at an early stage, reducing the usual 'research into practice' delay.

### Author affiliations

- <sup>1</sup>Centre for Health Systems and Safety Research, Australian Institute of Health Innovation, Macquarie University, Sydney, New South Wales, Australia
- <sup>2</sup>Graduate School of Health, University of Technology Sydney, Sydney, New South Wales, Australia
- <sup>3</sup>The Royal College of Pathologists of Australasia Quality Assurance Program, St Leonards, New South Wales, Australia
- <sup>4</sup>Outcome Health, Burwood, Victoria, Australia
- <sup>5</sup>Gippsland Primary Health Network, Moe, Victoria, Australia
- <sup>6</sup>Eastern Melbourne Primary Health Network, Box Hill, Victoria, Australia
- <sup>7</sup>South Eastern Melbourne Primary Health Network, Heatherton, Victoria, Australia

**Contributors** This study is jointly undertaken through a collaboration between Centre for Health Systems and Safety Research at Macquarie University, the Royal College of Pathologists of Australasia Quality Assurance Programs, POLAR Data Space (Outcome Health, Eastern Melbourne PHN, Gippsland PHN and South Eastern Melbourne PHN). AG initiated the project and led the development of the Quality Use of Pathology Program (QUPP) grant proposal. AG, JIW, LL, LGP, TB, CP and NR are chief investigators on the project and have contributed to the grant proposal and protocol in their area of expertise. From the PHNs, MS, RW and ED contributed their expertise in general practice to the protocol. AM, from Outcome Health, has contributed his expertise in EHR data to the protocol. R-AH, GS and GSF are members of the project team and contributed to the protocol in relation to describing the procedures of data collection, validation, and analyses procedures. R-AH and GS prepared the first draft of this protocol based on the grant proposal. All authors have reviewed and approved the final version of this protocol.

**Funding** This study is independently executed through funding by a grant from the Australian Government Department of Health, Quality Use of Pathology Program (Agreement ID: 4-2QFVW4M). This Australian national grants programme aims to improve health and economic outcomes from pathology tests.

**Competing interests** None declared.

**Patient consent** Not required.

**Ethics approval** Ethics clearance to collect data from general practice facilities has been obtained by the data provider from the RACGP National Research and Evaluation Ethics Committee (NREEC 17-008). Approval for the research group to use these data has been obtained from Macquarie University (5201700872).

**Provenance and peer review** Not commissioned; externally peer reviewed.

**Open access** This is an open access article distributed in accordance with the Creative Commons Attribution Non Commercial (CC BY-NC 4.0) license, which permits others to distribute, remix, adapt, build upon this work non-commercially, and license their derivative works on different terms, provided the original work is properly cited, appropriate credit is given, any changes made indicated, and the use is non-commercial. See: <http://creativecommons.org/licenses/by-nc/4.0/>.

### REFERENCES

1. Harris MF, Jayasinghe UW, Taggart JR, *et al*. Multidisciplinary team care arrangements in the management of patients with chronic disease in Australian general practice. *Med J Aust* 2011;194:236-9.
2. Britt H, Miller G, Bayram C, *et al*. A decade of Australian general practice activity 2006-07 to 2015-16: BEACH : bettering the evaluation and care of health. 2016.
3. Zhi M, Ding EL, Theisen-Toupal J, *et al*. The landscape of inappropriate laboratory testing: a 15-year meta-analysis. *PLoS One* 2013;8:e78962.
4. Elshaug AG, Watt AM, Mundy L, *et al*. Over 150 potentially low-value health care practices: an Australian study. *Med J Aust* 2012;197:556-60.
5. Bhatia RS, Levinson W, Shortt S, *et al*. Measuring the effect of Choosing Wisely: an integrated framework to assess campaign impact on low-value care. *BMJ Qual Saf* 2015;24:523-31.
6. National Institute for Health and Care Excellence. Clinical Guidelines. [www.nice.org.uk](http://www.nice.org.uk) (cited 26 Mar 2018).

7. RACGP. *Redbook Guidelines for preventive activities in general practice*. 9th edn. (cited 26 Mar 2018).
8. PerryUndem Research/Communication, 2014. What Physicians Say about the Problem, the Causes, and the Solutions: Results from a National Survey of Physicians. <http://www.choosingwisely.org/wp-content/uploads/2015/04/Final-Choosing-Wisely-Survey-report.pdf> (cited 26 Mar 2018).
9. Boyages SC. Vitamin D testing: new targeted guidelines stem the overtesting tide. *Med J Aust* 2016;204:18.
10. Pearce C, Shearer M, Gardner K, et al. A division's worth of data. *Aust Fam Physician* 2011;40:167–70.
11. MedicinesInsight. *MedicineInsight Data Book version 1.1*. Sydney: NPS MedicineWise. (cited 26 Mar 2018).
12. Gadzhanova S, Pratt N, Roughead E. Use of SGLT2 inhibitors for diabetes and risk of infection: Analysis using general practice records from the NPS MedicineWise MedicinesInsight program. *Diabetes Res Clin Pract* 2017;130:180–5.
13. Vandembroucke JP, von Elm E, Altman DG, et al. Strengthening the Reporting of Observational Studies in Epidemiology (STROBE): explanation and elaboration. *PLoS Med* 2007;4:e297.
14. Gerson SL. Clozapine—deciphering the risks. *N Engl J Med* 1993;329:204–5.
15. Filia SL, Wheelhouse A, Lee SJ, et al. Transitioning patients taking clozapine from the public to private/GP shared-care setting: barriers and criteria. *Aust N Z J Psychiatry* 2012;46:225–31.
16. Government of South Australia. *Clozapine Management Clinical Guideline*: Government of South Australia, 2017. (cited 26 Mar 2018).
17. State of Queensland, 2016. Safe and quality use of clozapine therapy in mental health services. Report No: QH-GDL-437 [https://www.health.qld.gov.au/\\_\\_data/assets/pdf\\_file/0025/625354/qh-gdl-437.pdf](https://www.health.qld.gov.au/__data/assets/pdf_file/0025/625354/qh-gdl-437.pdf)
18. Wysowski DK, Nourjah P, Swartz L. Bleeding complications with warfarin use: a prevalent adverse effect resulting in regulatory action. *Arch Intern Med* 2007;167:1414–9.
19. Tideman PA, Tirimacco R, St John A, et al. How to manage warfarin therapy. *Aust Prescr* 2015;38:44–8.
20. State of Queensland, 2016. Guidelines for warfarin management in the community;27 <https://www.health.qld.gov.au/clinical-practice/guidelines-procedures/medicines?a=165945>
21. Menzin J, Hoesche J, Friedman M, et al. Failure to correct International Normalized Ratio and mortality among patients with warfarin-related major bleeding: an analysis of electronic health records. *J Thromb Haemost* 2012;10:596–605.
22. Australian Government Department of Health. National Strategic Framework for Chronic Conditions. <http://www.health.gov.au/internet/main/publishing.nsf/content/nsfcc> (cited 26 Mar 2018).
23. Choby B. Diabetes Update: Prevention and Management of Diabetes Complications. *FP Essent* 2017;456:36–40.
24. The Royal Australian College of General Practitioners (RACGP), 2018. General practice management of type 2 diabetes. <https://www.racgp.org.au/your-practice/guidelines/diabetes/> (cited 26 Mar 2018).
25. Bilinski K, Boyages S. Evidence of overtesting for vitamin D in Australia: an analysis of 4.5 years of Medicare Benefits Schedule (MBS) data. *BMJ Open* 2013;3:e002955.
26. The Royal College of Pathologists of Australia (RCPA), 2013. Position Statement: Use and Interpretation of Vitamin D testing. <https://www.rcpa.edu.au/Library/College-Policies/Position-Statements/Use-and-Interpretation-of-Vitamin-D-Testing> (cited 25 Mar 2018).
27. Gowda U, Smith BJ, Wluka AE, et al. Vitamin D testing patterns among general practitioners in a major Victorian primary health care service. *Aust N Z J Public Health* 2016;40:144–7.
28. Surks MI, Ortiz E, Daniels GH, et al. Subclinical thyroid disease: scientific review and guidelines for diagnosis and management. *JAMA* 2004;291:228–38.
29. Walsh JP. Managing thyroid disease in general practice. *Med J Aust* 2016;205:179–84.
30. Ladenson PW, Singer PA, Ain KB, et al. American Thyroid Association guidelines for detection of thyroid dysfunction. *Arch Intern Med* 2000;160:1573–5.
31. Liaw ST, Taggart J, Yu H, et al. Data extraction from electronic health records - existing tools may be unreliable and potentially unsafe. *Aust Fam Physician* 2013;42:820–3.
32. Young J, Eley D, Fahey P, et al. Enabling research in general practice—increasing functionality of electronic medical records. *Aust Fam Physician* 2010;39:506–9.