BMJ Open Relationship between hospital or surgeon volume and outcomes in joint arthroplasty: protocol for a suite of systematic reviews and dose-response meta-analyses

Xiang-Dong Wu,^{1,2} Meng-Meng Liu,³ Ya-Ying Sun,⁴ Zhi-Hu Zhao,⁵ Quan Zhou,⁶ Joey S W Kwong,^{7,8} Wei Xu,¹ Mian Tian,^{1,9} Yao He,^{1,10} Wei Huang¹

To cite: Wu X-D, Liu M-M, Sun Y-Y, *et al.* Relationship between hospital or surgeon volume and outcomes in joint arthroplasty: protocol for a suite of systematic reviews and dose-response meta-analyses. *BMJ Open* 2018;**8**:e022797. doi:10.1136/bmjopen-2018-022797

► Prepublication history for this paper is available online. To view these files, please visit the journal online (<http://dx.doi.org/10.1136/bmjopen-2018-022797>).

Received 6 March 2018
Revised 11 September 2018
Accepted 7 November 2018

© Author(s) (or their employer(s)) 2018. Re-use permitted under CC BY-NC. No commercial re-use. See rights and permissions. Published by BMJ.

For numbered affiliations see end of article.

Correspondence to

Dr Xiang-Dong Wu;
xiangdong_wu@outlook.com,
xiangdong_wu@foxmail.com
and Professor Wei Huang;
drhuangwei68@gmail.com,
huangwei68@263.net

ABSTRACT

Introduction Joint arthroplasty is a particularly complex orthopaedic surgical procedure performed on joints, including the hip, knee, shoulder, ankle, elbow, wrist and even digit joints. Increasing evidence from volume-outcomes research supports the finding that patients undergoing joint arthroplasty in high-volume hospitals or by high-volume surgeons achieve better outcomes, and minimum case load requirements have been established in some areas. However, the relationships between hospital/surgeon volume and outcomes in patients undergoing arthroplasty are not fully understood. Furthermore, whether elective arthroplasty should be restricted to high-volume hospitals or surgeons remains in dispute, and little is known regarding where the thresholds should be set for different types of joint arthroplasties.

Methods and analyses This is a protocol for a suite of systematic reviews and dose-response meta-analyses, which will be amended and updated in conjunction with the Preferred Reporting Items for Systematic Reviews and Meta-Analyses Protocols. Electronic databases, including PubMed and Embase, will be searched for observational studies examining the relationship between the hospital or surgeon volume and clinical outcomes in adult patients undergoing primary or revision of joint arthroplasty. We will use records management software for study selection and a predefined standardised file for data extraction and management. Quality will be assessed using the Newcastle-Ottawa Scale, and the meta-analysis, subgroup analysis and sensitivity analysis will be performed using Stata statistical software. Once the volume-outcome relationships are established, we will examine the potential non-linear relationships between hospital/surgeon volume and outcomes and detect whether thresholds or turning points exist.

Ethics and dissemination Ethical approval is not required, because these studies are based on aggregated published data. The results of this suite of systematic reviews and meta-analyses will be submitted to peer-reviewed journals for publication.

PROSPERO registration number CRD42017056639.

Strengths and limitations of this study

- To the best of our knowledge, this will be the first suite of systematic reviews and dose-response meta-analyses to explore the relationship between the hospital/surgeon volume and outcomes of the dominant types of arthroplasties.
- Once the volume-outcome relationships are established, we will examine the potential non-linear relationships between hospital/surgeon volume and outcomes, and determine whether thresholds or turning points exist.
- A comprehensive literature search, developed in consultation with a librarian with experience in systematic review search strategies, will be performed to include all eligible studies to present comprehensive systematic reviews and meta-analyses of the currently available evidence.
- It is highly likely that many studies will be included in the systematic reviews but excluded from the meta-analyses due to a paucity of data, which will introduce some bias.

INTRODUCTION

In previous decades, the relationships between the number of patients treated by a physician/surgeon (physician/surgeon volume) or in a hospital (hospital volume) and patient outcomes have been extensively documented under many medical and surgical conditions.^{1–14} In these relationships, patients admitted to a higher volume hospital or treated by a higher volume physician/surgeon are thought to be associated with a lower rate of adverse events and better health outcomes.³ And a posteriori defined volume threshold for such an association usually is artificially determined to optimise the correlation when the volume-outcome relationship has been established, indicating that there is a hospital/surgeon volume above

which any increase will no longer be associated with improved outcomes. However, controversy pertaining to such volume–outcome relationships persists, and recent published studies have conveyed inconsistent results, thus fuelling a continuing debate.²

Supporters believe that volume–outcome relationship may be a sensible surrogate for quality assessment when choosing where to obtain surgical and interventional care,¹⁵ while opponents argue that such volume–outcome relationship is an imperfect indicator of healthcare quality and that generalisability of these results is uncertain.^{16 17} Despite the heated controversy that it provokes, the volume–outcome relationship remains a good point of departure in the exploration of optimal care in health services delivery. Understanding these relationships remains critical for clinicians and policy-makers because they are under increasing pressure to elucidate the performances of hospitals/surgeons, analyse the processes of care that lead to optimal outcomes and identify strategies to improve the quality of care.^{18 19} Considering that volume is not an immutable determinant of the incidence of adverse events and that the volume–outcome relationship can change with the development of the healthcare provider and improvements in the quality of care, updating these volume–outcome relationships should be a priority when new research released.

Joint arthroplasty or joint replacement surgery is a particularly complex orthopaedic surgical procedure that is performed when severe joint pain or dysfunction cannot be alleviated by less invasive therapies. The goal of this procedure is to relieve pain, restore joint function and enhance quality of life.^{20–23} Until now, joint arthroplasty has been performed on joints including the hip, knee, shoulder, ankle, elbow, wrist and even digit joints.

Hip and knee arthroplasties are the most common types of procedures performed.²⁴ According to estimates for the USA, the demands for primary total hip arthroplasty (THA) and revision THA will reach 572 000 and 96 700, respectively, by 2030, while the demands for primary total knee arthroplasty (TKA) and revision TKA will reach 3.48 million and 268 000, respectively.^{25 26} Although shoulder arthroplasty is less common than hip and knee arthroplasties, it is still an exceptional procedure with excellent results,²⁷ and more than 53 000 shoulder arthroplasties are performed each year in the USA.²⁸ Because ankle arthrodesis has long been considered the gold standard of surgical treatment for ankle arthritis, total ankle arthroplasty is not frequently performed.²⁹ Only 2608 procedures were performed in the USA in 2010.³⁰ The elbow joint cannot be easily replaced or bypassed by external aids as can the lower extremity joints, and total elbow arthroplasty remains a relatively uncommon surgical procedure.^{31 32} Approximately 3000 procedures were performed in the USA in 2015.³³ Similar to the ankle joint, wrist arthrodesis is the most frequently recommended treatment,³⁴ with only approximately 1000 total wrist arthroplasty procedures performed annually in the USA.³⁵

Increasing evidence from volume–outcome research supports the finding that patients undergoing arthroplasty in high-volume hospitals or by high-volume surgeons achieve better outcomes,^{36–49} but the actual definitions of high-volume hospitals and surgeons are highly variable among studies.⁵⁰ To improve clinical outcomes and deliver the best healthcare, the German Federal Joint Committee has established minimum case load requirements. The volume standards for primary TKA proposed by the Committee are 25 TKAs per year for surgeons and 50 TKAs per year for hospitals.⁵¹ Although accumulating evidence supports these interventions, many researchers question how the minimum case load requirements should be exactly determined in clinical practice.^{50 52}

The relationships between the hospital/surgeon volume and the outcomes in patients undergoing arthroplasty are not fully understood; whether elective arthroplasty should be restricted to high-volume hospitals or surgeons remains in dispute; and little is known regarding where exactly the thresholds should fall for different types of joint arthroplasties. Therefore, we decided to explore the volume–outcome relationships and thresholds to address this issue.

OBJECTIVE

This is a protocol for a suite of systematic reviews and dose–response meta-analyses to explore the relationship between hospital/surgeon volume and outcomes in patients undergoing arthroplasty with the following objectives:

1. To examine the relationships between hospital/surgeon volume and outcomes in different types of joint arthroplasties.
2. To investigate the dose–response relationship between the volume and outcomes and to propose meaningful hospital/surgeon arthroplasty volume thresholds.
3. To compare the volume–outcome relationships among different procedure volumes for joint arthroplasties (primary hip and knee arthroplasties *vs* revision hip and knee arthroplasties *vs* shoulder arthroplasty *vs* ankle, elbow and wrist arthroplasties).

METHODS AND ANALYSIS

Our systematic reviews and meta-analyses will be performed in accordance with guidelines from the Meta-analysis Of Observational Studies in Epidemiology (MOOSE) group and the methods prescribed in the Cochrane Handbook for Systematic Reviews of Interventions and will be reported following the Preferred Reporting Items for Systematic Reviews and Meta-Analyses (PRISMA) Statement.^{53 54}

Registration information

This protocol will be amended and updated in conjunction with the PRISMA-Protocols.⁵⁵

Eligibility criteria

Types of studies

We will include observational studies that examined the relationship between hospital or surgeon volume and clinical outcomes, mainly including prospective cohort studies, retrospective cohort studies, case-control studies and cross-sectional studies, and meeting abstracts will also be included if eligible. There will be no restrictions regarding publication status or language.

Types of participants

Adult patients undergoing primary or revision joint arthroplasty will be eligible, and will specifically include those undergoing primary and revision THA, primary and revision TKA, and primary shoulder, ankle, elbow or wrist arthroplasty.

Types of outcome measures

The outcomes of interest are as follows: rate of mortality, readmission, periprosthetic joint infection, dislocation, revision, as well as wound complication, urinary tract infection, length of hospital stay, hospitalisation cost and functional score. The list is not exhaustive and will be modified based on the evidence compiled from the systematic reviews. We will not exclude studies due to paucity of data and we will include these studies in systematic reviews.

Patient and public involvement

In this study, data will not be collected directly from patients but instead will be obtained from published studies available in the main databases. Therefore, patients will not be involved in the completion of the systematic review protocol or subsequent research.

Information sources

We will search the electronic bibliographic databases PubMed and Embase from inception to March 2018 to ensure that all recent relevant studies are captured. No language restrictions will be imposed. In addition, we will search the clinical trial registry for ongoing and unpublished studies. Reference lists of all the identified studies as well as relevant reviews will be manually searched for potentially relevant studies. Potential grey literature sources (eg, conference abstracts) will be screened to identify any eligible published and unpublished studies.

Search strategy

Electronic search terms for each part will include both exploded Medical Subject Headings (MeSHs) terms and corresponding keywords. Search terms will include those related to 'Volume', 'Caseload', 'Arthroplasty', 'Replacement' and their variants. The search will be broad, and no restrictions will be applied. After retrieving and combining the corresponding subject terms using 'OR', the two parts will be combined using 'AND'. The detailed librarian-assisted search strategy is shown in [table 1](#).

Table 1 Search strategy

PubMed	Search	Query
#1		'Arthroplasty'[Mesh]
#2		Arthroplasty[Title/Abstract]
#3		Replacement[Title/Abstract]
#4		'Arthroplasty, Replacement'[Mesh]
#5		#1 OR #2 OR #3 OR #4
#6		'Hospitals, High-Volume'[Mesh]
#7		'Hospitals, Low-Volume'[Mesh]
#8		Volume[Title/Abstract]
#9		Volumes[Title/Abstract]
#10		Caseload[Title/Abstract]
#11		Caseloads[Title/Abstract]
#12		#6 OR #7 OR #8 OR #9 OR #10 OR #11
#13		hemofiltration[Title/Abstract]
#14		hematoma[Title/Abstract]
#15		Brain[Title/Abstract]
#16		Blood[Title/Abstract]
#17		Search Platelet[Title/Abstract]
#18		Gastric[Title/Abstract]
#19		Ventricle[Title/Abstract]
#20		Pressure[Title/Abstract]
#21		Lung[Title/Abstract]
#22		Stroke[Title/Abstract]
#23		hemodialysis[Title/Abstract]
#24		Tidal[Title/Abstract]
#25		#12 NOT #13 NOT #14 NOT #15 NOT #16 NOT #17 NOT #18 NOT #19 NOT #20 NOT #21 NOT #22 NOT #23 NOT #24
#26		#12 AND #25
Embase	Search	Query
#1		arthroplasty'/exp
#2		replacement arthroplasty'/exp
#3		arthroplasty:ti,ab
#4		replacement:ti,ab
#5		#1 OR #2 OR #3 OR #4
#6		volume'/exp
#7		volume:ti,ab
#8		volumes:ti,ab
#9		caseload:ti,ab
#10		caseloads:ti,ab
#11		#6 OR #7 OR #8 OR #9 OR #10
#12		hemofiltration:ti,ab
#13		hematoma:ti,ab
#14		brain:ti,ab
#15		blood:ti,ab
#16		platelet:ti,ab
#17		gastric:ti,ab
#18		ventricle:ti,ab

Continued

Table 1 Continued

Embase	Search	Query
#19		pressure:ti,ab
#20		lung:ti,ab
#21		stroke:ti,ab
#22		hemodialysis:ti,ab
#23		tidal:ti,ab
#24		#12 OR #13 OR #14 OR #15 OR #16 OR #17 OR #18 OR #19 OR #20 OR #21 OR #22 OR #23
#25		#11 NOT #24
#26		#5 AND #25

Data collection and synthesis

Study selection

The obtained study records will be exported from medical databases and imported into a software package (EndNote V.X7, Thomson Reuters, California, USA) for records management. We will use a three-stage process for study screening and selection using standardised and piloted screening forms (figure 1). First, two reviewers

(X-DW and YYS) will jointly remove duplicate records from the initial search results. Second, the two reviewers will independently screen the titles and abstracts of each record to determine the eligibility, and identify the studies as included, excluded or requiring further assessment. Third, the full text of potentially eligible records will be retrieved and reviewed independently with reference to the predetermined inclusion and exclusion criteria. Differences of opinions will be resolved by discussion and consensus with a third reviewer (WH).

Data extraction and management

A predefined standardised Excel (Microsoft Corporation, Washington, USA) file will be applied for data extraction, and separate sheets will be applied for each type of arthroplasty. Two independent reviewers (Z-HZ and YH) will extract the following information from each included study: first author, publication year, study location, study design, database, study period, number of patients, volume grouping and category, multivariate effect estimate, covariates in the fully adjusted model, as well as outcome measures mentioned above. The online supplementary files of the included studies will

Figure 1 PRISMA flow diagram showing the process of literature screening, study selection and reasons for study exclusion. The PRISMA statement is used worldwide to improve the reporting of systematic reviews and meta-analyses. PRISMA, Preferred Reporting Items for Systematic Reviews and Meta-Analyses.

also be examined for data extraction. In cases of missing data, we will contact the authors of the study. If we fail to obtain the missing data, the study will not be included in data analysis. Any discrepancies between the two reviewers will be resolved through consensus by discussion with an independent adjudicator (WH) as required.

Quality assessment

As recommended by the MOOSE checklist, the quality of the included studies will be assessed using the Newcastle-Ottawa Scale, which is a validated scale for evaluating the quality of non-randomised studies in meta-analyses. This scale contains eight items with a maximum score of nine stars, which are awarded based on three domains: four stars for selection, two stars for comparability and three stars for outcomes. We will assign studies with scores of 0–3, 4–6 and 7–9 as low, moderate and high-quality studies, respectively. Two authors (MT and WX) will independently perform the quality appraisal, and disagreements will be resolved by a third investigator (WH).

Data synthesis

Included studies that provide sufficient data to calculate an effect size measure will be included in the quantitative analysis. To quantify the degree of heterogeneity across studies, we will use Cochran's Q test with its p value and the Higgins I^2 statistic with its 95% CI.^{56 57} The I^2 statistic is used to quantify the proportion of total variation in the effect estimation that is due to between-study variation. I^2 values of 25%, 50% and 75% will be used as indicators of low, moderate and high heterogeneity, respectively.^{57 58} Multivariate ORs with corresponding 95% CIs between extreme levels of hospital/surgeon volume (highest *vs* lowest) will be pooled using a random-effects model accounting for clinical heterogeneity. We will evaluate the possible presence of publication bias by using a funnel plot for meta-analyses including at least 10 studies.⁵⁹ We will also use tests proposed by Egger *et al.*, and by Begg and Mazumdar to measure funnel plot asymmetry.^{60 61}

Subgroup analysis and sensitivity analysis

To explore the potential sources of heterogeneity among the studies and to test the robustness of the volume–outcome relationships, we will further carry out subgroup analyses, primarily including the study design (cohort studies *vs* cross-sectional studies), adjusted factors (uncertain), sample size (uncertain) and period (?–1998 *vs* 1999–2008 *vs* 2009–2018). Additional 'leave-one-out' sensitivity analyses will be performed to explore whether the results are dominated by a single study; this issue will be investigated by omitting each study in turn and examining the influence of each individual study on the overall risk estimate (the 'leave-one-out' approach). This approach will enable an evaluation of the influence of individual studies on the overall risk estimate, and a two-sided $p < 0.05$ will be considered as statistically significant. All of the above analyses will be performed using the Stata statistical software V.13.0 (StataCorp).

Dose–response analysis and threshold effect analysis

Once the volume–outcome relationships are established, we will use two-step random-effects meta-regression models to examine potential non-linear relationships between hospital/surgeon volume and outcomes. To derive the dose–response curve, study-specific slopes (non-linear trends) with 95% CIs from the natural logs of the reported ORs and CIs across the hospital/surgeon volume categories will be calculated. The details of the methods that will be used have been described by Greenland and Longnecker and Orsini *et al.*^{62 63} In particular, the mean or median level of volume for each category of the annual hospital/surgeon volume will be assigned to each corresponding OR for each study. If the data are not available, we will assign the midpoint of the upper and lower boundaries in each category as the annual hospital/surgeon volume. In cases where the upper or lower boundary of the category is open ended or extreme upper or lower values are present, we will assume that the absent boundary has the same amplitude as the adjacent category, meaning that the highest boundary has the same amplitude as the closest category, and the lowest boundary will be assumed to be 0. Additionally, only studies that report the number of events and control subjects (rather than the event rate), and the OR and its variance estimate for at least three categories will be eligible for the dose–response analysis.

For different types of arthroplasties, if the dose–response relationships are available, we will further develop a two-piecewise linear regression model to detect whether there exist thresholds or turning points of the annual hospital/surgeon volume on outcome using a smoothing function. The threshold level will be determined using trial and error, primarily by including the selection of turning points along a predefined interval and choosing the turning point that yields the maximum model likelihood. All analyses will be performed using Empower (R) (www.empowerstats.com, X&Y solutions, Massachusetts, USA).

DISCUSSION

The relationship between hospital/surgeon volume and clinical outcomes has been proposed in arthroplasty for more than two decades.³⁹ Currently, minimum case load requirements or certificate-of-need programmes have been implemented to improve healthcare quality or to prevent overutilisation of healthcare resources. These measures are intended to align the supply of facilities with demand, but the advantages and disadvantages of regionalisation or centralisation in joint arthroplasty have not been fully elaborated. Therefore, this suite of systematic reviews and meta-analyses will provide new knowledge that is essential for healthcare service planning as well as knowledge on the implementation process and adjustment of programmes to improve the quality of care.

We anticipate challenges in conducting this research. First, it is highly likely that many studies will be included

in the systematic reviews but excluded from the meta-analyses due to a paucity of data, which will introduce some bias. Second, most of the included studies will be retrospective, which will limit the ability to control for confounders. Additionally, the numbers of eligible studies for ankle, elbow and wrist arthroplasties are predicted to be small, and the findings of the meta-analyses may be restricted.

ETHICS AND DISSEMINATION

The findings will be disseminated in peer-reviewed journals and will also be shared with all stakeholders. Knowledge dissemination workshops will be conducted with relevant stakeholders to transfer the evidence, which will be tailored to the stakeholder (eg, policy briefs, publications and information booklets).

Author affiliations

¹Department of Orthopaedic Surgery, The First Affiliated Hospital of Chongqing Medical University, Chongqing, China

²Evidence-Based Perioperative Medicine 07 Collaboration Group, China

³Department of Pathology, Anhui Medical University, Hefei, China

⁴Department of Sports Medicine, Huashan Hospital, Fudan University, Shanghai, China

⁵Department of orthopaedic, Tianjin Hospital, Tianjin, China

⁶Department of Science and Education, First People's Hospital of Changde City, Changde, China

⁷Department of Health Policy, National Center for Child Health and Development, Tokyo, Japan

⁸Department of Clinical Epidemiology, National Center for Child Health and Development, Tokyo, Japan

⁹Department of Orthopaedic Surgery, Dianjiang People's Hospital, Chongqing, China

¹⁰Department of Orthopaedic Surgery, Banan People's Hospital of Chongqing, Chongqing, China

Contributors X-DW and WH conceived and designed the study. X-DW drafted the protocol. M-ML, Y-YS, Z-HZ, QZ, JSWK, WX, MT, YH and WH revised the protocol. X-DW and Y-YS will search for and select eligible studies. Z-HZ and MT will extract the data. M-ML will check the data. YH and WX will assess the risk of bias. QZ and JSWK will perform the data synthesis. X-DW and WH act as guarantors of the protocol. All authors approved the publication of the protocol.

Funding The authors have not declared a specific grant for this research from any funding agency in the public, commercial or not-for-profit sectors.

Competing interests None declared.

Patient consent Not required.

Ethics approval Because this suite of systematic reviews and dose-response meta-analyses involves analysis of anonymous secondary data that are available in the public domain, and does not involve human participants or encroach on the privacy of individual patients, ethical approval is not required.

Provenance and peer review Not commissioned; externally peer reviewed.

Open access This is an open access article distributed in accordance with the Creative Commons Attribution Non Commercial (CC BY-NC 4.0) license, which permits others to distribute, remix, adapt, build upon this work non-commercially, and license their derivative works on different terms, provided the original work is properly cited, appropriate credit is given, any changes made indicated, and the use is non-commercial. See: <http://creativecommons.org/licenses/by-nc/4.0/>.

REFERENCES

- Luft HS, Bunker JP, Enthoven AC. Should operations be regionalized? The empirical relation between surgical volume and mortality. *N Engl J Med* 1979;301:1364-9.
- Epstein AM. Volume and outcome--it is time to move ahead. *N Engl J Med* 2002;346:1161-4.
- Halm EA, Lee C, Chassin MR. Is volume related to outcome in health care? A systematic review and methodologic critique of the literature. *Ann Intern Med* 2002;137:511-20.
- Post PN, Kuijpers M, Ebels T, et al. The relation between volume and outcome of coronary interventions: a systematic review and meta-analysis. *Eur Heart J* 2010;31:1985-92.
- Schelbert EB, Vaughan-Sarrazin MS, Welke KF, et al. Hospital volume and selection of valve type in older patients undergoing aortic valve replacement surgery in the United States. *Circulation* 2005;111:2178-82.
- Holt PJ, Poloniecki JD, Gerrard D, et al. Meta-analysis and systematic review of the relationship between volume and outcome in abdominal aortic aneurysm surgery. *Br J Surg* 2007;94:395-403.
- Nguyen Y-L, Wallace DJ, Yordanov Y, et al. The volume-outcome relationship in critical care. *Chest* 2015;148:79-92.
- Gu WJ, Wu XD, Zhou Q, et al. Relationship between Annualized Case Volume and Mortality in Sepsis: A Dose-Response Meta-analysis. *Anesthesiology* 2016;125:168-79.
- Brusselaers N, Mattsson F, Lagergren J. Hospital and surgeon volume in relation to long-term survival after oesophagectomy: systematic review and meta-analysis. *Gut* 2014;63:1393-400.
- Eskander A, Merdad M, Irish JC, et al. Volume-outcome associations in head and neck cancer treatment: a systematic review and meta-analysis. *Head Neck* 2014;36:1820-34.
- Gooiker GA, van Gijn W, Wouters MW, et al. Systematic review and meta-analysis of the volume-outcome relationship in pancreatic resection. *Br J Surg* 2011;98:485-94.
- Goossens-Laan CA, Gooiker GA, van Gijn W, et al. A systematic review and meta-analysis of the relationship between hospital/surgeon volume and outcome for radical cystectomy: an update for the ongoing debate. *Eur Urol* 2011;59:775-83.
- Gooiker GA, van Gijn W, Post PN, et al. A systematic review and meta-analysis of the volume-outcome relationship in the surgical treatment of breast cancer. Are breast cancer patients better off with a high volume provider? *Eur J Surg Oncol* 2010;36:S27-35.
- Markar SR, Karthikesalingam A, Thrumurthy S, et al. Volume-outcome relationship in surgery for esophageal malignancy: systematic review and meta-analysis 2000-2011. *J Gastrointest Surg* 2012;16:1055-63.
- Ross JS, Normand SL, Wang Y, et al. Hospital volume and 30-day mortality for three common medical conditions. *N Engl J Med* 2010;362:1110-8.
- Ghali WA, Ash AS, Hall RE, et al. Statewide quality improvement initiatives and mortality after cardiac surgery. *JAMA* 1997;277:379-82.
- Birkmeyer JD, Siewers AE, Finlayson EV, et al. Hospital volume and surgical mortality in the United States. *N Engl J Med* 2002;346:1128-37.
- Silber JH, Rosenbaum PR, Brachet TJ, et al. The Hospital Compare mortality model and the volume-outcome relationship. *Health Serv Res* 2010;45(5 Pt 1):1148-67.
- Auerbach AD, Hilton JF, Maselli J, et al. Follow the crowd or shop for the best?: Volume, quality, and outcomes of coronary artery bypass surgery. *Ann Intern Med* 2009;150:696-704.
- Skou ST, Roos EM, Laursen MB, et al. A randomized, controlled trial of total knee replacement. *N Engl J Med* 2015;373:1597-606.
- Jacobs JJ, Roebuck KA, Archibeck M, et al. Osteolysis: basic science. *Clin Orthop Relat Res* 2001;393:71-7.
- Pivec R, Johnson AJ, Mears SC, et al. Hip arthroplasty. *Lancet* 2012;380:1768-77.
- Carr AJ, Robertsson O, Graves S, et al. Knee replacement. *Lancet* 2012;379:1331-40.
- Maradit Kremers H, Larson DR, Crowson CS, et al. Prevalence of total hip and knee replacement in the United States. *J Bone Joint Surg Am* 2015;97:1386-97.
- Kurtz S, Ong K, Lau E, et al. Projections of primary and revision hip and knee arthroplasty in the United States from 2005 to 2030. *J Bone Joint Surg Am* 2007;89:780-5.
- Kurtz SM, Lau E, Ong K, et al. Future young patient demand for primary and revision joint replacement: national projections from 2010 to 2030. *Clin Orthop Relat Res* 2009;467:2606-12.
- Kim SH, Wise BL, Zhang Y, et al. Increasing incidence of shoulder arthroplasty in the United States. *J Bone Joint Surg Am* 2011;93:2249-54.
- American Academy of Orthopaedic Surgeons. G. M. Shoulder joint replacement- OrthoInfo- AAOS. <https://orthoinfo.aaos.org/en/treatment/shoulder-joint-replacement> (accessed 05 Sep 2018).

29. Raikin SM, Rasouli MR, Espandar R, *et al.* Trends in treatment of advanced ankle arthropathy by total ankle replacement or ankle fusion. *Foot Ankle Int* 2014;35:216–24.
30. Singh JA, Ramachandran R. Time trends in total ankle arthroplasty in the USA: a study of the National Inpatient Sample. *Clin Rheumatol* 2016;35:239–45.
31. de Putter CE, Selles RW, Haagsma JA, *et al.* Health-related quality of life after upper extremity injuries and predictors for suboptimal outcome. *Injury* 2014;45:1752–8.
32. Triplett JJ, Kurowicki J, Momoh E, *et al.* Trends in total elbow arthroplasty in the Medicare population: a nationwide study of records from 2005 to 2012. *J Shoulder Elbow Surg* 2016;25:1848–53.
33. Day JS, Lau E, Ong KL, *et al.* Prevalence and projections of total shoulder and elbow arthroplasty in the United States to 2015. *J Shoulder Elbow Surg* 2010;19:1115–20.
34. Hinds RM, Capo JT, Rizzo M, *et al.* Total Wrist Arthroplasty Versus Wrist Fusion: Utilization and Complication Rates as Reported by ABOS Part II Candidates. *Hand* 2017;12:376–81.
35. Melamed E, Marascalchi B, Hinds RM, *et al.* Trends in the utilization of total wrist arthroplasty versus wrist fusion for treatment of advanced wrist arthritis. *J Wrist Surg* 2016;5:211–6.
36. Ravi B, Jenkinson R, Austin PC, *et al.* Relation between surgeon volume and risk of complications after total hip arthroplasty: propensity score matched cohort study. *BMJ* 2014;348:g3284.
37. Ries MD. CORR Insights®: Higher Volume Surgeons Have Lower Medicare Payments, Readmissions, and Mortality After THA. *Clin Orthop Relat Res* 2018.
38. Koltsov JCB, Marx RG, Bachner E, *et al.* and Surgeon-Volume Categories for Total Hip Arthroplasty. *J Bone Joint Surg Am* 2018;100:1203–8.
39. Kreder HJ, Deyo RA, Koepsell T, *et al.* Relationship between the volume of total hip replacements performed by providers and the rates of postoperative complications in the state of Washington. *J Bone Joint Surg Am* 1997;79:485–94.
40. Katz JN, Losina E, Barrett J, *et al.* Association between hospital and surgeon procedure volume and outcomes of total hip replacement in the United States medicare population. *J Bone Joint Surg Am* 2001;83-A:1622–9.
41. Hervey SL, Purves HR, Guller U, *et al.* Provider volume of total knee arthroplasties and patient outcomes in the HCUP-Nationwide Inpatient Sample. *J Bone Joint Surg Am* 2003;85-A:1775–83.
42. Katz JN, Barrett J, Mahomed NN, *et al.* Association between hospital and surgeon procedure volume and the outcomes of total knee replacement. *J Bone Joint Surg Am* 2004;86-A:1909–16.
43. Doro C, Dimick J, Wainess R, *et al.* Hospital volume and inpatient mortality outcomes of total hip arthroplasty in the United States. *J Arthroplasty* 2006;21:10–16.
44. Ames JB, Lurie JD, Tomek IM, *et al.* Does surgeon volume for total hip arthroplasty affect outcomes after hemiarthroplasty for femoral neck fracture? *Am J Orthop* 2010;39:E84–9.
45. Baker P, Jameson S, Critchley R, *et al.* Center and surgeon volume influence the revision rate following unicondylar knee replacement: an analysis of 23,400 medial cemented unicondylar knee replacements. *J Bone Joint Surg Am* 2013;95:702–9.
46. Zaidi R, Macgregor AJ, Goldberg A. Quality measures for total ankle replacement, 30-day readmission and reoperation rates within 1 year of surgery: a data linkage study using the NJR data set. *BMJ Open* 2016;6:e011332.
47. Basques BA, Bitterman A, Campbell KJ, *et al.* Influence of surgeon volume on inpatient complications, cost, and length of stay following total ankle arthroplasty. *Foot Ankle Int* 2016;37:1046–51.
48. Clark JC, Simon P, Clark RE, *et al.* The influence of patient- and surgeon-specific factors on operative duration and early postoperative outcomes in shoulder arthroplasty. *J Shoulder Elbow Surg* 2017;26:1011–6.
49. Ramkumar PN, Navarro SM, Frankel WC, *et al.* Evidence-Based Thresholds for the Volume and Length of Stay Relationship in Total Hip Arthroplasty: Outcomes and Economies of Scale. *J Arthroplasty* 2018;33:2031–7.
50. Wilson S, Marx RG, Pan TJ, *et al.* Meaningful Thresholds for the Volume-Outcome Relationship in Total Knee Arthroplasty. *J Bone Joint Surg Am* 2016;98:1683–90.
51. Peschke D, Nimptsch U, Mansky T. Achieving minimum caseload requirements—an analysis of hospital discharge data from 2005–2011. *Dtsch Arztebl Int* 2014;111:556.
52. Browne JA, Cancienne JM, Casp AJ, *et al.* Certificate-of-Need State Laws and Total Knee Arthroplasty. *J Arthroplasty* 2018;33:2020–4.
53. Stroup DF, Berlin JA, Morton SC, *et al.* Meta-analysis of observational studies in epidemiology: a proposal for reporting. *JAMA* 2000;283:2008–12.
54. Moher D, Liberati A, Tetzlaff J, *et al.* Preferred reporting items for systematic reviews and meta-analyses: the PRISMA statement. *PLoS Med* 2009;6:e1000097.
55. Shamseer L, Moher D, Clarke M, *et al.* Preferred reporting items for systematic review and meta-analysis protocols (PRISMA-P) 2015: elaboration and explanation. *BMJ* 2015;349:g7647.
56. Cochran WG. The combination of estimates from different experiments. *Biometrics* 1954;10:101–29.
57. Higgins JP, Thompson SG. Quantifying heterogeneity in a meta-analysis. *Stat Med* 2002;21:1539–58.
58. Higgins JP, Green S. *Cochrane handbook for systematic reviews of interventions version 5.1.0*. The cochrane collaboration, 2011:5.
59. Harbord RM, Harris RJ, Sterne JA. Updated tests for small-study effects in meta-analyses. *Stata J* 2009;9:197.
60. Egger M, Davey Smith G, Schneider M, *et al.* Bias in meta-analysis detected by a simple, graphical test. *BMJ* 1997;315:629–34.
61. Begg CB, Mazumdar M. Operating characteristics of a rank correlation test for publication bias. *Biometrics* 1994;50:1088–101.
62. Greenland S, Longnecker MP. Methods for trend estimation from summarized dose-response data, with applications to meta-analysis. *Am J Epidemiol* 1992;135:1301–9.
63. Orsini N, Bellocco R, Greenland S. Generalized least squares for trend estimation of summarized dose-response data. *Stata J* 2006;6:40.