

Table A2.2: Statin utilisation according to age and indication among female TILDA participants (Wave 1) aged over 50 years (n=3,014)

	Age	Number	Number on statins	% taking statin [95%CI]	MI	IHD	Stroke	PAC	Diabetes	Hyper-tension	High cholesterol	No diagnosis
	50-64	1,868	399	21.4% [19.6%-23.3%]	12	10	14	40	33	132	139	18
	65-74	747	295	39.5% [36.0%-43.1%]	13	19	20	28	23	79	95	17
	75+	399	175	43.9% [39.0%-48.8%]	18	22	14	26	12	51	20	9
	Total	3,014	869	28.8% [27.2%-30.5%]	43	51	48	94	68	262	254	44
Total with diagnosis whether on statins or not [95% CI]					71.7% [58.7%-81.8%]	68.9% [57.3%-78.6%]	60.0% [48.7%-70.3%]	33.9% [28.6%-39.7%]	56.2% [47.1%-64.9%]	35.9% [32.5%-39.4%]	45.9% [41.8%-50.1%]	4.0% [3.0%-5.3%]
% of those with indication on statins					5.0%	5.9%	5.6%	10.9%	7.9%	30.3%	29.4%	5.1%
% of those on statins with												

indication												
------------	--	--	--	--	--	--	--	--	--	--	--	--

Source: Analysis of The Irish Longitudinal Study on Ageing (TILDA) Wave 1 data.

Notes: The column 'Number' represents the total number of individuals in the TILDA dataset by age group and overall, whether they are taking a statin or not, while the column 'Number on statins' represents the total number of individuals taking a statin. The column '% taking statins' is the number on statins as a percentage of all individuals by age group and overall. The subsequent eight columns present the numbers or percentages on statins for each of the indication categories. MI= myocardial infarction; IHD = Ischaemic Heart Disease; PAC= potential atherosclerotic conditions.

Table A2.3: Statin utilisation according to age and indication among male TILDA participants (Wave 1) aged over 50 years (n=2,604)

	Age	Number	Number on statins	% taking statin [95% CI]	MI	IHD	Stroke	PAC	Diabetes	Hyper-tension	High cholesterol	No diagnosis
	50-64	1,519	366	24.1% [22.0%-26.3%]	44	19	11	33	38	95	112	11
	65-74	741	316	42.6% [39.1%-46.2%]	59	37	19	32	38	68	41	17
	75+	344	164	47.7% [42.4%-53.0%]	39	25	14	25	16	22	10	10
	Total	2,604	846	32.5% [30.7%-34.3%]	142	81	44	90	92	185	163	38
Total with diagnosis whether on statins or not					190	105	70	207	156	554	392	908
% of those with indication on statins [95% CI]					74.7% [68.0%-80.5%]	77.1% [68.0%-84.3%]	62.9% [50.7%-73.6%]	43.5% [36.8%-50.4%]	59.0% [51.0%-66.5%]	33.4% [29.6%-37.4%]	41.6% [36.8%-46.5%]	4.2% [3.1%-5.7%]
% of those on statins with indication [95% CI]					17.0%	9.7%	5.3%	10.8%	11.0%	22.2%	19.5%	4.6%

Source: Analysis of The Irish Longitudinal Study on Ageing (TILDA) Wave 1 data.

Notes: The column 'Number' represents the total number of individuals in the TILDA dataset by age group and overall, whether they are taking a statin or not, while the column 'Number on statins' represents the total number of individuals taking a statin. The column '% taking statins' is the number on statins as a percentage of all individuals by age group and overall. The subsequent eight columns present the numbers or percentages on statins for each of the indication categories. MI= myocardial infarction; IHD = Ischaemic Heart Disease; PAC= potential atherosclerotic conditions.