

BMJ Open Assessing the nutrient intake of a low-carbohydrate, high-fat (LCHF) diet: a hypothetical case study design

Caryn Zinn,¹ Amy Rush,² Rebecca Johnson²

To cite: Zinn C, Rush A, Johnson R. Assessing the nutrient intake of a low-carbohydrate, high-fat (LCHF) diet: a hypothetical case study design. *BMJ Open* 2018;**8**:e018846. doi:10.1136/bmjopen-2017-018846

► Prepublication history for this paper is available online. To view these files, please visit the journal online (<http://dx.doi.org/10.1136/bmjopen-2017-018846>).

Received 27 July 2017
Revised 18 December 2017
Accepted 15 January 2018

ABSTRACT

Objective The low-carbohydrate, high-fat (LCHF) diet is becoming increasingly employed in clinical dietetic practice as a means to manage many health-related conditions. Yet, it continues to remain contentious in nutrition circles due to a belief that the diet is devoid of nutrients and concern around its saturated fat content. This work aimed to assess the micronutrient intake of the LCHF diet under two conditions of saturated fat thresholds.

Design In this descriptive study, two LCHF meal plans were designed for two hypothetical cases representing the average Australian male and female weight-stable adult. National documented heights, a body mass index of 22.5 to establish weight and a 1.6 activity factor were used to estimate total energy intake using the Schofield equation. Carbohydrate was limited to <130 g, protein was set at 15%–25% of total energy and fat supplied the remaining calories. One version of the diet aligned with the national saturated fat guideline threshold of <10% of total energy and the other included saturated fat ad libitum.

Primary outcomes The primary outcomes included all micronutrients, which were assessed using FoodWorks dietary analysis software against national Australian/New Zealand nutrient reference value (NRV) thresholds.

Results All of the meal plans exceeded the minimum NRV thresholds, apart from iron in the female meal plans, which achieved 86%–98% of the threshold. Saturated fat intake was logistically unable to be reduced below the 10% threshold for the male plan but exceeded the threshold by 2 g (0.6%).

Conclusion Despite macronutrient proportions not aligning with current national dietary guidelines, a well-planned LCHF meal plan can be considered micronutrient replete. This is an important finding for health professionals, consumers and critics of LCHF nutrition, as it dispels the myth that these diets are suboptimal in their micronutrient supply. As with any diet, for optimal nutrient achievement, meals need to be well formulated.

INTRODUCTION

The low-carbohydrate, high-fat (LCHF) diet is becoming increasingly employed in clinical practice as a dietary means to achieve a variety of health goals, from weight reduction to management of chronic disease, in particular diabetes.^{1–3} This style of eating has been shown to be efficacious both short term and long term for its beneficial outcomes

Strengths and limitations of this study

- A strength of this study is that we used an accurate, professional and local food composition database for dietary analysis.
- A strength of this study is that specialty foods, fortified foods and generally unpopular tasting food (ie, liver and mussels) were specifically excluded from the food selection to avoid bias towards nutrient density.
- A limitation of the study is that these results are specific to the two case studies selected, and inference to population groups cannot be made.

on metabolic health.^{4–8} However, LCHF continues to remain a highly contentious topic in nutrition circles. Two likely reasons for this are as follows: (1) the supposition that LCHF diets are devoid of certain nutrients, and therefore increase risk of nutrient deficiencies^{9 10} and (2) concern around the saturated fat content of the diets and the speculation that high intakes might increase risk of cardiovascular disease. Full-fat versions of animal fat-containing whole foods are not purposefully minimised in an LCHF diet; as a result, the saturated fat intake can exceed the maximum 10% of total energy intake threshold set by the National Australian and New Zealand Nutrient Reference Values (NRV) guidelines.¹¹ Recently, the science supporting the long-standing diet-heart hypothesis and the 10% threshold for saturated fat intake has been challenged; both epidemiological studies and randomised controlled trials have come under criticism for being flawed in research methodology and outcome interpretation.^{12 13} This is an ongoing debate that indicates this area of public health and nutrition guidance needs further work to resolve.^{12 14 15}

The LCHF nutrition approach tends not to exclude any food groups specifically but rather focuses on reducing intake from high-load carbohydrate foods in general;



¹Human Potential Centre, Faculty of Health and Environmental Sciences, Auckland University of Technology, Auckland, New Zealand

²Telethon Type 1 Diabetes Family Centre, Stirling, Western Australia, Australia

Correspondence to
Dr Caryn Zinn;
caryn.zinn@aut.ac.nz

and when carbohydrate is eaten, whole-food sources are preferred to processed ones. In contrast, vegetarian and vegan diets are styles of eating that do exclude several food groups that contain vital micronutrients. Recently, a lacto-ovo vegetarian diet has been shown to be nutrient replete, if well planned using an array of non-animal sources containing these potentially missing nutrients.¹⁶ However, a traditional vegan diet is deficient in vitamin B₁₂, as this vitamin is derived only from foods of animal origin; it is also low in the fat-soluble vitamins A and D, with consumption of fortified foods and supplementation necessary for their repletion.^{17 18} Despite the known nutrient deficiency risks that arise with vegan and some vegetarian eating styles, national and international nutrition organisations are not dissuasive of these diets; nutrition professionals merely address any dietary issues in clinical practice.^{19 20} By contrast, carbohydrate-restricted diets are still frowned on by many dietitians and associated national organisations,¹⁰ despite their endorsement by some organisations such as the Commonwealth Scientific and Industrial Research Organisation.²¹ This study aimed to assess the micronutrient thresholds of two versions of the LCHF diet against national NRV thresholds, as set by the Australian National Health and Medical Research Council (NHMRC) and New Zealand Ministry of Health (MOH)¹¹ under two conditions of saturated fat thresholds.

METHODS

In this descriptive study, we designed two LCHF meal plans for each of two hypothetical case studies representing the average Australian male and female as closely as possible. Using the body mass index (BMI) equation, $BMI = (\text{weight-kg}) / (\text{height-m} \times \text{height-m})$, we inputted national average heights recorded by the Australian Bureau of Statistics 2011–2013 for male and female adults²² and used the midpoint for a healthy range BMI of 22.5 to calculate body weight. To estimate total energy expenditure, we used the Schofield equation,²³ where weight and height variables and an activity factor of 1.6 (light level) were inputted. The adult age range category of 19–50 years was selected from the Australian NHMRC and New Zealand MOH set of categories.¹¹ We used computer analysis software FoodWorks Professional V.8 (Xyris software), which uses an Australian and New Zealand food database. The only NRV that varies within this age category is magnesium, which is slightly higher, for both males and females in the 35–50 years age subcategory than in the 19–35 year age subcategory; this was accounted for in the analysis. Total energy intake would likely change within the age groups and was accounted for in the analysis by ensuring that energy intake was met within 95% of requirements for both a 25-year-old and a 45-year-old male and female, respectively. We assumed weight stability and matched the energy intake with the calculated energy expenditure.

For the macronutrient thresholds, we selected a value of carbohydrate that aligned with the 'low carbohydrate' definition as per Feinman and colleagues,²⁴ that is, <130 g per day (or <26% of total energy). The protein threshold was calculated based on the upper limit of the acceptable macronutrient distribution range (AMDR) as per the Australian NHMRC and New Zealand MOH guidelines (ie, 25% of total energy intake).¹¹ Fat was set as the remaining calories. The two sample meal plans differed only by saturated fat content, which was set at the threshold of 10% or less of total energy intake. All micronutrients were required to achieve at least 100% of the NRVs. The NRVs are a set of recommendations for nutritional intake based on currently available scientific knowledge. NRVs include recommended dietary intakes (RDIs), a term used to represent the average daily micronutrient level sufficient to meet the requirements of almost all (97%–98%) healthy individuals of a certain gender and life stage, and adequate intakes (AIs), where the nutrient intake level is based on observed or experimentally determined nutrient estimates of apparently healthy people and are assumed to be adequate.¹¹ Where the RDI value was not available, the AI value was used. **Table 1** presents the demographic data used for the case studies.

We created two different meal plans for the purpose of ensuring variety in food options, using the same macronutrient and micronutrient thresholds and targets for both sets of plans. For each sample meal plan, the male and female versions differ only by portion sizes to align with personalised energy requirements. For the dual purpose of preventing duplication in **table 2** and wanting to illustrate dietary variety, we have elected to present meal plan sample 1 for females and meal plan sample 2 for males, along with their corresponding diets with saturated fat limits. All meals have been developed with a whole-food principle (ie, using foods that have been minimally processed) as a foundation. We also opted to include foods that we considered to be, generally, popular and acceptable, rather than any specialty or unusual food that would demand an acquired taste.

RESULTS

Tables 2 and 3 present the LCHF sample meal plans for females and males, and the nutrient analysis of the meal plans, respectively, with their corresponding plans aligning with the saturated fat threshold of <10% total energy.

Both of the meal plans successfully exceeded the NRV thresholds for all nutrients apart from two instances. The first was iron intake in females, the two meal plans achieving 86%–98% of the RDI threshold value for this mineral. The second instance was an inability to meet the <10% of total energy intake saturated fat threshold in males (meal plan 2 only). In this plan, saturated fat amounted to 10.6% of total energy, exceeding the threshold by 0.6% (or 2 g).

Table 1 Case study demographics

	Age range(years)	Reference height (cm)	Reference weight (kg)	PAL	Energy (kcal)	Carbohydrate(g)	Protein UL 25% TE (g)	Saturated fat 10% TE (g)
Male	19–50	175	63–71	1.6	2820	65	165.4	30.5
Female	19–50	162	56.3	1.6	2203	65	129.4	23.8

PAL, physical activity level; TE, total energy; UL, upper limit.

DISCUSSION

Overall, the LCHF meal plans successfully achieved almost all of the NRV thresholds. There are several important points about iron that warrant discussion. First, in our meal plan development, we specifically selected whole, unprocessed foods that were not fortified with nutrients, such as iron. We also decided to purposely exclude liver and mussels from these plans, despite their rich nutrient density, as we are aware that these foods may not be considered appealing for the majority of the population. However, it is worthwhile noting that the addition of a small amount of chicken liver (ie, 5g and 25g or 1–4 mussels, in meal plans 1 and 2, respectively) would have resulted in the RDI being met for iron. Our strategy differs somewhat to the work of Reid *et al*, who, knowing iron is a nutrient of concern for vegetarians, intentionally incorporated iron-fortified foods in their vegetarian meal plans. Hence, they were able to meet iron requirements, apart from during pregnancy, where the RDI is higher.¹⁶

Second, iron bioavailability is affected by dietary composition and iron status, two aspects that are not considered comprehensively in RDI threshold generation. For industrialised countries, like Australia and New Zealand, a mean iron bioavailability factor is used to generate the iron RDI for all population groups, irrespective of dietary composition.²⁵ Iron bioavailability is reduced by phytates, found predominantly in wholegrains, such as breads and cereals.²⁵ Other compounds that reduce bioavailability are polyphenols and oxalates, and while found in vegetables and fruit, are also present in wholegrains. The LCHF diet is typically very low (or devoid) in grains, which could mean higher iron bioavailability for those consuming such a diet. This, along with other factors that influence nutrient status, plus the natural variation in food intake raises caution about the use of the RDI threshold alone to assess an individual’s diet at a glimpse. While not presented in this work, our nutrient analysis work on isocaloric diets aligning with the Australian NHMRC and New Zealand MOH guidelines also indicate a failure to meet the RDI threshold for iron. As such, while the LCHF diet is often targeted for being inadequate, using NRVs as a dietary adequacy tool, one can assume a similar inadequacy for iron under mainstream dietary guidance using unfortified foods where possible. It is worthwhile to note that this point can also be applied to other micronutrients, in that it is unlikely that any diet will achieve over 100% of the NRV thresholds each day; hence the reason why dietitians encourage the consumption of a varied diet.

In the instance where the saturated fat threshold of <10% of total energy was not met in one of the male meal plans, comprehensive dietary manipulation of this meal plan for the specific purpose of meeting this target was attempted. To achieve carbohydrate and protein targets, the only way to achieve the energy requirements with a <10% saturated fat threshold was to add an alcoholic beverage into the meal plan, as it is the only food item that provides energy without any other macronutrient. We viewed this as being

Table 2 LCHF sample meal plans

LCHF meal plan 1 (females)	LCHF meal plan 1 (females) (saturated fat <10% TE)	LCHF meal plan 2 (males)	LCHF meal plan 2 (males) (saturated fat <10% TE)
Breakfast ¾ cup frozen mixed berries, 150g plain, unsweetened, full-fat yoghurt, 2 T each sunflower and pumpkin seeds, 3 macadamia nuts, 2 Brazil nuts, 6 almonds	Breakfast ¾ cup frozen mixed berries, 150g plain, unsweetened low-fat yoghurt, 2 T each sunflower and pumpkin seeds, 3 macadamia nuts, 2 Brazil nuts, 6 almonds	Breakfast Omelette: 3 eggs, 60g mushrooms, 100g tomato, 1 cup baby spinach, cooked in 2 tsp butter Coffee made with 200 mL full- fat milk	Breakfast Omelette: 3 eggs, 60g mushrooms, 100g tomato, 1 cup baby spinach, cooked in 3 tsp olive oil Coffee made with 200 mL low- fat milk
Lunch Tuna salad: 95g tin tuna, canned in brine (drained), 1 cup baby spinach leaves, 60g English cucumber, 5 cherry tomatoes, 30g cheddar cheese, 2 T linseeds, 1 tsp basil pesto, 3 tsp olive oil	Lunch Tuna salad: 95g tin tuna, canned in brine (drained), 1 cup baby spinach leaves, 60g English cucumber, 5 cherry tomatoes, 30g low fat cottage cheese, 2 T linseeds, 10 green olives, 5 tsp avocado oil	Lunch Beef salad: 120g eye fillet, 1 cup spinach leaves, half red capsicum, 60g English cucumber, 5 cherry tomatoes, half large avocado, 5 walnuts, 1 T linseeds, 30g parmesan cheese, 2 T olive oil	Lunch Beef salad: 150g eye fillet, 1 cup spinach leaves, half red capsicum, 60g English cucumber, 5 cherry tomatoes, half large avocado, 3 T sunflower seeds, 12 green olives, 30g low fat cottage cheese, 2 T olive oil, 2 tsp avocado oil
Dinner 150g sirloin steak, fat not trimmed, grilled, roasted vegetables: 8 florets cauliflower, 1 medium-sized beetroot, 1 medium courgette, 1 medium carrot, coated in 2 T olive oil	Dinner 150g beef eye fillet, grilled, roasted vegetables: 8 florets cauliflower, 1 medium-sized beetroot, 1 medium courgette, 1 medium carrot, coated in 2 T olive oil	Dinner 130g grilled salmon, 100g green beans, 150g broccoli, 200g grilled pumpkin, half cup peas, 1 T olive oil	Dinner 150g grilled salmon, 100g green beans, 150g broccoli, 200g grilled pumpkin, half cup peas, 1 T olive oil
Snacks 10 medium strawberries 3 T pistachio nuts Coffee made with 200 mL full- fat milk	Snacks 10 medium strawberries 3 T pistachio nuts 100g low fat plain, unsweetened yoghurt Coffee made with 200 mL low- fat milk	Snacks 20 macadamia nuts Smoothie made with 200 mL full-fat milk, 10 medium strawberries, crushed nuts (20g almonds, 2 T linseeds)	Snacks 20 macadamia nuts Smoothie made with 200 mL low-fat milk, 10 medium strawberries, crushed nuts (20g almonds, 2 T linseeds, 100g low-fat plain, unsweetened yoghurt)

LCHF, low-carbohydrate, high-fat diet; TE, total energy; T, tablespoon; tsp, teaspoon.

somewhat futile so decided against this option. It is also important to note that during this dietary manipulation exercise, to reduce the saturated fat contribution, once all the relevant animal foods were altered to low-fat or non-fat versions (ie, dairy products and meats) and coconut products were removed, the saturated fat content still slightly exceeded the 10% threshold by 0.6%. It was only when we reduced the amount of avocados, certain seeds, olive oil and macadamia nuts, that is, foods that contain predominantly unsaturated fats, did the thresholds align. In New Zealand and Australia, saturated fat guidelines exist in the form of a percentage of total energy threshold, and it is not known whether the public health caution for saturated fat relates to an absolute amount consumed in grams or is relative to total energy only, suggesting that as a guideline it is rather arbitrary.

There are three other nutrient components that warrant discussion in the context of LCHF diets: B vitamins and fibre, protein and essential fatty acids.

B vitamins and fibre

The LCHF diet is frequently criticised for being deficient in B vitamins in particular vitamin B₁ or thiamin and fibre, two key components of grain-based foods. We have demonstrated that these meal plans do indeed meet the RDI threshold for thiamin due to the incorporation of alternative, every day thiamin-rich foods such as animal protein, nuts and seeds and several green vegetables. Despite RDIs met for thiamin, one could call into question the minimum threshold of thiamin required in an LCHF context. A key function of thiamin is the metabolism of carbohydrate²⁶; it could be speculated that with a reduced intake of carbohydrate, less thiamin is required. However, considering that in the absence of exogenous carbohydrate, glucose is still made internally through gluconeogenic precursors, research is warranted to determine whether this theory holds any truth. We also demonstrate that the AIs for fibre are surpassed in all of these meal plans, and while devoid of wholegrains, fibre, both soluble and insoluble, can be easily derived from vegetables, some fruit, nuts and seeds.

Table 3 Nutrient analysis of LCHF meal plans

Nutrient	Female meal plans			Male meal plans		
	Meal plan 1	Meal plan 1 (saturated fat<10% TE)	NRV/goal	Meal plan 2	Meal plan 2 (saturated fat<10% TE)	NRV/goal
Energy (calories)	2145	2053	2203	2675	2758	2820
Carbohydrate (g)	61	67	248–358	66	69	303–439
% TE	11	13	45–65	10	10	45–65
Protein (g)	115	135	83–138	149	164	106–176
% TE	22	26	15–25	22	24	15–25
Fat (g)	153	129	49–86	194	195	63–110
% TE	63	57	20–35	65	64	20–35
Saturated fat (g)	40	21	24	46	33	31
% TE	28	9.6	10	15	10.6	10
Trans fats (g)	2.2	0.7	<2.4	1.4	0.8	<3g
% TE	0.9	0.3	<1%*	0.4	0.3	<1%*
MUFA (g)	75	71	–	101	117	–
% total fat	53	59	–	56	65	–
PUFA (g)	27	28	–	32	31	–
% total fat	19	23	–	18	17	–
Linoleic acid (O6 PUFA) (g)	20.4	18.6	8†	19.1	18.5	13†
Alpha-linoleic acid (omega-3 PUFA) (g)	5.8	5.5	0.8†	9.3	6.2	1.3†
Omega-6:omega-3 ratio	3.5	3.4	10	2.1	3.0	10
Fibre (g)	38	39	25†	45	44	30†
Thiamin (mg)	1.4	1.4	1.1	1.6	1.8	1.2
Riboflavin (mg)	2.4	2.4	1.1	3.3	3.5	1.3
Niacin (mg)	23.4	18.8	14	16.2	17.5	16
Vitamin C (mg)	371	370	45	394	398	45
Vitamin A (µg)	2247	2095	700	2374	2047	900
Vitamin E (mg)	23	22	7†	32	41	10†
Vitamin B ₁₂ (µg)	3.9	6.4	2.4	11.6	12.8	2.4
Folate, total (µg)	568	583	400	788	757	400
Calcium (mg)	1093	1224	1000	1216	1251	1000
Iron (mg)	16	16	18	20	21	8
Magnesium (mg)	553	589	310–320	582	598	400–420
Zinc (mg)	16	22	8	23	24	14
Sodium (mg)	2183	2250	460†	1554	2032	460†
Potassium (mg)	4639	5154	2800†	5585	6107	3800†
Phosphorous (mg)	1848	2076	1000	2478	2644	1000
Selenium (µg)	166	169	60	113	117	70
Iodine (µg)	225	190	150	223	207	150

*WHO recommendation for trans fats.

†AIs were used as RDIs were unavailable.

AI, adequate intake; LCHF, low-carbohydrate, high-fat diet; MUFA, monounsaturated fat; NRV, nutrient reference value; PUFA, polyunsaturated fat; RDI, recommended daily intake; TE, total energy.

Protein

Another misconception about the LCHF diet is that it is excessive in protein. We used the protein AMDR as per the Australian NHMRC and New Zealand MOH in the development of these meal plans; however, the female meal plan 1 (<10% saturated fat threshold) exceeded the maximum AMDR protein threshold by 2% (or 7g of protein). In the dietary manipulation required to reduce saturated fat

(ie, a swap from untrimmed sirloin steak to lean eye fillet steak for dinner and a swap from full-fat to non-fat dairy products), what resulted was a higher proportion of protein being derived from those foods at the same quantities so it was unavoidable. While the LCHF diet is not intended to be any higher in protein than current dietary recommendations, in this case, protein only exceeded the AMDR when saturated fat was restricted.

Omega-6:omega-3 polyunsaturated fat (PUFA) ratio

A key characteristic of the LCHF way of eating is the discouragement of consumption of seed oils (eg, canola, sunflower, soybean, corn oil and rice bran oil) to minimise intake of linoleic acid (omega-6 PUFA), and consequently, to achieve an optimal omega-6:omega-3 ratio. Evidence suggests that a high ratio is proinflammatory and has a role to play in promoting the pathogenesis of chronic diseases such as cardiovascular disease, cancer, inflammatory and autoimmune diseases.^{27 28} Furthermore, data indicates that humans evolved on a diet with a 1:1 omega-6:omega-3 ratio, whereas Western dietary patterns typically reflect a ratio of around 15-20:1.²⁸ Along with the promotion of olive oil use, which is unanimously endorsed, NHMRC/MOH guidelines recommend the use and consumption of vegetable fats (ie, margarine, canola, sunflower, soybean, corn oil and rice bran oil) in place of fats with a predominantly saturated fat make-up (ie, butter, coconut oil). As a result, most packaged supermarket foods including foods recommended by the NHMRC/MOH, such as liquid breakfasts, wholegrain bread and cereals contain omega-6-rich seed oils. This would not necessarily be problematic if omega-3 intakes were increasing on a population level. However, this is not the case; it was recently reported in the Australian 2011–2012 National Nutrition and Physical Activity Survey that 80% of the population was not meeting the NRV threshold for omega-3.²⁹ In our sample meal plans, we demonstrate favourable omega-6:omega-3 ratios, that is, less than the 10:1 NRV thresholds and substantially closer to that of our dietary composition prior to the agricultural revolution.

Finally, it is important to note that LCHF eating is frequently adopted for weight loss purposes. In this context, it would be highly likely that energy intakes would be lower than that of these hypothetical healthy weight case studies for a certain period of time, while weight is being lost. This poses a risk to achieving 100% of all NRVs on a daily basis; however, this would not be unique to the LCHF approach but would apply to any energy-restricted eating style, including mainstream national nutrition guidelines. During the active weight loss period, nutrient density should be a priority and if suboptimal nutrient status becomes a concern, this could be addressed by the inclusion of nutrient-fortified foods or supplementation.

CONCLUSION

There were two key limitations to this work. First, the average height for Australian male and female, respectively, and the midpoint BMI values were used to extrapolate the weights of the two hypothetical case studies. These weights do not align with documented weights for Australian males (85.9 kg) and females (71.1 kg), respectively.²² This underestimation of weight would have underestimated energy intake to a small extent. Second, our analyses were limited to the available values in the

FoodWorks database; consequently, an accurate estimate of vitamin D intake was not available. It is reported that due to current eating patterns, it is almost impossible to get sufficient vitamin D (in this case 5 µg/day) from the diet alone,³⁰ and it is the assumption that it will be derived from sun exposure. It is likely that the LCHF diet supplies more vitamin D than mainstream nutrition guidelines due to its greater intake of high fat, vitamin D-rich foods, such as full-fat dairy, butter, eggs and fatty fish.

We have demonstrated that a well-formulated LCHF diet can provide sufficient intakes of all of the micronutrients profiled in the FoodWorks database, apart from iron for females. This marginal shortfall along with the acknowledged limitations of using NRVs in estimating dietary adequacy leads us to believe that this is not a nutrient of concern for those consuming the LCHF diet. Irrespective of the ongoing saturated fat/heart disease scientific debate, it is still possible to adopt the LCHF diet while keeping saturated fat intake around the 10% of total energy threshold. Considering this way of eating provides a replete set of nutrients and has been shown to be effective for improving metabolic health, particularly for people with diabetes, it should at least be considered a suitable dietary option for populations, alongside that of mainstream MOH guidelines.

Contributors All authors contributed to the conception and design of the research, interpretation of data; all authors read and approved the final manuscript; CZ, AR contributed to the diet development and analysis. CZ drafted the manuscript.

Funding This research received no specific grant from any funding agency in the public, commercial or not-for-profit sectors.

Competing interests CZ has coauthored two books called *What the Fat?—Fat's In, Sugar's Out* and *What the Fat—Sports Performance*, which both assume an LCHF nutrition approach.

Patient consent Not required.

Provenance and peer review Not commissioned; externally peer reviewed.

Data sharing statement Data (nutrient analysis) are unable to be placed on a data sharing system due to it being embedded in specific nutrient analysis software that is unable to be shared outside of the software program. However, a print screen version of the data is available on request from the author.

Open Access This is an Open Access article distributed in accordance with the Creative Commons Attribution Non Commercial (CC BY-NC 4.0) license, which permits others to distribute, remix, adapt, build upon this work non-commercially, and license their derivative works on different terms, provided the original work is properly cited and the use is non-commercial. See: <http://creativecommons.org/licenses/by-nc/4.0/>

© Article author(s) (or their employer(s) unless otherwise stated in the text of the article) 2018. All rights reserved. No commercial use is permitted unless otherwise expressly granted.

REFERENCES

1. Unwin D, Unwin J. Low carbohydrate diet to achieve weight loss and improve HbA1c in type 2 diabetes and pre-diabetes: experience from one general practice. *Practical Diabetes* 2014;31:76–9.
2. Noakes TD. Low-carbohydrate and high-fat intake can manage obesity and associated conditions: occasional survey. *S Afr Med J* 2013;103:826–30.
3. McKenzie AL, Hallberg SJ, Creighton BC, et al. A novel intervention including individualized nutritional recommendations reduces hemoglobin A1c Level, medication use, and weight in type 2 diabetes. *JMIR Diabetes* 2017;2:e5.

4. Fan Y, Di H, Chen G, *et al.* Effects of low carbohydrate diets in individuals with type 2 diabetes: systematic review and meta-analysis. *Int J Clin Exp Med* 2016;9:11166–74.
5. Forsythe CE, Phinney SD, Fernandez ML, *et al.* Comparison of low fat and low carbohydrate diets on circulating fatty acid composition and markers of inflammation. *Lipids* 2008;43:65–77.
6. Sackner-Bernstein J, Kanter D, Kaul S. Dietary intervention for overweight and obese adults: comparison of low-carbohydrate and low-fat diets. A Meta-Analysis. *PLoS One* 2015;10:e0139817.
7. Foster GD, Wyatt HR, Hill JO, *et al.* Weight and metabolic outcomes after 2 years on a low-carbohydrate versus low-fat diet: a randomized trial. *Ann Intern Med* 2010;153:147–57.
8. Shai I, Schwarzfuchs D, Henkin Y, *et al.* Weight loss with a low-carbohydrate, Mediterranean, or low-fat diet. *N Engl J Med* 2008;359:229–41.
9. Diabetes UK. Is it time to stop promoting carbohydrates to people with diabetes? 2017. cited 8 Mar 2017 https://www.diabetes.org.uk/About_us/News/Carbohydrates-and-diabetes-debate/ (accessed 15 Sep 2015).
10. Nursing Review. Fad diets: what do dietitians say about the latest crop? 2015 <http://www.nursingreview.co.nz/issue/june-2015-vol-15-3/fad-diets-what-do-dietitians-say-about-the-latest-crop/#.WL8yITXJKnk> (accessed 5 Jun 2015).
11. National Health and Medical Research Council, Ministry of Health. Nutrient reference values for Australia and New Zealand Including Recommended Dietary Intakes. 2005.
12. Harcombe Z, Baker JS, Davies B. Evidence from prospective cohort studies does not support current dietary fat guidelines: a systematic review and meta-analysis. *British J Sports Med* 2016;0:1–8.
13. Harcombe Z, Baker JS, DiNicolantonio JJ, *et al.* Evidence from randomised controlled trials does not support current dietary fat guidelines: a systematic review and meta-analysis. *Open Heart* 2016;3:e000409.
14. Ramsden CE, Zamora D, Majchrzak-Hong S, *et al.* Re-evaluation of the traditional diet-heart hypothesis: analysis of recovered data from Minnesota Coronary Experiment (1968-73). *BMJ* 2016;353:i1246.
15. Siri-Tarino PW, Chiu S, Bergeron N, *et al.* Saturated fats versus polyunsaturated fats versus carbohydrates for cardiovascular disease prevention and treatment. *Annu Rev Nutr* 2015;35:517–43.
16. Reid MA, Marsh KA, Zeuschner CL, *et al.* Meeting the nutrient reference values on a vegetarian diet. *Med J Aust* 2013;199(4 Suppl):S33–40.
17. Pawlak R, Lester SE, Babatunde T. The prevalence of cobalamin deficiency among vegetarians assessed by serum vitamin B12: a review of literature. *Eur J Clin Nutr* 2014;68:541–8.
18. Craig WJ. Nutrition concerns and health effects of vegetarian diets. *Nutr Clin Pract* 2010;25:613–20.
19. American Dietetic Association/Dietitians of Canada. Position of the American Dietetic Association and Dietitians of Canada: Vegetarian diets. *J Am Diet Assoc* 2003;103:748–65.
20. Webb D. *Defending vegan diets—RDs aim to clear up common misconceptions about vegan diets. Today's dietitian.* Spring City, PA: Great Valley Publishing Company, Inc., 2010. 20.
21. CSIRO. CSIRO Low Carb Diet Book 2017 18 April, 2017. <https://www.csiro.au/en/Research/Health/CSIRO-diets/CSIRO-Low-Carb-Diet-Book> (accessed 18 Apr 2017).
22. Australian Bureau of Statistics. Profiles of Health, Australia, 2011-13 2013. cited 3 Feb 2017 <http://www.abs.gov.au/ausstats/abs@.nsf/Lookup/4338.0main+features212011-13> (accessed 3 Feb 2017).
23. Schofield WN. Predicting basal metabolic rate, new standards and review of previous work. *Hum Nutr Clin Nutr* 1985;39(Suppl 1):5–41.
24. Feinman RD, Pogozelski WK, Astrup A, *et al.* Dietary carbohydrate restriction as the first approach in diabetes management: critical review and evidence base. *Nutrition* 2015;31:1–13.
25. Hurrell R, Egli I. Iron bioavailability and dietary reference values. *Am J Clin Nutr* 2010;91:1461S–7.
26. Manzetti S, Zhang J, van der Spoel D. Thiamin function, metabolism, uptake, and transport. *Biochemistry* 2014;53:821–35.
27. Kang JX. The omega-6/omega-3 fatty acid ratio in chronic diseases: animal models and molecular aspects. *World Rev Nutr Diet* 2011;102:22–9.
28. Simopoulos AP. The importance of the omega-6/omega-3 fatty acid ratio in cardiovascular disease and other chronic diseases. *Exp Biol Med* 2008;233:674–88.
29. Meyer BJ. Australians are not meeting the recommended intakes for omega-3 long chain polyunsaturated fatty acids: results of an analysis from the 2011-2012 National Nutrition and Physical Activity Survey. *Nutrients* 2016;8:111.
30. Fuller KE, Casparian JM. Vitamin D: balancing cutaneous and systemic considerations. *South Med J* 2001;94:58–64.