

**Appendix 3:** Prevalence, n (%) of the most common comorbidities in each country

<b>Country</b>	<b>Myocardial Infarction (MI)</b>	<b>Angina</b>	<b>Stroke</b>	<b>Transient Ischemic Attack (TIA)</b>	<b>Heart Failure (HF)</b>	<b>Peripheral Vascular Disease (PVD)</b>	<b>Revascularisation</b>	<b>Atrial Fibrillation (AF)</b>	<b>Hypertension</b>	<b>Diabetes</b>	<b>Epilepsy</b>	<b>Osteoporosis</b>
Ireland	7 (6.1)	7 (6.1)	4 (3.5)	3 (2.6)	3 (2.6)	1 (1.0)	17 (14.8)	10 (8.7)	75 (65.2)	17 (14.8)	0 (0)	10 (8.7)
Switzerland	13 (8.0)	4 (2.5)	5 (3.1)	12 (7.4)	7 (4.3)	11 (6.8)	29 (18.0)	22 (13.6)	83 (51.2)	20 (12.3)	4 (2.5)	21 (13.0)
The Netherlands	21 (8.2)	25 (9.8)	15 (5.9)	16 (6.3)	12 (4.7)	9 (3.5)	26 (10.2)	32 (12.5)	114 (44.7)	38 (15)	2 (1)	32 (12.5)