BMJ Open Impact of parental weight status on children's body mass index in early life: evidence from a Chinese cohort

Hong Mei,^{1,2} Siyu Guo,¹ Hongyan Lu,¹ Yunhong Pan,³ Wenhua Mei,⁴ Bin Zhang,² Jianduan Zhang¹

To cite: Mei H, Guo S, Lu H, *et al.* Impact of parental weight status on children's body mass index in early life: evidence from a Chinese cohort. *BMJ Open* 2018;**8**:e018755. doi:10.1136/bmjopen-2017-018755

► Prepublication history for this paper is available online. To view these files, please visit the journal online (<http://dx.doi.org/10.1136/bmjopen-2017-018755>).

HM and SG contributed equally.

Received 24 July 2017

Revised 22 November 2017

Accepted 24 November 2017

¹Department of Maternal and Child Health Care, School of Public Health, Tongji Medical College, Huazhong University of Science and Technology, Wuhan, China

²Wuhan Children's Hospital (Wuhan Maternal and Child Healthcare Hospital), Tongji Medical College, Huazhong University of Science and Technology, Wuhan, China

³Department of cardiology, Tianyou Hospital, Wuhan University of Science and Technology, Wuhan, Hubei, China

⁴Department of Information, Public Hospital Administration of Zhuhai Municipality, Zhuhai, China

Correspondence to

Dr Jianduan Zhang;
jd_zh@hust.edu.cn

ABSTRACT

Objectives To understand whether parents' weight status before conception predicts body mass index (BMI) of their offspring in early life and the differences between the mother-child and father-child associations.

Design A birth cohort study.

Setting Conducted at the Community Health Service Centre in Shenyang, Wuhan and Guangzhou.

Participants A total of 2220 live birth newborns were recruited randomly after consent of their parents, and 1178 were followed up until 2 years old.

Methods Parental demographics, maternal characteristics during pregnancy, children's anthropometric data and feeding patterns at 1 month old were collected. BMI was calculated and BMI Z-scores (BMI_Z) were generated by referring to WHO growth standard. Parental weight status was categorised into underweight, normal weight, and overweight and obese according to the Working Group of Obesity in China. General linear models and generalised linear models were used to assess the associations between parents and offspring.

Outcome measures The primary outcomes were descriptive data on child's sex-specific anthropometric variables. The secondary outcomes were BMI_Z and weight status of children at each time point.

Results No gender difference was observed in BMI_Z or overweight or obesity rates from birth to 24 months old, although boys were significantly heavier and had a greater length/height than girls ($P < 0.05$). The overweight and obesity rates of children peaked at 12 months old. Maternal BMI/weight status had a significant but small effect on BMI_Z at birth, but not on the paternal side. The impact of parental BMI on child's BMI_Z after birth was similar at each follow-up. Offspring with underweight mothers tend to have reduced BMI_Z after birth while overweight/obese fathers had children with a greater BMI_Z.

Conclusions Maternal weight status had small effect on both fetal and child growth after birth. Significant but mild paternal influence was only detected after birth.

INTRODUCTION

Childhood obesity is a serious public health concern¹ because of its adverse long-term health consequences^{2,3} and is related directly and indirectly to lifetime economic burden.⁴ Data from the 2014 WHO show that the

Strengths and limitations of this study

- The study is among the few studies in an Asian population that have compared the impacts of parental body mass index (BMI) and weight status on offspring BMI Z-score (BMI_Z) at multiple time points during early age.
- The comparatively large proportion of underweight mothers in this study allowed us to investigate the association between maternal underweight and reduced offspring BMI_Z after birth.
- As BMI is correlated with both lean and fat mass, BMI could not precisely reflect the adiposity level of our subjects.
- The self-reported weight and height of parents in our study may have resulted in an underestimate of parental overweight and obesity.

overweight or obese rate among Chinese children <5 years old is about 10%,⁵ which is much higher than that from two decades earlier.⁶ As it is difficult to lose weight or to sustain successful weight loss,^{7,8} detecting and addressing the causes of overweight or obesity from the early years of life, especially during the critical stage of adiposity development in the prenatal period⁹ and early infancy,¹⁰ is of great public health importance.

Both genetic factors and an obesogenic environment play roles in the development of obesity.^{11–13} Genetic studies have identified more than 20 genetic variations that are related to the development of childhood obesity.^{14,15} Meanwhile, environmental obesogenic-related factors such as a high energy diet, short sleep duration and insufficient physical activity, also contribute to childhood obesity.^{16,17} Parental overweight/obesity is considered as a proxy for the genetic predisposition and/or the obesogenic exposure, and should be considered when tackling the prevalence of childhood obesity. A large number of studies have confirmed a significant correlation between maternal or/and paternal weight status and the growth

measures of their school-age offspring.^{18 19} The parental weight status' impact on children through similar genetic preposition and/or obesogenic environmental exposure may 'programme' the growth and development of children from a very early stage of life.^{20 21} However, the connection between parental weight status and growth of children <2 years old has been rarely discussed. The limited number of studies that have analysed the association between parental body mass index (BMI) and toddler growth, have mostly yielded inconsistent results.^{22 23} For example, Regnault *et al* and Knight *et al* reported a positive association between maternal BMI and their offspring's birth weight and/or BMI,^{24 25} whereas Stunkard *et al* revealed a null association at 2 years old.²³ Of the studies that have examined the impact of paternal BMI on offspring birth weight, only one reported any effect.²⁶ Fewer studies have compared the effect of maternal and paternal weight status on postpartum growth in offspring, but with discrepant results.^{27 28} So far, conclusions have predominately been drawn from populations of European descent,²⁹ which may not be applicable to other populations, as the parent-offspring association in BMI could vary given significant differences in social and cultural conditions.^{5 30 31} Furthermore, there is a lack of longitudinal data to examine the parental-offspring association in BMI and obesity during the early years of life, especially the father-child association in an Asian population.

Based on data from a 2-year birth cohort in three cities in China, we (1) explored the association between parental BMI/weight status and child's BMI Z-score (BMI_Z) over the first 2 years of life and (2) compared the difference, if any, between the impacts from maternal versus paternal weight status.

METHODS

Study population and design

A total of 2220 newborns (1155 boys and 1065 girls) were recruited within 1 month after birth from April 2009 to March 2010 at randomly selected Community Health Service Centres (CHSC) in Shenyang, Wuhan and Guangzhou. The participating children were registered and routinely received health checks and follow-ups at the project's CHSC. Details of participant selection have been published elsewhere.^{32 33} After recruitment, the study subjects were followed at 3, 12 and 24 months (figure 1). The permanent withdrawals included consented withdrawals and those lost to follow-up. Missing data resulted from those who could not attend all follow-ups because of parent unavailability for the regular healthcare visit, child's sickness, inclement weather or other unknown reasons. No significant differences were detected in the socioeconomic variables, children's birth variables or parental BMI and weight status between the initially recruited subjects and those available at each follow-up.

All protocol-required procedures were standardised across the three field sites and data collectors were trained

Figure 1 Follow-up chart from birth to 24 months old.

and monitored centrally. Informed consent was obtained from the parents.

Data collection

Characteristics of children and parents

Socioeconomic and health-related variables were obtained at recruitment from the Perinatal Health Booklet (PHB) held by parents and from standardised interviews at 1-month-old and 3-month-old follow-ups. Data extracted from the PHB were gestational weight gain (GWG), health condition during pregnancy (eg, gestational hypertension and diabetes), gestational age at birth, delivery mode and the child's sex and date of birth. Children's birth weight and crown-heel length, as measured by nurses at birth, were also extracted from the PHB. Data collected through interviews included maternal lifestyle during pregnancy (such as maternal gestational smoking, secondhand smoking and alcohol consumption), parental age and educational level at childbirth, number of children in the household and infant feeding practices within the first month.

Anthropometric measurements

Children's weight and length/height were measured at 3 and 12 months old within 10 days of the exact month, and within 15 days of 24 months old, by the centrally trained child healthcare staff from the project CHSC. Body weight was measured to the nearest 50 g with an electronic scale (WHS-I, Wuhan Computer Software Development), with children wearing light indoor clothes. Recumbent length was measured using an infant length measurement instrument (WHS-I) before 24 months old, and a measuring tape (S-RT-Y1 Healthy kid Intelligent Health Check

Wuhan Computer Software Development) attached to a wall was used for height at 24 months old, both to the nearest 0.1 cm. Both weight and length/height were taken in duplicate, and the means of the replicates were used in the analyses. Parental weights before pregnancy were self-reported, while heights were measured using a stadiometer (Leicester height measure; Invicta Plastic, Leicester, UK) during the first-month-old follow-up home visit.

Variables

Outcome variables

Children's BMI at birth and at 3, 12 and 24 months old as well as BMI_Z were employed as the primary outcome variables. BMI was calculated as weight (kilogram) divided height (metre) squared and the age-specific and gender-specific Z-score (BMI_Z) was generated according to WHO Child Growth Standards 2006, using the SAS macro provided to assess child growth and development.³⁴ Overweight and obesity for children were the secondary outcome and were defined as BMI $\geq$ 85th and 95th percentiles, respectively, according to WHO gender-specific and age-specific BMI reference values for children.³⁵

Independent variables

Parental BMI was set as a continuous independent variable and further categorised into weight status groups, based on BMI cut-offs for Chinese adults recommended by the Working Group of Obesity in China: BMI <18.5 kg/m² as underweight, BMI ≥ 24 kg/m² as overweight/obese and BMI of 18.5–24 kg/m² as normal weight.³⁶

Potential confounders

The originally included potential confounders were selected based on previous reports. Low birth weight and macrosomia were defined as birth weight <2500 g or ≥ 4000 g, respectively. GWG was set as a continuous variable. Delivery mode was categorised as vaginal birth or caesarean section (C-section) and preterm birth referred to gestational age at birth <37 weeks. Parental education level was classified as middle school or under, high school/technical, university/college and master's degree or advanced. The feeding pattern at 1 month of age was categorised as exclusive breast feeding, mixed feeding and formula feeding. Exclusive breast feeding was defined according to WHO definition.³⁷ Mixed-fed children were fed with both breast milk and formula milk, or other food. Formula-fed children were those fed exclusively with formula milk or other food but no breast milk. Variables such as single-child family, gestational hypertension, gestational diabetes, smoking, secondhand smoking and alcohol consumption during pregnancy were set as dichotomous variables (yes/no). The act of smoking a cigarette or cigar at least once per day on average was defined as smoking, and secondhand smoking was defined as being exposed to cigarette smoke accumulatively for at least 15 min for at least 1 day per week by

non-smoking pregnant women³⁸; alcohol consumption as consumption at least one alcoholic drink per day.³⁹

Statistical analysis

Continuous variables are presented as mean and SD except for BMI with median and IQR, which was in a skewed distribution, and categorical variables are presented as percentages or rates. A tabulated procedure was used to capture the descriptive characteristics of children and parents at recruitment and at the endpoint. Student's t-test was used to compare differences in child weight, length/height, BMI and BMI_Z between boys and girls at birth and at 3, 12 and 24 months old; and sex-specific overweight and obesity rates of children at each age point were generated using the χ^2 test.

General linear models were adopted to model child BMI_Z at the four follow-ups as a function of parental BMI and covariates. Covariates were selected according to previous reports and a stepwise regression analysis. The covariates of child's sex, gestational age at birth, maternal age and GWG were included in the model for BMI_Z at birth. Delivery mode was also adjusted for in the BMI_Z models at 3, 12 and 24 months old. In addition, to examine the mediating effects of birth size on the association between parental BMI and the outcome variables, two BMI_Z models at 3, 12 and 24 months old were performed without and with adjustment for the BMI_Z at birth. A similar analytical strategy was applied for the generalised linear models to assess the association between parental weight status and children's BMI_Z.

The database was set up in Epidata 3.1 with all analyses done with SAS V.9.4 software. The level of statistical significance was set at $P < 0.05$ for all hypothesis testing.

RESULTS

A total of 2220 participants were recruited at birth, and 1800, 1677 and 1178 participants were successfully followed up until 3, 12 and 24 months old, respectively. At recruitment, 52% of the neonates were boys and more than 87% were the only child in the household. Low birthweight children accounted for 1.7%, whereas 6.8% of children were born with macrosomia. C-sections made up 68% of deliveries in our population. Maternal secondhand smoking during pregnancy was found in 19.9% of cases, compared with 0.7% firsthand smoking, and 1.6% of the study mothers consumed alcohol during pregnancy. More than 50% of parents had received a bachelor degree. More details about the characteristic of study subjects at recruitment and at 24 months old are presented in table 1. No significant differences were found in the parents' or children's characteristics between the baseline and the final stage.

Table 2 lists the crude sex-specific and age-specific weight, length/height, BMI, BMI_Z, overweight and obesity rates of children at the four age points. Child's BMI, BMI_Z, overweight and obesity rate increased during the first year but declined from 12 to 24 months.

Table 1 Baseline characteristics of the samples at recruitment and at the endpoint

Variables	At recruitment (N=2220)		At endpoint (N=1178)	
	n (%)	Mean (SD)	n (%)	Mean (SD)
Boys	1155 (52.0)		611 (51.9)	
Single-child family	1937 (87.3)		1032 (87.6)	
Birth weight (g)		3334 (419)		3325 (424)
Low birth weight	34 (1.5)		20 (1.7)	
Normal birth weight	2028 (91.4)		1078 (91.5)	
Macrosomia	158 (7.1)		80 (6.8)	
Maternal age at childbirth (years)	2217	28.4 (3.8)	1176	28.6 (3.8)
Paternal age at childbirth (years)	2211	31.1 (4.7)	1172	31.3 (4.8)
Maternal prepregnancy BMI (kg/m ²)	2215	20.4 (2.6)*	1174	20.4 (2.6)
Maternal underweight	541 (24.4)		487 (22.8)	
Maternal normal weight	194 (66.9)		1463 (68.5)	
Maternal overweight/obesity	194 (8.8)		187 (8.8)	
Paternal BMI at recruitment (kg/m ²)	2201	23.5 (3.3)*	1169	23.6 (3.4)
Paternal underweight	87 (4.0)		81 (3.8)	
Paternal normal weight	1232 (55.9)		1169 (55.0)	
Paternal overweight/obesity	884 (40.1)		877 (41.2)	
Gestational weight gain	2205	17.1 (5.4)	1168	17.2 (5.6)
Gestational age at birth	2206	39.5 (1.5)	1162	39.5 (1.5)
Preterm	80 (3.6)		47 (4.0)	
Full term	2126 (96.4)		1115 (96.0)	
Delivery mode				
Vaginal	699 (31.5)		376 (31.9)	
C-section	1521 (68.5)		802 (68.1)	
Maternal pregnancy hypertension	52 (2.6)		30 (2.7)	
Pregnancy smoking	15 (0.7)		8 (0.8)	
Pregnancy secondhand smoking	435 (19.9)		240 (20.7)	
Pregnancy alcohol consumption	35 (1.6)		20 (1.7)	
Maternal educational level				
Middle school or less	432 (19.5)		227 (19.3)	
High school/technical	584 (26.3)		321 (27.3)	
University/college	1117 (50.3)		582 (49.4)	
Master degrees or advanced	87 (3.9)		48 (4.1)	
Paternal educational level				
Middle school or less	343 (15.5)		180 (15.3)	
High school/technical	644 (29.0)		329 (27.9)	
University/college	1105 (49.8)		595 (50.5)	
Master degrees or advanced	128 (5.8)		74 (6.3)	
Feeding pattern at the first month				
Exclusive breast feeding	759 (36.7)		418 (37.1)	
Mixed feeding	1015 (49.1)		532 (48.2)	
Formula feeding	295 (14.3)		153 (13.9)	

*BMI is presented as median (IQR).

BMI, body mass index; C-section, caesarean section.

Table 2 Sex-specific anthropometric characteristics and overweight and obesity rates of children from birth to 24 months old

Age (month)	Weight* mean (SD)	Height/length* mean (SD)	BMI* median (IQR)	BMI_Z mean (SD)	Overweight, n (%)	Obesity, n (%)
Boys						
0	3.39 (0.01)	50.55 (0.05)	13.25 (0.04)	−0.19 (0.03)	91 (7.88)	26 (2.25)
3	6.94 (0.03)	62.77 (0.07)	17.50 (0.05)	0.34 (0.04)	110 (11.74)	115 (12.27)
12	10.56 (0.04)	77.08 (0.08)	17.77 (0.05)	0.64 (0.03)	171 (16.49)	172 (16.59)
24	13.03 (0.05)	88.28 (0.12)	16.72 (0.05)	0.53 (0.04)	212 (19.17)	174 (15.73)
Girls						
0	3.27 (0.01)	50.02 (0.05)	13.08 (0.04)	−0.27 (0.04)	59 (5.53)	39 (3.66)
3	6.37 (0.02)	61.29 (0.08)	16.88 (0.05)	0.27 (0.03)	117 (13.31)	71 (8.08)
12	9.93 (0.04)	75.67 (0.09)	17.32 (0.05)	0.60 (0.03)	181 (19.53)	116 (12.51)
24	12.47 (0.05)	87.45 (0.13)	16.30 (0.06)	0.45 (0.04)	133 (12.83)	124 (11.96)

*Refers to significant difference between boys and girls at each time point, $P < 0.05$.

BMI, body mass index; BMI_Z, BMI Z-score.

Significant sex differences in the growth measures (weight, length/height and BMI) and insignificant sex difference in BMI_Z and the rate of overweight and obesity were detected at all follow-ups.

Results from the general linear models show that maternal BMI had a significant effect on the child's BMI_Z at birth and later (3, 12 and 24 months old), and the impact of maternal BMI on child's BMI_Z at birth was significantly larger than that on the after-birth measures (0.07 vs 0.02–0.04, $P < 0.05$); No meaningful impact from the father's side was detected. The generalised linear models showed that children from underweight mothers had significantly lower BMI_Z from birth to 24 months old compared with those with normal weight mothers. Similarly, overweight/obese mothers tended to have heavier children, which was only significant at birth ($\beta = 0.30$, $P < 0.05$). Consistent with the general linear models,

paternal overweight/obesity had a significant effect on the children's BMI_Z after birth but not for BMI_Z at birth (at birth: $\beta = 0.04$, $P > 0.05$; after birth: $\beta = 0.04$ –0.20, $P < 0.05$). Both parents' BMI and weight status had almost unchanged coefficients in the unadjusted and adjusted models with child's sex, gestational age at birth, maternal age and GWG included being the major confounders (table 3).

DISCUSSION

In the present study, we examined the impacts of maternal and paternal weight status on offspring growth during their early ages. Our results show that maternal weight status was mildly but significantly associated with fetal growth, and this association persisted until after the child's birth. In contrast to normal weight mothers,

Table 3 Effect of parental weight status on child's BMI_Z from birth to 24 months old

		General linear model		Generalised linear model			
				Maternal weight status (reference: normal weight)		Paternal weight status (reference: normal weight)	
Age (month)		Maternal BMI	Paternal BMI	Underweight	Overweight/obesity	Underweight	Overweight/obesity
0	–	0.07*	0.01	−0.22*	0.30*	−0.12	0.04
3	Unadjusted	0.03*	0.03*	−0.17*	0.11	−0.31*	0.11*
	Adjusted	0.02*	0.02*	−0.14*	0.05	−0.29*	0.10*
12	Unadjusted	0.03*	0.03*	−0.13*	0.12	−0.18	0.16*
	Adjusted	0.02*	0.02*	−0.11*	0.10	−0.17	0.15*
24	Unadjusted	0.04*	0.03*	−0.16*	0.12	0.03	0.20*
	Adjusted	0.04*	0.03*	−0.15*	0.11	0.20*	0.04

Variables controlled for were the same in the general linear models and generalised linear models; in the birth models, child's sex, gestational age, maternal age, educational level and gestational weight gain were controlled for; in the unadjusted models for BMI_Z at aged 3–24 months old, delivery mode was also added to the factors controlled for in the birth models; in the adjusted models, BMI_Z at birth was also controlled for apart from the factors controlled for in the unadjusted models.

*Refers to significant effect, $P < 0.05$; variables controlled for were the same in the general linear models and generalised linear models.

BMI, body mass index; BMI_Z, BMI Z-score.

underweight mothers had children with significantly lower BMI_Z from birth to 24 months old, whereas overweight/obesity mothers were more likely to have children with significantly higher BMI_Z at birth. The impacts from the father's side were found to be mild and only significant after children were born.

Previous longitudinal studies reported inconsistent results regarding parental-offspring BMI associations. Both Adegboye *et al*⁴⁰ and Linabery *et al*²⁷ reported a significantly greater association between maternal weight and offspring BMI than for the paternal side. Li *et al* indicated that obesity-related genes inherited from parents may play a role during the fetal period,⁴¹ and results from a large Norwegian population-based study showed a similar parental-offspring BMI association when children were 3 years old.⁴² In our study, we found an association between maternal BMI and offspring's BMI_Z from birth to 24 months old, while the significant association between paternal BMI and offspring's BMI_Z only became apparent after birth. Our results seemed to suggest that the genetic influence from the father side, if any, might be overshadowed by the impact of maternal weight status through an intrauterine nutritional environment and/or a genetic influence from the mother's side.

The association between parental weight status and child's BMI_Z has been reported previously.¹¹ In our study, when parental prepregnancy BMI value was further stratified into three weight status groups (underweight, normal weight and overweight/obesity), maternal underweight was significantly associated with a reduced child's BMI_Z after birth, while the expected positive associations between parental overweight/obese and the outcome measure only existed in obese paternal group. The arguable results of the non-significant associations between maternal overweight/obesity and children's BMI_Z after birth could be attributed to several factors. First, it may be due to the small number of overweight/obese mothers (8.7%) and relatively large number of underweight mothers (24.4%), which may reduce the power to detect the mother-offspring association. Second, it could be attributed to the termination of the mother's role as the direct and only nutrition provider, the involvement of more environmental factors (eg, feeding practices) and the probability of interactions between environmental factors and genetic predispositions if stimulated.^{43–45} Interestingly, in our study, only about 10% of children had a BMI >85th percentile at birth and this figure increased rapidly and reached nearly 30% at both 12 and 24 months old, indicating that at least some of the children may have experienced rapid growth during infancy. According to a study by Wells *et al* on a group of Brazilian boys, early rapid weight gain remained the dominant risk factor for later obesity, which may weaken the effect from the maternal side.⁴⁶

According to the National Report on Nutrition and Chronic Diseases in Chinese Residents in 2015, the prevalence of overweight and obesity rate in Chinese adults were up to 30.1% and 11.9% in 2012.⁴⁷ The relatively

lower overweight/obesity rate in the current maternal population (7.39% and 1.35%, respectively, with the mean prepregnancy BMI being 20.4 kg/m²) and high underweight rate (24.4%) compared with the national figure might be attribute to different characteristics such as population age (about 28 years old) and the study area (mostly central and southern China). The large proportion of thin women made the current cohort distinguishable from previous studies mainly performed in high-income counties where the average BMI was heavily shifted to the right.⁴⁸ This allowed the detection of the association between maternal undernourishment and the reduced toddler BMI. Thinness (wasting) in early childhood has been significantly related to reduced cognitive development, and increased risk for impaired immune system, poor school performance and a higher risk for morality.^{49–51} As a country that has been experiencing nutrition transition, China is now facing a double burden of nutrition problems. While a great deal of attention has been attached to addressing the emerging obesity issue, undernourishment, due to an unfavourable nutrition status from the early years of life or as a result of a mindset of 'thinner is prettier', appears to be neglected in China especially in the affluent urban area. More attention should be redirected to the group of thin women who are at their childbearing age given the health implication of maternal nutrition in both mother and child.

The current study had some strengths. First, this was a longitudinal study with a relatively large sample size. Previous studies were mostly cross-sectional in design,^{42 52} whereas others were longitudinal studies but with a smaller sample size.⁵³ Second, we confirmed the significant association between parental BMI and child's BMI_Z at first, then classified the parental BMI as underweight or overweight/obesity to better introduce the clinical implications of our findings. This made our results more convenient for identifying high-risk individuals based on their parents' weight status, either overweight/obesity or underweight. Furthermore, this is among the first studies to compare the parental influence on the BMI_Z of children at an early age in an Asian population.

The use of parental BMI based on their self-reported weight may be one of our weaknesses, as it may be less precise compared with anthropometric data collected under a more carefully controlled research setting. Low maternal overweight/obesity rate and high paternal overweight/obesity rate in our study may limit the generalisability of our finding. Parental overweight and obesity is considered to be due to both genetic predisposition and obesogenic environmental exposure. Using parental weight status, for example, overweight/obesity or underweight as exposure, could not distinguish the impacts of genetic predisposition from that of environmental exposure or the interaction of the two. More genetic studies are needed to explore the genetic impact on the growth of children from an early stage of life.

Research on body composition in infancy suggests that most of the variance in BMI is explained by variance in

lean mass.⁴⁶ In our study, BMI was used as the proxy of obesity in children, which is correlated with both lean and fat mass. There was no measurement of body composition of infants available. Because of the comparatively short follow-up durations, we could not verify the association between birth weight, rapid weight gain with body composition of children later in life; and we could not detect whether this presents any inherited similarity in term of maternal and paternal impacts on offspring's body composition. More studies should include precisely measured body composition with longer observation duration to clarify the clinical implications of BMI and the variation in the development of body composition in early childhood. Last, the comparatively high lost to follow-up rate may limit the generalisability of our findings and reduce the power to detect associations. However, no significant differences were detected in the sociodemographic or birth variables, parental BMI or weight status between the initially recruited and the remaining participants indicating the randomness of those lost to follow-ups. We believe the chance of random lost to follow-up errors affecting our conclusions was slim. Furthermore, the consistency between our findings and those of other related studies supports the generalisability of the findings.

CONCLUSIONS

Our results suggest that the parents' BMI and weight status affected the growth of their offspring in different ways from the intrauterine period through the first 24 months of life. Maternal BMI and weight status appeared to influence intrauterine growth, and this mild effect persisted during the child's first 24 months of life. Paternal weight status had less of a significant impact on fetal growth, but showed a small but significant and continuous impact on child's postpartum growth, although the precise mechanism remains unknown. Nevertheless, our results stress that maintaining a healthy prepregnancy weight for both mother and father is critical for a healthy prenatal and postnatal growth of children. Further studies are needed to more precisely evaluate the genetic and environmental impacts and their interactions on fetal and child growth, and to clarify the underlying mechanisms of the different maternal-offspring and paternal-offspring associations in BMI.

Acknowledgements The authors acknowledge the Community Health Service Centres at Shenyang, Wuhan and Guangzhou for providing the data for this study. The contributions of National Natural Science Fund of China and Special Fund for Basic Scientific Research of Central Colleges and individuals involved are gratefully acknowledged.

Contributors MH collected the data, performed the statistical analyses and the interpretation of data, and drafted, revised and finalised the article. SG contributed to data collecting, writing the background, results and discussions sections of the article. HL, YP, WM and BZ helped finalise the drafted and revised manuscripts. JZ was responsible for the design and implementation of the study, and supervised the data collection, data analysis and manuscript writing.

Funding This study was supported by the National Natural Science Fund of China (grant nos: 81373017 and 81673182). It was also partly supported by the Special Fund for Basic Scientific Research of Central Colleges (grant no: 2016YXZD041).

Competing interests None declared.

Patient consent Obtained.

Ethics approval The Ethics Committee of Tongji Medical College, Huazhong University of Science and Technology (IORG0003571).

Provenance and peer review Not commissioned; externally peer reviewed.

Data sharing statement No data are available.

Open access This is an open access article distributed in accordance with the Creative Commons Attribution Non Commercial (CC BY-NC 4.0) license, which permits others to distribute, remix, adapt, build upon this work non-commercially, and license their derivative works on different terms, provided the original work is properly cited and the use is non-commercial. See: <http://creativecommons.org/licenses/by-nc/4.0/>

© Article author(s) (or their employer(s) unless otherwise stated in the text of the article) 2018. All rights reserved. No commercial use is permitted unless otherwise expressly granted.

REFERENCES

1. Song Y, Wang HJ, Ma J, *et al.* Secular trends of obesity prevalence in urban Chinese children from 1985 to 2010: gender disparity. *PLoS One* 2013;8:e53069.
2. Valerio G, Licenziati MR, Manco M, *et al.* [Health consequences of obesity in children and adolescents]. *Minerva Pediatr* 2014;66:381–414.
3. Berenson GS, Bogalusa Heart Study group. Health consequences of obesity. *Pediatr Blood Cancer* 2012;58:117–21.
4. Sonntag D, Ali S, De Bock F. Lifetime indirect cost of childhood overweight and obesity: A decision analytic model. *Obesity* 2016;24:200–6.
5. Swinburn B, Vandevijvere S. WHO report on ending childhood obesity echoes earlier recommendations. *Public Health Nutr* 2016;19:1–2.
6. Ji CY. Working Group on Obesity in China (WGOC). Report on childhood obesity in China (4) prevalence and trends of overweight and obesity in Chinese urban school-age children and adolescents, 1985–2000. *Biomed Environ Sci* 2007;20:1–10.
7. Oken E, Gillman MW. Fetal origins of obesity. *Obes Res* 2003;11:496–506.
8. Ebbeling CB, Pawlak DB, Ludwig DS. Childhood obesity: public-health crisis, common sense cure. *Lancet* 2002;360:473–82.
9. Dietz WH. Critical periods in childhood for the development of obesity. *Am J Clin Nutr* 1994;59:955–9.
10. Gillman MW. Early infancy as a critical period for development of obesity and related conditions. *Nestle Nutr Workshop Ser Pediatr Program* 2010;65:13–20.
11. Woo Baidal JA, Locks LM, Cheng ER, *et al.* Risk Factors for Childhood Obesity in the First 1,000 Days: A Systematic Review. *Am J Prev Med* 2016;50:761–79.
12. Gardner KR, Sapienza C, Fisher JO. Genetic and epigenetic associations to obesity-related appetite phenotypes among African-American children. *Pediatr Obes* 2015;10:476–82.
13. Engeli S, Lehmann AC, Kaminski J, *et al.* Influence of dietary fat intake on the endocannabinoid system in lean and obese subjects. *Obesity* 2014;22:E70–6.
14. Felix JF, Bradfield JP, Monnereau C, *et al.* Genome-wide association analysis identifies three new susceptibility loci for childhood body mass index. *Hum Mol Genet* 2016;25:389–403.
15. Speliotes EK, Willer CJ, Berndt SI, *et al.* MAGICProcardis Consortium. Association analyses of 249,796 individuals reveal 18 new loci associated with body mass index. *Nat Genet* 2010;42:937–48.
16. Birdsey J, Sussell AL. Prevalence of Obesity, No Leisure-Time Physical Activity, and Short Sleep Duration Among Occupational Groups in 29 States. *J Occup Environ Med* 2017;59:1221–8.
17. Ventura AK. Does Breastfeeding Shape Food Preferences? Links to Obesity. *Ann Nutr Metab* 2017;70(Suppl 3):8–15.
18. Savva SC, Kourides Y, Tornaritis M, *et al.* Obesity in children and adolescents in Cyprus. Prevalence and predisposing factors. *Int J Obes Relat Metab Disord* 2002;26:1036–45.
19. Krassas GE, Tzotzas T, Tsamietis C, *et al.* Determinants of body mass index in Greek children and adolescents. *J Pediatr Endocrinol Metab* 2001;14(Suppl 5):1327–33.
20. Craig CL, Cameron C, Tudor-Locke C. Relationship between parent and child pedometer-determined physical activity: a sub-study of the CANPLAY surveillance study. *Int J Behav Nutr Phys Act* 2013;10:8.

21. Swaminathan S, Thomas T, Yusuf S, *et al.* Clustering of diet, physical activity and overweight in parents and offspring in South India. *Eur J Clin Nutr* 2013;67:128–34.
22. Zalbazar N, Jan Mohamed HJ, Loy SL, *et al.* Association of parental body mass index before pregnancy on infant growth and body composition: Evidence from a pregnancy cohort study in Malaysia. *Obes Res Clin Pract* 2016;10 Suppl 1:S35–47.
23. Stunkard AJ, Berkowitz RI, Schoeller D, *et al.* Predictors of body size in the first 2 y of life: a high-risk study of human obesity. *Int J Obes Relat Metab Disord* 2004;28:503–13.
24. Regnault N, Botton J, Forhan A, *et al.* Determinants of early ponderal and statural growth in full-term infants in the EDEN mother-child cohort study. *Am J Clin Nutr* 2010;92:594–602.
25. Knight B, Shields BM, Hill A, *et al.* The impact of maternal glycemia and obesity on early postnatal growth in a nondiabetic Caucasian population. *Diabetes Care* 2007;30:777–83.
26. Xue F, Willett WC, Rosner BA, *et al.* Parental characteristics as predictors of birthweight. *Hum Reprod* 2008;23:168–77.
27. Linabery AM, Nahhas RW, Johnson W, *et al.* Stronger influence of maternal than paternal obesity on infant and early childhood body mass index: the Fels Longitudinal Study. *Pediatr Obes* 2013;8:159–69.
28. Kivimäki M, Lawlor DA, Smith GD, *et al.* Substantial intergenerational increases in body mass index are not explained by the fetal overnutrition hypothesis: the Cardiovascular Risk in Young Finns Study. *Am J Clin Nutr* 2007;86:1509–14.
29. Botton J, Scherdel P, Regnault N, *et al.* EDEN Mother-Child Cohort Study Group. Postnatal weight and height growth modeling and prediction of body mass index as a function of time for the study of growth determinants. *Ann Nutr Metab* 2014;65:156–66.
30. Sommer I, Griebler U, Mählknecht P, *et al.* Socioeconomic inequalities in non-communicable diseases and their risk factors: an overview of systematic reviews. *BMC Public Health* 2015;15:914.
31. Lakshman R, Zhang J, Zhang J, *et al.* Higher maternal education is associated with favourable growth of young children in different countries. *J Epidemiol Community Health* 2013;67:595–602.
32. Mei H, Guo B, Yin B, *et al.* Interactive effects of early exclusive breastfeeding and pre-pregnancy maternal weight status on young children's BMI - a chinese birth cohort. *PLoS One* 2015;10:e0144357.
33. Zhang J, Himes JH, Guo Y, *et al.* Birth weight, growth and feeding pattern in early infancy predict overweight/obesity status at two years of age: a birth cohort study of Chinese infants. *PLoS One* 2013;8:e64542.
34. De Onis M. *WHO Child Growth Standards: Length/height-for-age, weight-for-age, weight-for-length, weight-for-height and body mass index-for-age: Methods and development*, 2006.
35. Krebs NF, Himes JH, Jacobson D, *et al.* Assessment of child and adolescent overweight and obesity. *Pediatrics* 2007;120(Suppl 4):S193–228.
36. Wen CP, David Cheng TY, Tsai SP, *et al.* Are Asians at greater mortality risks for being overweight than Caucasians? Redefining obesity for Asians. *Public Health Nutr* 2009;12:497–506.
37. Jones AD, Ickes SB, Smith LE, *et al.* World Health Organization infant and young child feeding indicators and their associations with child anthropometry: a synthesis of recent findings. *Matern Child Nutr* 2014;10:1–17.
38. World Health Organisation. *Guidelines for controlling and monitoring the tobacco epidemic*. Geneva: World Health Organisation, 1997.
39. Mikkelsen EM, Riis AH, Wise LA, *et al.* Alcohol consumption and fecundability: prospective Danish cohort study. *BMJ* 2016;354:i4262.
40. Adegboye AR, Andersen LB, Wedderkopp N, *et al.* Influence of parental overweight on the association of birth weight and fat distribution later in childhood. *Obes Facts* 2012;5:784–94.
41. Li A, Robiou-du-Pont S, Anand SS, *et al.* Parental and child genetic contributions to obesity traits in early life based on 83 loci validated in adults: the FAMILY study. *Pediatr Obes* 2016.
42. Fleten C, Nystad W, Stigum H, *et al.* Parent-offspring body mass index associations in the Norwegian Mother and Child Cohort Study: a family-based approach to studying the role of the intrauterine environment in childhood adiposity. *Am J Epidemiol* 2012;176:83–92.
43. Mei H, Johansson E, Hagströmer M, *et al.* Physical Activity Levels in Chinese One-Year-Old Children and Their Parents, an Early STOPP China Study. *PLoS One* 2016;11:e0153605.
44. Waterland RA. Epigenetic epidemiology of obesity: application of epigenomic technology. *Nutr Rev* 2008;66(Suppl 1):S21–3.
45. Davey Smith G, Steer C, Leary S, *et al.* Is there an intrauterine influence on obesity? Evidence from parent child associations in the Avon Longitudinal Study of Parents and Children (ALSPAC). *Arch Dis Child* 2007;92:876–80.
46. Wells JC, Hallal PC, Wright A, *et al.* Fetal, infant and childhood growth: relationships with body composition in Brazilian boys aged 9 years. *Int J Obes* 2005;29:1192–8.
47. Prevention C. *National report on nutrition and chronic diseases in Chinese Residents*. 2015.
48. Barisic T, Mandic V, Barac I. Associations of body mass index and gestational weight gain with term pregnancy outcomes. *Mater Sociomed* 2017;29:52–7.
49. Faruque AS, Ahmed AM, Ahmed T, *et al.* Nutrition: basis for healthy children and mothers in Bangladesh. *J Health Popul Nutr* 2008;26:325–39.
50. Murray CJ, Lopez AD. Global mortality, disability, and the contribution of risk factors: global burden of disease study. *Lancet* 1997;349:1436–42.
51. Pelletier DL, Frongillo EA, Schroeder DG, *et al.* A methodology for estimating the contribution of malnutrition to child mortality in developing countries. *J Nutr* 1994;124:2106S–22.
52. McCarthy K, Ye YL, Yuan S, *et al.* Parental weight status and offspring cardiovascular disease risks: a cross-sectional study of Chinese children. *Prev Chronic Dis* 2015;12:E01.
53. Park MH, Falconer CL, Croker H, *et al.* Predictors of health-related behaviour change in parents of overweight children in England. *Prev Med* 2014;62:20–4.