

Supplementary table 2: Adherence to protocols

	n (%)	NP			PA		
		T0	T1	T2	T0	T1	T2
General written protocols present		31 (70.5)	97 (81.5)	145 (84.8)	16 (66.7)	44 (68.8)	74 (76.3)
Completely satisfying		15 (55.6)	51 (56.7)	81 (58.3)	10 (66.7)	21 (51.2)	44 (63.8)
Partly satisfying		11 (40.7)	37 (41.1)	54 (38.8)	5 (33.3)	18 (43.9)	24 (43.8)
Not satisfying		1 (3.7)	2 (2.2)	4 (2.9)	0 (0.0)	2 (4.9)	1 (1.4)
Ambiguities about							
Work instructions		2 (8.3)	6 (5.0)	9 (6.0)	0 (0.0)	7 (11.1)	4 (4.7)
Distribution of tasks		1 (4.2)	14 (11.7)	34 (21.9)*	0 (0.0)	5 (7.9)	14 (14.7)
Distribution of responsibilities		8 (33.3)	12 (10.0)	35 (22.6)*	1 (7.1)	8 (12.7)	10 (10.6)
Personalized protocols present, including		19 (42.2)	67 (57.3)	89 (57.4)	11 (42.3)	23 (37.7)	33 (36.7)
Specified reserved procedures allowed to perform on own authority		11 (61.1)	45 (38.5)	70 (55.6)	8 (72.2)	13 (22.4)	28 (38.4)*
Specified patient groups, allowed to treat on own authority		10 (55.6)	41 (35.0)	57 (41.0)	5 (45.5)	7 (12.1)	19 (23.8)*
Specified medication, allowed to prescribe on own authority		9 (50.0)	49 (41.9)	81 (58.3)	4 (36.4)	12 (20.7)	29 (36.3)
Method to assess competence		3 (16.7)	13 (11.1)	29 (20.9)	1 (9.1)	9 (15.5)	4 (5.0)
Distribution of responsibilities		2 (11.1)	16 (13.7)	25 (18.0)	2 (18.2)	6 (10.3)	8 (10.0)
Adherence to personalized protocols examined		11 (55.0)	24 (36.9)	43 (43.0)	5 (45.5)	15 (57.7)	15 (38.5)

* statistically significant difference between T0-T2 (X2 (1) p<0,05)