



BMJ Open is committed to open peer review. As part of this commitment we make the peer review history of every article we publish publicly available.

When an article is published we post the peer reviewers' comments and the authors' responses online. We also post the versions of the paper that were used during peer review. These are the versions that the peer review comments apply to.

The versions of the paper that follow are the versions that were submitted during the peer review process. They are not the versions of record or the final published versions. They should not be cited or distributed as the published version of this manuscript.

BMJ Open is an open access journal and the full, final, typeset and author-corrected version of record of the manuscript is available on our site with no access controls, subscription charges or pay-per-view fees (<http://bmjopen.bmj.com>).

If you have any questions on BMJ Open's open peer review process please email [info.bmjopen@bmj.com](mailto:info.bmjopen@bmj.com)

# BMJ Open

## The Accuracy and Utility of Using Administrative Healthcare Databases to Identify People with Epilepsy: A Protocol for a Systematic Review and Meta-Analysis

Journal:	<i>BMJ Open</i>
Manuscript ID	bmjopen-2017-020824
Article Type:	Protocol
Date Submitted by the Author:	15-Jan-2018
Complete List of Authors:	Mbizvo, Gashirai; University of Edinburgh Centre for Clinical Brain Sciences, Muir Maxwell Epilepsy Centre, Bennett, Kyle; University of Edinburgh Centre for Clinical Brain Sciences, Muir Maxwell Epilepsy Centre, Simpson, Colin; Victoria University of Wellington, Faculty of Health,; University of Edinburgh School of Molecular Genetic and Population Health Sciences, Usher Institute of Population Health Sciences and Informatics, Duncan, Susan; Western General Hospital, Department of Clinical Neurosciences, Chin, Richard; University of Edinburgh Centre for Clinical Brain Sciences, Muir Maxwell Epilepsy Centre,
Keywords:	Epilepsy < NEUROLOGY, Factual database, Administrative claims, Algorithms, Validation studies

SCHOLARONE™  
Manuscripts

**Title: The Accuracy and Utility of Using Administrative Healthcare Databases to Identify People with Epilepsy: A Protocol for a Systematic Review and Meta-Analysis**

**Authors:** Gashirai K Mbizvo<sup>1</sup>, Kyle Bennett<sup>1</sup>, Colin R Simpson<sup>2,3</sup>, Susan E Duncan<sup>4</sup>, Richard FM Chin<sup>1,5</sup>

<sup>1</sup> University of Edinburgh Centre for Clinical Brain Sciences, Muir Maxwell Epilepsy Centre, Edinburgh, UK

<sup>2</sup> Victoria University of Wellington, Faculty of Health, Wellington, NZ

<sup>3</sup> University of Edinburgh School of Molecular Genetic and Population Health Sciences, Usher Institute of Population Health Sciences and Informatics, Edinburgh, UK

<sup>4</sup> Western General Hospital, Department of Clinical Neurosciences, Edinburgh, UK

<sup>5</sup> Royal Hospital for Sick Children, Edinburgh, UK

**Keywords:** Epilepsy, factual database, administrative claims, algorithms, validation studies

**Corresponding Author**

Dr Gashirai K Mbizvo, *University of Edinburgh Centre for Clinical Brain Sciences, Muir Maxwell Epilepsy Centre, Edinburgh, UK*  
*Tel: +44 (0) 131 536 0801, Fax: N/A*  
*email: [gashirai.mbizvo@ed.ac.uk](mailto:gashirai.mbizvo@ed.ac.uk)*

**Word count:**

Abstract – 299 words,

Body – 3,860 words,

Tables – 2,

Illustrations/figures – 0,

References – 32.

## Abstract

**Introduction:** In an increasingly digital age for healthcare around the world, administrative data have become rich and accessible tools for potentially identifying and monitoring population trends in diseases including epilepsy. However, it remains unclear (1) how accurate administrative data are at identifying epilepsy within a population, and (2) the optimal algorithms needed for administrative data to correctly identify people with epilepsy within a population. To address this knowledge gap, we will conduct a novel systematic review of all identified studies validating administrative healthcare data in epilepsy identification. We provide here a protocol that will outline the methods and analyses planned for the systematic review.

**Methods and analysis:** The systematic review described in this protocol will be conducted to follow the Preferred Reporting Items for Systematic Reviews and Meta-Analyses (PRISMA) guidelines. MEDLINE and EMBASE will be searched for studies validating administrative data in epilepsy published between 01/01/1975 and 03/01/2018. Included studies will validate the International Classification of Disease (ICD) 9th revision onwards (ICD-9 code 345 and ICD-10 codes G40-G41). The primary outcome will be providing pooled estimates of accuracy for identifying epilepsy within the administrative databases validated using sensitivity, specificity, positive and negative predictive values, and area under the receiver operating characteristic curves. Heterogeneity will be assessed using the  $I^2$  statistic and descriptive analyses used where this is present. The secondary outcome will be the optimal administrative data algorithms for correctly identifying epilepsy. These will be identified using multivariable logistic regression models. 95% confidence intervals will be quoted throughout. We will make an assessment of risk of bias, quality of evidence, and completeness of reporting for included studies.

**Ethics and dissemination:** Ethical approval is not required as primary data will not be collected. Results will be disseminated in peer-reviewed journals, conference presentations, and in press releases.

**PROSPERO registration:** CRD42017081212.

## Strengths and limitations of this study

### Strengths

- The protocol describes what will be the first systematic review to conduct a worldwide assessment of the accuracy of administrative data in identifying epilepsy and the optimal disease-identification algorithms.
- This protocol also describes what will be the first systematic review to make an assessment of risk of bias, quality of evidence, and completeness of reporting for studies validating administrative healthcare data in epilepsy identification.

### Limitations

- The review described in this protocol will be limited to assessing the use of administrative data in diagnosing epilepsy within observational studies rather than randomised controlled trials. Observational studies are more prone to biases such as selection bias, which may unevenly distribute clinical or demographic characteristics of the study population and thereby influence the coding practices. We will therefore include a risk of bias assessment for each study. A systematic review of the diagnostic accuracy of administrative data within randomised controlled trials in epilepsy remains to be completed.

## Introduction

Administrative healthcare databases are electronic data sources that consist of demographic, diagnostic, and clinical information routinely collected about patients when they use a healthcare

service<sup>3</sup>. They are often national and mandatory, and therefore they have the potential to provide a relatively cheap, widely-available, and less intrusive resource for medical research<sup>3</sup>. However, the accuracy of the information held in an administrative database needs to be validated before such use can be made. This is because the administrative data were not originally collected for research, but for other purposes such as assisting in health insurance claims. The clinical information held may therefore lack the rigor in accuracy that might be expected in scientifically collected data. Furthermore, the data may be limited by inaccurate or incomplete hospital discharge letters or clinical coding transcription errors<sup>4</sup>.

The validation of administrative data involves comparing the diagnostic codes held within the administrative database against a reference standard (such as medical records) in order to quantify the number of instances in which the administrative diagnosis made matches the diagnosis in the reference standard (deemed to be the true diagnosis). In this way, the administrative database can be handled like a diagnostic test and measures of disease-identification accuracy calculated. These measures usually include the sensitivity, specificity, and the positive or negative predictive value (PPV or NPV, respectively). Optimal disease-identification algorithms can also be determined by making relative comparisons of predictive values after adding in data from other variables recorded in an administrative database, such as drug combinations, investigations, and procedures.

There are many administrative databases worldwide in which the process of validation has been successfully performed for many diagnostic codes<sup>5-11</sup>. There are also examples of where the results of these have been pooled successfully into systematic review to increase confidence in the estimates made and scrutinise the quality of evidence, and this has led to changes in practice<sup>6</sup>. There has been limited systematic review of the validation of administrative databases in capturing epilepsy as a diagnosis. The only systematic review<sup>12</sup> on this subject included only studies from the USA or Canada and therefore excluded 121 studies because the data sources were not from these two countries. Furthermore, the eleven studies included were published between 2000 and 2010, making the conclusions nearly a decade old now. With health informatics now at the forefront of epidemiological disease surveillance it is important to have an update on performance of the administrative disease-identification codes. Only one of the included studies evaluated the performance of the ICD-10 system in capturing epilepsy within administrative datasets<sup>13</sup> –the remainder evaluated the older ICD-9 system<sup>12</sup>. The review also made no assessment of risk of bias, quality of evidence, and completeness of reporting for included studies. This limits the confidence with which conclusions can be interpreted. There is now need for a more contemporary systematic review of the validation of administrative databases in capturing epilepsy. This should include evaluating performance of the ICD-10 system, as well as other non-ICD disease classification systems used, such as Read Codes in the United Kingdom<sup>14 15</sup>. The review should include studies from anywhere in the world in order to give clinicians and researchers a representative picture of the performance of administrative data in capturing epilepsy as a diagnosis and in order to allow more generalisable diagnostic algorithms to be suggested. Furthermore, the review should make an assessment of risk of bias, quality of evidence, and completeness of reporting for included studies. These are the aims of the proposed systematic review described in this protocol. This will help researchers and clinicians better understand the accuracy of global estimates for incidence, prevalence, and population characteristics in epilepsy – which have largely been made using administrative data.

Aims and objectives:

The study hypotheses are:

1. Administrative data can correctly identify people with epilepsy within a population with a high degree of accuracy. We predict the PPV to be above 80%.

2. The optimal disease-identification algorithms for epilepsy within administrative datasets take into account diagnoses, investigations, and drug combinations<sup>13 16-22</sup>.

The aim of the systematic review is to quantify the disease-identification accuracy and algorithm performance of administrative healthcare data in epilepsy. To this end, the research questions are:

1. How accurately do administrative data identify epilepsy within a population? –as measured by sensitivity, specificity, PPV, NPV, or area under the receiver operating characteristic curve (AUC) analysis (which are the approved accuracy measures described in the Standards for Reporting Diagnostic Accuracy (STARD) statement<sup>23</sup>). This will be the primary outcome.
2. What are the optimal administrative data algorithms for correctly identifying epilepsy within a population? This will be the secondary outcome.

A preliminary feasibility search of the MEDLINE database via PubMed identifies at least nine studies validating diagnostic epilepsy codes held within administrative databases around the world that could be utilised to answer these research questions<sup>13 15-22</sup>.

## Methods and analysis

The systematic review will be designed and conducted to follow the PRISMA checklist<sup>24</sup>.

### Eligibility criteria

We will include studies according to the following criteria:

- *Language*: There will be no language restriction on full-length articles although abstracts will need to be in English to allow the authors to screen them. We will seek translations for full length articles not written in English that appear eligible in abstract. These will remain in the section for “studies still awaiting classification” and will feature in subsequent updates to the review if not translated by the time of initial publication.
- *Setting*: There will be no restrictions by study location worldwide. Where possible, we will show pooled accuracy and best algorithm data for administrative datasets from individual countries in addition to pooled global estimates of these measures.
- *Databases*: The data sources will be routine administrative healthcare databases. This means that the data should have been routinely and passively collected without an *a priori* research question<sup>6</sup>. We will include databases containing diagnostic codes for epilepsy classified on the ICD system – where we will restrict this to studies using the ninth ICD revision onwards (ICD-9, active from 1975 to 1994)<sup>25</sup>. This is because although the ICD system is currently in its tenth revision (ICD-10, active from 1994 to present) and this the primary coding system utilised by many countries around the world, a significant proportion of countries, particularly developing ones, still use the ICD-9 system<sup>13</sup>. ICD-9 code 345 and ICD-10 codes G40 and or G41 will be used to identify epilepsy diagnoses. We will provide summary measures of accuracy and best algorithms for any other disease classification systems used in studies separately, e.g. the primary care Read Code system used in the UK<sup>14 15</sup>.
- *Study design*: Prospective or retrospective observational studies including cohort or case-control designs that are community-based/population-based or primary/secondary/tertiary care-based and have utilised administrative databases.
- *Participants*: People with epilepsy of all ages. Where available, we will additionally report data for adults and children separately.
- *Observations and outcomes*: Studies will need to employ a validation process for diagnostic epilepsy codes, i.e. estimate the disease-identification accuracy of the epilepsy codes held within the database using sensitivity, specificity, PPV, NPV, or AUC analysis<sup>23</sup>. In this, true positives (TPs) and false negatives (FNs) will be considered as the patient has the disease and the administrative diagnosis is positive or negative, respectively. True negatives (TNs) and false

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

positives (FPs) will be considered as the patient does not have the disease and the administrative diagnosis is negative or positive, respectively. Sensitivity will be considered as the ability of the administrative database to correctly identify those patients with the disease (TP/(TP + FN). Specificity will be considered as the ability of the administrative database to correctly identify those patients without the disease (TN/(TN + FP). PPV will be considered as how likely it is that a patient has the disease if the administrative diagnosis is positive (TP/(TP + FP). NPV will be considered as how likely it is that this patient does not have the disease if the administrative diagnosis is negative (TN/(TN + FN). AUC analysis will be considered as where TP and FP are plotted against each other in a perfect correlation as reference to show poor test accuracy, then accuracy measured as the area under the curve created by plotting the actual values against each other<sup>26</sup>.

- Studies may also use diagnostic as well as other variables (e.g. admissions, drugs or investigations) to calculate optimal disease-identification algorithms for epilepsy within the database. Studies will need to provide a clinical reference standard. An appropriate clinical reference standard will be medical records, clinical assessment, or a validated disease registry<sup>23</sup>.
- *Timeframe*: Studies conducted from 01/01/1975 to 03/01/2018. 1975 represents the advent of ICD-9<sup>25</sup>.

We will exclude studies according to the following criteria:

- Data reported in systematic reviews unless we can identify the primary data, e.g. by contacting authors of the original source.
- Conference proceedings abstracts or studies not written in English where we are unable to obtain the meta-data from authors or full-length manuscript translations remain awaited, respectively.

Information Sources

Studies will be identified from the following sources

- *Electronic databases*: We will search 01/01/1975 to 03/01/2018 for studies meeting the inclusion criteria within the MEDLINE (OVID interface) and EMBASE (OVID interface) databases. The search strategies are outlined in table 1.

Ovid MEDLINE(R) Epub Ahead of Print, In-Process & Other Non-Indexed Citations, Ovid MEDLINE(R) Daily and Ovid MEDLINE(R) 1946 to 03 January 2018	Embase 1974 to 03 January 2018
1      Epilepsy, Complex Partial/ or Epilepsy, Reflex/ or Epilepsy, Absence/ or Drug Resistant Epilepsy/ or Epilepsy/ or Epilepsy, Rolandic/ or Epilepsy, Partial, Motor/ or Epilepsy, Benign Neonatal/ or Epilepsy, Tonic-Clonic/ or Epilepsy, Post-Traumatic/ or Epilepsy, Partial, Sensory/ or Epilepsy.mp. or Epilepsy, Temporal Lobe/ or Epilepsy, Frontal Lobe/ or Myoclonic Epilepsy, Juvenile/ or Epilepsy, Generalized/	1      reflex epilepsy/ or photosensitive epilepsy/ or grand mal epilepsy/ or epilepsy.mp. or drug resistant epilepsy/ or experimental epilepsy/ or severe myoclonic epilepsy in infancy/ or childhood absence epilepsy/ or benign childhood epilepsy/ or catamenial epilepsy/ or symptomatic epilepsy/ or startle epilepsy/ or generalized epilepsy/ or epilepsy/ or mesial temporal lobe epilepsy/ or rolandic epilepsy/ or traumatic epilepsy/ or myoclonic astatic epilepsy/ or temporal lobe epilepsy/ or intractable epilepsy/ or focal epilepsy/ or "seizure, epilepsy and convulsion"/ or myoclonus epilepsy/ or lateral temporal lobe epilepsy/ or frontal lobe epilepsy/
2      Databases, Factual/	2      factual database/ or data base/
3      "Reproducibility of Results"/	3      reproducibility/
4      Algorithms/ or algorithm*.mp.	4      algorithm/ or algorithm*.mp.
5      3 or 4	5      3 or 4
6      2 and 5	6      2 and 5
7      1 and 6	7      1 and 6
8      Administrative Claims, Healthcare/ or administrativ*.mp. or insurance data*.mp. or claims data*.mp. or Veterans Health Administration.mp.	8      "administrative claims (health care)"/ or administrative*.mp. or insurance data*.mp. or claims data*.mp. or Veterans Health Administration.mp.
9      administrat* data*.mp.	
10      routin* data*.mp.	



11	big data.mp.	9	administrat* data*.mp.
12	8 or 9 or 10 or 11	10	routin* data*.mp.
13	1 and 12	11	big data.mp.
14	Algorithms/ or algorithm*.mp.	12	8 or 9 or 10 or 11
15	"Sensitivity and Specificity"/ or "Predictive Value of Tests"/ or predictive value.mp.	13	1 and 12
16	positive* predict* value*.mp.	14	algorithm/ or algorithm*.mp.
17	negative* predict* value*.mp.	15	(specificity or sensitivity).mp. or "sensitivity and specificity"/ or positive* predict* value*.mp. or negative* predict* value*.mp.
18	sensitivity.mp.	16	area under the curve/ or area* under* curve*.mp.
19	specificity.mp.	17	roc curve/ or receiver operating characteristic/ or ROC curve*.mp.
20	area* under* curve*.mp. or Area Under Curve/	18	14 or 15 or 16 or 17
21	ROC Curve/ or ROC curve*.mp.	19	code*.mp. or "Read code"/
22	14 or 15 or 16 or 17 or 18 or 19 or 20 or 21	20	ICD-10.mp. or "International Classification of Diseases"/ or ICD-10/ or disease classification/ or ICD-9.mp. or ICD-9/
23	code*.mp.	21	19 or 20
24	(ICD-9 or ICD-10).mp. or "International Classification of Diseases"/ or Clinical Coding/ or read code*.mp.	22	18 and 21
25	23 or 24	23	1 and 22
26	22 and 25	24	validation study/ or validation process/ or validat*.mp. or validity/ or predictive validity/ or validity.mp.
27	1 and 26	25	electronic medical record/ or medical record/ or medical record*.mp. or medical case note*.mp.
28	(validat* or validity).mp. or Validation Studies/	26	Registries.mp. or register/
29	Medical Records/ or medical record*.mp. or medical case note*.mp.	27	25 or 26
30	electronic health records.mp. or Medical Records Systems, Computerized/ or Electronic Health Records/	28	24 and 27
31	Registries.mp. or Registries/	29	1 and 28
32	29 or 30 or 31	30	7 or 13 or 23 or 29
33	28 and 32	31	limit 30 to yr="1975 -Current"
34	1 and 33	32	animal/ not human/
35	7 or 13 or 27 or 34	33	31 not 32
36	limit 35 to yr="1975 -Current"		
37	Animals/ not Humans/		
38	36 not 37		

Table 1: Search strategies for MEDLINE and EMBASE

- *Reference lists:* We will also identify studies meeting the inclusion criteria from the reference lists of included studies and relevant reviews identified through the electronic database searches.
- *Conference proceedings:* For conference abstracts that appear to meet the inclusion criteria but do not have a full-length article published, we will contact authors directly to request meta-data.

### Study Records

*Data management:* Literature search results will be uploaded onto Review Manager 5, an internet-based software program that facilitates collaboration among reviewers. Citation titles and abstracts



will be uploaded. GM will then conduct an initial screen across individual studies to identify and exclude duplicate publications identified from the various electronic databases we will utilise by comparing author names, citation titles, sample sizes, outcomes used, and any other information held in abstract. All reviewer authors will have access to the systematic review process via the internet-based review software program and this will create an audit trail of studies included/excluded, data analysis steps, and subsequent manuscript revisions. All data will be within the management software and password-protected.

*Selection process:* Once duplicates have been excluded, two review authors (GM and KB) will independently screen the titles and abstracts yielded from the databases searches against the inclusion criteria. Where titles and abstracts indicate that a study may meet the inclusion criteria or where there is uncertainty about this, the full-length manuscripts will be downloaded and used to help decide. Where details in the manuscript are still insufficient for a decision to be made about eligibility, we will seek additional information from the study authors and automatically exclude studies where there is no response from authors after three weeks. We will record the reasons for excluding all excluded studies. The two review authors will compare their list of included and excluded trials and any disagreements will be resolved by mutual discussion and, where necessary, adjunction by a third reviewer (RC/SD/CS). Review authors will not be blind to the journal titles, study authors, or institutions.

*Data collection process and items:* Two review authors (GM and KB) will independently abstract data about the primary and secondary outcomes from included studies using the data collection tool shown below. The additional information extracted maps onto the items contained within the STARD Guidelines modified for epilepsy studies reporting diagnostic accuracy of administrative databases<sup>23</sup>. This will allow us to extract sufficient information to make a quality assessment of the completeness of included studies against the STARD Checklist<sup>23</sup>. Any disagreements in the contents of data abstraction will be resolved by mutual discussion and, where necessary, adjunction by a third reviewer (RC/SD/CS).

*Data collection tool*

- What is the study title?
- Who are the study authors?
- What is the year of study publication?
- What is the journal of publication?
- What country(s) was the study conducted in?
- Does the study explicitly identify as utilising “administrative data” (yes/no)?
  - o If not, how is this identified by the reviewer? – e.g. from descriptions given of the databases utilised and background knowledge about them or from correspondence with authors.
- Does the abstract provide a structured summary of study design, methods, results and conclusions (yes/no/unclear)?
- Does the introduction give a scientific and clinical background including intended use and clinical role of administrative data (yes/no/unclear)?
- Are the study objectives and hypotheses described (yes/no/unclear)?
  - o If so, what are they?
- What was the intended study sample size and how was it determined?
- What is the study design?
- Was a study/participant flow diagram used?
- What are the eligibility criteria for participation in the validation cohort (i.e. the cohort of patients to which the reference standard will be applied)?

- Where, when, and how were potentially eligible validation cohort participants identified? Include within this:
  - What is the name of the administrative database(s) on which the validation cohort was identified?
  - What are the setting and location of the administrative database(s) from which the validation cohort was identified? – e.g. is it primary care, secondary care, tertiary care, outpatient care, emergency care?
  - What are the names of any hospitals/organisations affiliated with or using the administrative database routinely?
  - What is the size of the administrative database(s) on which the validation cohort was identified? – i.e. how many people/records does it hold in total?
  - What were the epilepsy ICD-codes (or other disease classification system codes) used to identify the validation cohort within the administrative database(s)? – i.e. what are the diagnostic epilepsy codes that will be validated by the study?
  - What was the size of the validation cohort identified by the epilepsy codes? – i.e. give the number of participants identified by these diagnostic epilepsy codes.
  - Did the validation cohort include identifying a sample of people 1) without epilepsy? 2) with epilepsy “mimicker codes”?
    - If so, give details of the codes used and the number of participants for each of these groups.

Samples of people without epilepsy are often used to help calculate the specificity and NPV of an administrative database. “Mimicker codes” are often interrogated as the conditions may resemble epilepsy. These may include, for example, classical migraine (ICD-9 code 346.x, ICD-10 code G43.1), transient cerebral ischemia (TIA) (ICD-9 code 435, ICD-10 code G45), syncope (ICD-9 code 780.2, ICD-10 code R55), or convulsion (ICD-9 code 780.3, ICD-10 code R56.0 or R56.8); which are intended to be used for organic convulsions but not for epilepsy<sup>13</sup>.
  - What other information was obtained about an individual to help identify epilepsy on the administrative database? For example, describe if they linked an individual’s ICD epilepsy codes with the investigations they underwent (such as EEG) or the AED they were taking.
- What were the demographic and clinical characteristics of the validation cohort? – i.e. age, gender, type of epilepsy, comorbidities, and AEDs.
- Describe the reference standard. Include the following:
  - Name of the reference standard used;
  - What type of reference standard it was, e.g. clinical assessment, medical records, or validated disease registry;
  - Any rationale given for choosing this reference standard (if alternatives exist).
- What were the number, training and expertise of persons reading the reference standard?
  - If more than one person read the reference standard, what were the measures of consistency given? – e.g. kappa statistic.
- Describe any methods used to blind persons reading the reference standard to how the validation cohort were coded diagnostically on the administrative database, i.e. how a person reviewing the medical records diagnosis of an individual was made unaware of their administrative ICD diagnosis.
- Describe any methods used to blind persons reading the diagnostic administrative data codes to results of the reference standard diagnoses.

- What method was used to estimate the disease-identification accuracy of the administrative database? (i.e. sensitivity, specificity, PPV, NPV, or AUC).
  - o What were the results of this?
    - i.e. provide figures for these estimates and, where possible, also the individual true positive, false positive, true negative, and false negative figures;
    - Include the results of any cross-tabulation of the administrative data diagnoses results against the results of the reference standard diagnoses;
    - Describe the methods used and results of the measures used to estimate variability or precision of the diagnostic accuracy results (e.g. 95% confidence intervals).
- What method was used to compare the diagnostic accuracy of variables within the administrative database? – i.e. describe the method used to determine an optimal diagnostic algorithm.
  - o What were the results of this?
  - o Describe the methods used and results of the measures used to estimate variability or precision of the algorithm estimates (e.g. 95% confidence intervals)
- How were indeterminate or missing administrative database diagnoses or reference standard results handled?
- What were the time interval and any clinical interventions given between reference standard diagnosis and administrative database diagnosis?
- Describe any adverse events found from using the administrative database or reference standard.
- Summarise the study limitations described by the authors including any sources of potential bias, statistical uncertainty, generalisability limitation, and what they described as implications for practice.
- What were the study’s sources of funding?

Systematic review and meta-analysis outcomes

The primary outcome will be providing pooled disease-identification accuracy estimates of the included administrative databases using sensitivity, specificity, PPV, NPV, and AUC as the measures of accuracy. This will answer research question 1) How accurately do administrative data identify epilepsy within a population? We will provide a summary estimate of accuracy for ICD-9 and a summary estimate of accuracy for ICD-10 by pooling data across all studies using these classification systems provided there is no significant heterogeneity between studies as measured using the  $I^2$  statistic. The preferred estimators will be means with standard errors or medians with interquartile ranges, dependent on distribution. We will also quote the 95% confidence intervals for estimates. Where there is significant heterogeneity ( $I^2$  statistic > 50%), we will provide a descriptive analysis of the results and include ranges. Results will also be stratified by country and by age (adults or children less than 18 years of age).

The secondary outcome will be the optimal administrative data algorithms for correctly identifying epilepsy within a population. For this, we will assign a dummy variable with a binary 0 = “no” or 1 = “yes” category to participants having the following:

- A. a reference standard diagnosis of epilepsy (yes/no);
- B. an administrative diagnosis code for epilepsy (yes/no);
- C. multiple administrative epilepsy diagnoses codes over time (yes/no)
- D. having previously had an EEG (yes/no);
- E. having previously had a CT/MRI brain (yes/no);
- F. having previously had epilepsy surgery (yes/no);
- G. being on an individual AED (yes/no);
- H. being on 2 or more AEDs (yes/no).

Multivariable logistic regression models with A as the outcome variable and B to H individually and in combinations as the independent variables will be used in order to demonstrate the algorithm(s) best fitting the data across the included studies and to assess the significance of each variable's contribution to the model. The results of the logistic models will be displayed as sensitivity, specificity, PPV, NPV and AUC where possible, with 95% confidence intervals and measures of inter-study heterogeneity provided using the  $I^2$  statistic.

### Risk of bias analysis

We will utilise the Quality Assessment of Diagnostic Accuracy Studies 2 (QUADAS-2)<sup>27</sup> tool to assess risk of bias within and across studies, modified for studies validating administrative data. This is summarised in the table 2 and will be completed independently by two review authors (GM and KB) for each study with disagreements resolved by mutual discussion and, where necessary, adjunction by a third reviewer (RC/SD/CS). The tool consists of four key domains (see row 1) covering i) patient selection, ii) the administrative database, iii) the reference standard, and iv) flow of patients through the study and timing of the administrative database and reference standard. Each domain is assessed in terms of the risk of bias (graded as high, low, or unclear, see row 4) and the first three domains are also assessed in terms of concerns regarding applicability (see row 5). The description (see row 2) contains information used to support the risk of bias judgment. To help reach a judgement on the risk of bias, signalling questions are included (see row 3). These flag aspects of study design related to the potential for bias and aim to help reviewers make risk of bias judgements. If all signalling questions for a domain are answered “yes”, then risk of bias is judged “low”. If any signalling question is answered “no”, then risk of bias is judged “high”. If any signalling question is answered “unclear”, then risk of bias is judged “unclear”. Applicability sections are structured in a similar way to the bias sections, but do not include signalling questions. Review authors are asked to record the information on which the judgment of applicability is made and then to rate their concern that the study does not match the review question<sup>27 28</sup>.

DOMAIN	PATIENT SELECTION	ADMINISTRATIVE DATABASE	REFERENCE STANDARD	FLOW AND TIMING
Description	Describe methods of patient selection:  What is the review question?	Describe the administrative database and how it was used and interpreted:	Describe the reference standard and how it was conducted and interpreted:	Describe any patients in the validation cohort who were not found within the reference standard or who were excluded from cross-tabulation of the administrative data diagnoses results against the results of the reference standard diagnoses:  Describe the time interval and any interventions between administrative database diagnosis and reference standard diagnosis:
Signalling questions (yes/no/unclear)	Was a consecutive or random sample of patients enrolled?	Were the administrative database diagnosis results interpreted without knowledge of the results of the reference standard diagnosis?	Is the reference standard likely to correctly classify the epilepsy?	Was there an appropriate interval between administrative database diagnosis and reference standard diagnosis?
	Was a case-control design avoided?	If a diagnostic threshold was used, was it pre-specified?	Were the reference standard results interpreted without knowledge of the results of the administrative database diagnosis?	Did all patients receive a reference standard?
	Did the study avoid inappropriate exclusions?			Did all patients receive the same reference standard?
Risk of bias: High/low/unclear	Could the selection of patients have introduced bias?	Could the conduct or interpretation of the administrative database have introduced bias?	Could the reference standard, its conduct, or its interpretation have introduced bias?	Were all patients included in the analysis? Could the patient flow have introduced bias?
Concerns regarding applicability: High/low/unclear	Are there concerns that the included patients do not match the review question?	Are there concerns that the administrative database, its conduct, or interpretation differ from the review question?	Are there concerns that epilepsy, as defined by the reference standard, does not match the review question?	

Table 2: Risk of bias and applicability judgements in QUADAS-2

Upon completing the QUADAS-2 table, we will provide a risk of bias and applicability concerns graph demonstrating the review authors' judgements about each domain presented as percentages across included studies. We will also provide a risk of bias and applicability concerns summary demonstrating review authors' judgements about each domain for each included study. We will use

the Deek’s test<sup>29</sup> to interrogate for publication bias. This test is specifically designed for detecting funnel plot asymmetry in reviews of diagnostic studies<sup>28</sup>.

Confidence in cumulative evidence

We will utilise the Grading of Recommendations Assessment, Development, and Evaluation (GRADE) approach to assess strength of the body of evidence<sup>30</sup>. The GRADE system classifies the quality of evidence into one of four grades:

- 1. *High*: further research is very unlikely to change our confidence in the estimate of effect;
- 2. *Moderate*: further research is likely to have an important impact on our confidence in the estimate of effect and may change the estimate;
- 3. *Low*: further research is very likely to have an important impact on our confidence in the estimate of effect and is likely to change the estimate;
- 4. *Very low*: any estimate of effect is very uncertain<sup>31</sup>.

A judgement is made on the individual studies used to provide the pooled effect estimates and the quality of evidence is then downgraded by the cumulative presence of i) bias (see risk of bias analysis), ii) inconsistency (i.e. heterogeneity present on  $I^2$  statistic), iii) indirectness (i.e. high concerns regarding applicability, see table 2), iv) imprecision (small sample sizes, wide confidence intervals, inadequately powered studies), and v) publication bias (see Deek’s test<sup>29</sup> comments)<sup>32</sup>. GRADE classifications will be independently conducted by two review authors (GM and KB) with any disagreements resolved by mutual discussion and, where necessary, adjunction by a third reviewer (RC/SD/CS).

We will rate the completeness of reporting for each study out of 30 using the STARD 2015 checklist<sup>23</sup>. A score of 0-10, 11-20, 21-30 will indicate a low, moderate, and high quality of completeness of reporting, respectively.

Ethics and dissemination

Ethical approval is not required as primary data will not be collected. Results will be disseminated in peer-reviewed journals, conference presentations, and in press releases

References

1. Shamseer L, Moher D, Clarke M, et al. Preferred reporting items for systematic review and meta-analysis protocols (PRISMA-P) 2015: elaboration and explanation. *BMJ* 2015;349:g7647. doi: 10.1136/bmj.g7647

2. Moher D, Shamseer L, Clarke M, et al. Preferred reporting items for systematic review and meta-analysis protocols (PRISMA-P) 2015 statement. *Syst Rev* 2015;4:1. doi: 10.1186/2046-4053-4-1

3. Garratt E, Barnes H, Dibben C. Health administrative data: Exploring the potential for academic research. St Andrews: Administrative Data Liaison Service. 2010

4. Loke YK. Use of databases for clinical research. *Archives of Disease in Childhood* 2014;99(6):587-89. doi: 10.1136/archdischild-2013-304466

5. St Germaine-Smith C, Metcalfe A, Pringsheim T, et al. Recommendations for optimal ICD codes to study neurologic conditions: a systematic review. *Neurology* 2012;79(10):1049-55. doi: 10.1212/WNL.0b013e3182684707

6. Abraha I, Montedori A, Eusebi P, et al. The Current State of Validation of Administrative Healthcare Databases in Italy: A Systematic Review. *Pharmacoepidemiology and Drug Safety* 2012;21:400-00.

7. Traversa G, Bianchi C, Da Cas R, et al. Cohort study of hepatotoxicity associated with nimesulide and other non-steroidal anti-inflammatory drugs. *BMJ* 2003;327(7405):18-22. doi: 10.1136/bmj.327.7405.18



8. Hottes TS, Skowronski DM, Hiebert B, et al. Influenza vaccine effectiveness in the elderly based on administrative databases: change in immunization habit as a marker for bias. *PLoS One* 2011;6(7):e22618. doi: 10.1371/journal.pone.0022618
9. Lalmohamed A, Vestergaard P, Cooper C, et al. Timing of stroke in patients undergoing total hip replacement and matched controls: a nationwide cohort study. *Stroke* 2012;43(12):3225-9. doi: 10.1161/STROKEAHA.112.668509
10. Schneeweiss S, Avorn J. A review of uses of health care utilization databases for epidemiologic research on therapeutics. *J Clin Epidemiol* 2005;58(4):323-37. doi: 10.1016/j.jclinepi.2004.10.012
11. Krarup LH, Boysen G, Janjua H, et al. Validity of stroke diagnoses in a National Register of Patients. *Neuroepidemiology* 2007;28(3):150-4. doi: 10.1159/000102143
12. Kee VR, Gilchrist B, Granner MA, et al. A systematic review of validated methods for identifying seizures, convulsions, or epilepsy using administrative and claims data. *Pharmacoepidemiology and Drug Safety* 2012;21:183-93. doi: 10.1002/pds.2329
13. Jette N, Reid AY, Quan H, et al. How accurate is ICD coding for epilepsy? *Epilepsia* 2010;51(1):62-9. doi: 10.1111/j.1528-1167.2009.02201.x
14. Benson T. The history of the Read Codes: the inaugural James Read Memorial Lecture 2011. *Inform Prim Care* 2011;19(3):173-82.
15. Fonferko-Shadrach B, Lacey AS, White CP, et al. Validating epilepsy diagnoses in routinely collected data. *Seizure* 2017;52:195-98. doi: 10.1016/j.seizure.2017.10.008
16. Tu K, Wang M, Jaakkimainen RL, et al. Assessing the validity of using administrative data to identify patients with epilepsy. *Epilepsia* 2014;55(2):335-43. doi: 10.1111/epi.12506
17. Franchi C, Giussani G, Messina P, et al. Validation of healthcare administrative data for the diagnosis of epilepsy. *J Epidemiol Community Health* 2013;67(12):1019-24. doi: 10.1136/jech-2013-202528
18. Reid AY, St Germaine-Smith C, Liu M, et al. Development and validation of a case definition for epilepsy for use with administrative health data. *Epilepsy Res* 2012;102(3):173-9. doi: 10.1016/j.eplepsyres.2012.05.009
19. Parko K, Thurman DJ. Prevalence of epilepsy and seizures in the Navajo Nation 1998-2002. *Epilepsia* 2009;50(10):2180-5. doi: 10.1111/j.1528-1167.2009.02140.x
20. Pugh MJ, Van Cott AC, Cramer JA, et al. Trends in antiepileptic drug prescribing for older patients with new-onset epilepsy: 2000-2004. *Neurology* 2008;70(22 Pt 2):2171-8. doi: 10.1212/01.wnl.0000313157.15089.e6
21. Christensen J, Vestergaard M, Olsen J, et al. Validation of epilepsy diagnoses in the Danish National Hospital Register. *Epilepsy Res* 2007;75(2-3):162-70. doi: 10.1016/j.eplepsyres.2007.05.009
22. Tan M, Wilson I, Braganza V, et al. Development and validation of an epidemiologic case definition of epilepsy for use with routinely collected Australian health data. *Epilepsy Behav* 2015;51:65-72. doi: 10.1016/j.yebeh.2015.06.031
23. Bossuyt PM, Reitsma JB, Bruns DE, et al. STARD 2015: An Updated List of Essential Items for Reporting Diagnostic Accuracy Studies. *Clin Chem* 2015;61(12):1446-52. doi: 10.1373/clinchem.2015.246280
24. Moher D, Liberati A, Tetzlaff J, et al. Preferred reporting items for systematic reviews and meta-analyses: the PRISMA statement. *BMJ* 2009;339:b2535. doi: 10.1136/bmj.b2535
25. World Health Organisation. Classification of Diseases (ICD), 2016. <http://www.who.int/classifications/icd/en/>.
26. Florkowski CM. Sensitivity, specificity, receiver-operating characteristic (ROC) curves and likelihood ratios: communicating the performance of diagnostic tests. *Clin Biochem Rev* 2008;29 Suppl 1:S83-7.
27. Whiting PF, Rutjes AW, Westwood ME, et al. QUADAS-2: a revised tool for the quality assessment of diagnostic accuracy studies. *Ann Intern Med* 2011;155(8):529-36. doi: 10.7326/0003-4819-155-8-201110180-00009
28. Leeflang MM, Deeks JJ, Takwoingi Y, et al. Cochrane diagnostic test accuracy reviews. *Syst Rev* 2013;2:82. doi: 10.1186/2046-4053-2-82

29. Deeks JJ, Macaskill P, Irwig L. The performance of tests of publication bias and other sample size effects in systematic reviews of diagnostic test accuracy was assessed. *J Clin Epidemiol* 2005;58(9):882-93. doi: 10.1016/j.jclinepi.2005.01.016

30. Schünemann H, Bro ek J, Oxman A. GRADE handbook for grading quality of evidence and strength of recommendation. 2009; 3.2. <http://www.ccims.net/gradepr>.

31. Mbizvo GK, Dixon P, Hutton JL, et al. Levetiracetam add-on for drug-resistant focal epilepsy: an updated Cochrane Review. *Cochrane Database Syst Rev* 2012;9:CD001901. doi: 10.1002/14651858.CD001901.pub2

32. Singh S, Chang SM, Matchar DB, et al. Chapter 7: grading a body of evidence on diagnostic tests. *J Gen Intern Med* 2012;27 Suppl 1:S47-55. doi: 10.1007/s11606-012-2021-9

**Authors' contributions**

GKM and RFMC conceived the idea for the protocol and made the main contribution to planning and preparation of timelines for its completion. GKM and KB put together and tested various search strategies for the protocol and, after consultation with CRS, had these reviewed and approved by Marshall Dozier at the University of Edinburgh library, who's support we acknowledge. GKM and RC planned the data extraction and statistical analysis, as well as of risk of bias, quality of evidence, and completeness of reporting assessments. GKM designed the tables and wrote the first draft of the manuscript, which was then reviewed and amended by KB, CRS, SD and RFMC. GKM, KB, CRS, SD and RFMC then approved the final written manuscript. RFMC is the guarantor for the work.

**Funding statement**

This work was supported by Epilepsy Research UK (R44007) and the Juliet Bergqvist Memorial Fund.

**Competing interests statement**

The authors declare no competing interests.

**Additional information**

This protocol follows the Preferred Reporting Items for Systematic review and Meta-Analysis Protocols (PRISMA-P) guidelines<sup>1, 2</sup>. Any future amendments of the protocol will be listed in this section along with a date, description, and rational for each amendment.



# BMJ Open

## The Accuracy and Utility of Using Administrative Healthcare Databases to Identify People with Epilepsy: A Protocol for a Systematic Review and Meta-Analysis

Journal:	<i>BMJ Open</i>
Manuscript ID	bmjopen-2017-020824.R1
Article Type:	Protocol
Date Submitted by the Author:	12-Mar-2018
Complete List of Authors:	Mbizvo, Gashirai; University of Edinburgh Centre for Clinical Brain Sciences, Muir Maxwell Epilepsy Centre, Bennett, Kyle; University of Edinburgh Centre for Clinical Brain Sciences, Muir Maxwell Epilepsy Centre, Simpson, Colin; Victoria University of Wellington, Faculty of Health,; University of Edinburgh School of Molecular Genetic and Population Health Sciences, Usher Institute of Population Health Sciences and Informatics, Duncan, Susan; Western General Hospital, Department of Clinical Neurosciences, Chin, Richard; University of Edinburgh Centre for Clinical Brain Sciences, Muir Maxwell Epilepsy Centre,
<b>Primary Subject Heading</b>:	Neurology
Secondary Subject Heading:	Health informatics, Neurology, Diagnostics, Epidemiology
Keywords:	Epilepsy < NEUROLOGY, Factual database, Administrative claims, Algorithms, Validation studies

SCHOLARONE™  
Manuscripts

**Title: The Accuracy and Utility of Using Administrative Healthcare Databases to Identify People with Epilepsy: A Protocol for a Systematic Review and Meta-Analysis**

**Authors:** Gashirai K Mbizvo<sup>A</sup>, Kyle Bennett<sup>A</sup>, Colin R Simpson<sup>B,C</sup>, Susan E Duncan<sup>D</sup>, Richard FM Chin<sup>A,E</sup>

<sup>A</sup> University of Edinburgh Centre for Clinical Brain Sciences, Muir Maxwell Epilepsy Centre, Edinburgh, UK

<sup>B</sup> Victoria University of Wellington, Faculty of Health, Wellington, NZ

<sup>C</sup> University of Edinburgh School of Molecular Genetic and Population Health Sciences, Usher Institute of Population Health Sciences and Informatics, Edinburgh, UK

<sup>D</sup> Western General Hospital, Department of Clinical Neurosciences, Edinburgh, UK

<sup>E</sup> Royal Hospital for Sick Children, Edinburgh, UK

**Keywords:** Epilepsy, factual database, administrative claims, algorithms, validation studies

**Corresponding Author**

Dr Gashirai K Mbizvo, *University of Edinburgh Centre for Clinical Brain Sciences, Muir Maxwell Epilepsy Centre, Edinburgh, UK*

*Tel: +44 (0) 131 536 0801, Fax: N/A*

*email: [gashirai.mbizvo@ed.ac.uk](mailto:gashirai.mbizvo@ed.ac.uk)*

**Word count:**

Abstract – 299 words,

Body – 3,860 words,

Tables – 2,

Illustrations/figures – 0,

References – 32.

## Abstract

**Introduction:** In an increasingly digital age for healthcare around the world, administrative data have become rich and accessible tools for potentially identifying and monitoring population trends in diseases including epilepsy. However, it remains unclear (1) how accurate administrative data are at identifying epilepsy within a population, and (2) the optimal algorithms needed for administrative data to correctly identify people with epilepsy within a population. To address this knowledge gap, we will conduct a novel systematic review of all identified studies validating administrative healthcare data in epilepsy identification. We provide here a protocol that will outline the methods and analyses planned for the systematic review.

**Methods and analysis:** The systematic review described in this protocol will be conducted to follow the Preferred Reporting Items for Systematic Reviews and Meta-Analyses (PRISMA) guidelines. MEDLINE and EMBASE will be searched for studies validating administrative data in epilepsy published between 01/01/1975 and 03/01/2018. Included studies will validate the International Classification of Disease (ICD) 9th revision onwards (ICD-9 code 345 and ICD-10 codes G40-G41). The primary outcome will be providing pooled estimates of accuracy for identifying epilepsy within the administrative databases validated using sensitivity, specificity, positive and negative predictive values, and area under the receiver operating characteristic curves. Heterogeneity will be assessed using the  $I^2$  statistic and descriptive analyses used where this is present. The secondary outcome will be the optimal administrative data algorithms for correctly identifying epilepsy. These will be identified using multivariable logistic regression models. 95% confidence intervals will be quoted throughout. We will make an assessment of risk of bias, quality of evidence, and completeness of reporting for included studies.

**Ethics and dissemination:** Ethical approval is not required as primary data will not be collected. Results will be disseminated in peer-reviewed journals, conference presentations, and in press releases.

**PROSPERO registration:** CRD42017081212.

## Strengths and limitations of this study

### Strengths

- The protocol describes what will be the first systematic review to conduct a worldwide assessment of the accuracy of administrative data in identifying epilepsy and the optimal disease-identification algorithms.
- This protocol also describes what will be the first systematic review to make an assessment of risk of bias, quality of evidence, and completeness of reporting for studies validating administrative healthcare data in epilepsy identification.

### Limitations

- The review described in this protocol will be limited to assessing the use of administrative data in diagnosing epilepsy within observational studies, which are more prone to bias than randomised controlled trials.
- A systematic review of the diagnostic accuracy of administrative data within randomised controlled trials in epilepsy remains to be completed, and is out of the scope of the current review.

## Introduction

Administrative healthcare databases are electronic data sources that consist of demographic, diagnostic, and clinical information routinely collected about patients when they use a healthcare service<sup>1</sup>. They are often national and mandatory, and therefore they have the potential to provide a

relatively cheap, widely-available, and less intrusive resource for medical research<sup>1</sup>. However, the accuracy of the information held in an administrative database needs to be validated before such use can be made. This is because the administrative data were not originally collected for research, but for other purposes such as assisting in health insurance claims. The clinical information held may therefore lack the rigor in accuracy that might be expected in scientifically collected data. Furthermore, the data may be limited by inaccurate or incomplete hospital discharge letters or clinical coding transcription errors<sup>2</sup>.

The validation of administrative data involves comparing the diagnostic codes held within the administrative database against a reference standard (such as medical records) in order to quantify the number of instances in which the administrative diagnosis made matches the diagnosis in the reference standard (deemed to be the true diagnosis). In this way, the administrative database can be handled like a diagnostic test and measures of disease-identification accuracy calculated. These measures usually include the sensitivity, specificity, and the positive or negative predictive value (PPV or NPV, respectively). Optimal disease-identification algorithms can also be determined by making relative comparisons of predictive values after adding in data from other variables recorded in an administrative database, such as drug combinations, investigations, and procedures.

There are many administrative databases worldwide in which the process of validation has been successfully performed for many diagnostic codes<sup>3-9</sup>. There are also examples of where the results of these have been pooled successfully into systematic review to increase confidence in the estimates made and scrutinise the quality of evidence, and this has led to changes in practice<sup>4</sup>. There has been limited systematic review of the validation of administrative databases in capturing epilepsy as a diagnosis. The only systematic review<sup>10</sup> on this subject included only studies from the USA or Canada and therefore excluded 121 studies because the data sources were not from these two countries. Furthermore, the eleven studies included were published between 2000 and 2010, making the conclusions nearly a decade old now. With health informatics now at the forefront of epidemiological disease surveillance it is important to have an update on performance of the administrative disease-identification codes. Only one of the included studies evaluated the performance of the ICD-10 system in capturing epilepsy within administrative datasets<sup>11</sup> –the remainder evaluated the older ICD-9 system<sup>10</sup>. The review also made no assessment of risk of bias, quality of evidence, and completeness of reporting for included studies. This limits the confidence with which conclusions can be interpreted. There is now need for a more contemporary systematic review of the validation of administrative databases in capturing epilepsy. This should include evaluating performance of the ICD-10 system, as well as other non-ICD disease classification systems used, such as Read Codes in the United Kingdom<sup>12 13</sup>. The review should include studies from anywhere in the world in order to give clinicians and researchers a representative picture of the performance of administrative data in capturing epilepsy as a diagnosis and in order to allow more generalisable diagnostic algorithms to be suggested. Furthermore, the review should make an assessment of risk of bias, quality of evidence, and completeness of reporting for included studies. These are the aims of the proposed systematic review described in this protocol. This will help researchers and clinicians better understand the accuracy of global estimates for incidence, prevalence, and population characteristics in epilepsy – which have largely been made using administrative data.

Aims and objectives:

The study hypotheses are:

1. Administrative data can correctly identify people with epilepsy within a population with a high degree of accuracy. We predict the PPV to be above 80%.

2. The optimal disease-identification algorithms for epilepsy within administrative datasets take into account diagnoses, investigations, and drug combinations<sup>11 14-20</sup>.

The aim of the systematic review is to quantify the disease-identification accuracy and algorithm performance of administrative healthcare data in epilepsy. To this end, the research questions are:

1. How accurately do administrative data identify epilepsy within a population? –as measured by sensitivity, specificity, PPV, NPV, or area under the receiver operating characteristic curve (AUC) analysis (which are the approved accuracy measures described in the Standards for Reporting Diagnostic Accuracy (STARD) statement<sup>21</sup>). This will be the primary outcome.
2. What are the optimal administrative data algorithms for correctly identifying epilepsy within a population? This will be the secondary outcome.

A preliminary feasibility search of the MEDLINE database via PubMed identifies at least nine studies validating diagnostic epilepsy codes held within administrative databases around the world that could be utilised to answer these research questions<sup>11 13-20</sup>.

### Methods and analysis

This protocol follows the Preferred Reporting Items for Systematic review and Meta-Analysis Protocols (PRISMA-P) checklist<sup>22 23</sup>. The systematic review will follow the PRISMA checklist<sup>24</sup>.

#### Eligibility criteria

We will include studies according to the following criteria:

- *Language*: There will be no language restriction on full-length articles although abstracts will need to be in English to allow the authors to screen them. We will seek translations for full length articles not written in English that appear eligible in abstract. These will remain in the section for “studies still awaiting classification” and will feature in subsequent updates to the review if not translated by the time of initial publication.
- *Setting*: There will be no restrictions by study location worldwide. Where possible, we will show pooled accuracy and best algorithm data for administrative datasets from individual countries in addition to pooled global estimates of these measures.
- *Databases*: The data sources will be routine administrative healthcare databases. This means that the data should have been routinely and passively collected without an *a priori* research question<sup>4</sup>. We will include databases containing diagnostic codes for epilepsy classified on the ICD system – where we will restrict this to studies using the ninth ICD revision onwards (ICD-9, active from 1975 to 1994)<sup>25</sup>. This is because although the ICD system is currently in its tenth revision (ICD-10, active from 1994 to present) and this the primary coding system utilised by many countries around the world, a significant proportion of countries, particularly developing ones, still use the ICD-9 system<sup>11</sup>. ICD-9 code 345 and ICD-10 codes G40 and or G41 will be used to identify epilepsy diagnoses. We will provide summary measures of accuracy and best algorithms for any other disease classification systems used in studies separately, e.g. the primary care Read Code system used in the UK<sup>12 13</sup>.
- *Study design*: Prospective or retrospective observational studies including cohort or case-control designs that are community-based/population-based or primary/secondary/tertiary care-based and have utilised administrative databases.
- *Participants*: People with epilepsy of all ages. Where available, we will additionally report data for adults and children separately.
- *Observations and outcomes*: Studies will need to employ a validation process for diagnostic epilepsy codes, i.e. estimate the disease-identification accuracy of the epilepsy codes held within the database using sensitivity, specificity, PPV, NPV, or AUC analysis<sup>21</sup>. In this, true positives (TPs) and false negatives (FNs) will be considered as the patient has the disease and the

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

administrative diagnosis is positive or negative, respectively. True negatives (TNs) and false positives (FPs) will be considered as the patient does not have the disease and the administrative diagnosis is negative or positive, respectively. Sensitivity will be considered as the ability of the administrative database to correctly identify those patients with the disease (TP/(TP + FN)). Specificity will be considered as the ability of the administrative database to correctly identify those patients without the disease (TN/(TN + FP)). PPV will be considered as how likely it is that a patient has the disease if the administrative diagnosis is positive (TP/(TP + FP)). NPV will be considered as how likely it is that this patient does not have the disease if the administrative diagnosis is negative (TN/(TN + FN)). AUC analysis will be considered as where TP and FP are plotted against each other in a perfect correlation as reference to show poor test accuracy, then accuracy measured as the area under the curve created by plotting the actual values against each other<sup>26</sup>.

- Studies may also use diagnostic as well as other variables (e.g. admissions, drugs or investigations) to calculate optimal disease-identification algorithms for epilepsy within the database. Studies will need to provide a clinical reference standard. An appropriate clinical reference standard will be medical records, clinical assessment, or a validated disease registry<sup>21</sup>.
- *Timeframe*: Studies conducted from 01/01/1975 to 03/01/2018. 1975 represents the advent of ICD-9<sup>25</sup>.

We will exclude studies according to the following criteria:

- Data reported in systematic reviews unless we can identify the primary data, e.g. by contacting authors of the original source.
- Conference proceedings abstracts or studies not written in English where we are unable to obtain the meta-data from authors or full-length manuscript translations remain awaited, respectively.

Information Sources

Studies will be identified from the following sources

- *Electronic databases*: We will search 01/01/1975 to 03/01/2018 for studies meeting the inclusion criteria within the MEDLINE (OVID interface) and EMBASE (OVID interface) databases. The search strategies are outlined in table 1.

Ovid MEDLINE(R) Epub Ahead of Print, In-Process & Other Non-Indexed Citations, Ovid MEDLINE(R) Daily and Ovid MEDLINE(R) 1946 to 03 January 2018	Embase 1974 to 03 January 2018
1      Epilepsy, Complex Partial/ or Epilepsy, Reflex/ or Epilepsy, Absence/ or Drug Resistant Epilepsy/ or Epilepsy/ or Epilepsy, Rolandic/ or Epilepsy, Partial, Motor/ or Epilepsy, Benign Neonatal/ or Epilepsy, Tonic-Clonic/ or Epilepsy, Post-Traumatic/ or Epilepsy, Partial, Sensory/ or Epilepsy.mp. or Epilepsy, Temporal Lobe/ or Epilepsy, Frontal Lobe/ or Myoclonic Epilepsy, Juvenile/ or Epilepsy, Generalized/	1      reflex epilepsy/ or photosensitive epilepsy/ or grand mal epilepsy/ or epilepsy.mp. or drug resistant epilepsy/ or experimental epilepsy/ or severe myoclonic epilepsy in infancy/ or childhood absence epilepsy/ or benign childhood epilepsy/ or catamenial epilepsy/ or symptomatic epilepsy/ or startle epilepsy/ or generalized epilepsy/ or epilepsy/ or mesial temporal lobe epilepsy/ or rolandic epilepsy/ or traumatic epilepsy/ or myoclonic astatic epilepsy/ or temporal lobe epilepsy/ or intractable epilepsy/ or focal epilepsy/ or "seizure, epilepsy and convulsion"/ or myoclonus epilepsy/ or lateral temporal lobe epilepsy/ or frontal lobe epilepsy/
2      Databases, Factual/	2      factual database/ or data base/
3      "Reproducibility of Results"/	3      reproducibility/
4      Algorithms/ or algorithm*.mp.	4      algorithm/ or algorithm*.mp.
5      3 or 4	5      3 or 4
6      2 and 5	6      2 and 5
7      1 and 6	7      1 and 6
8      Administrative Claims, Healthcare/ or administrativ*.mp. or insurance data*.mp. or claims data*.mp. or Veterans Health Administration.mp.	8      "administrative claims (health care)"/ or
9      administrat* data*.mp.	



10	routin* data*.mp.	administrative*.mp. or insurance data*.mp. or claims data*.mp. or Veterans Health Administration.mp.
11	big data.mp.	9 administrat* data*.mp.
12	8 or 9 or 10 or 11	10 routin* data*.mp.
13	1 and 12	11 big data.mp.
14	Algorithms/ or algorithm*.mp.	12 8 or 9 or 10 or 11
15	"Sensitivity and Specificity"/ or "Predictive Value of Tests"/ or predictive value.mp.	13 1 and 12
16	positive* predict* value*.mp.	14 algorithm/ or algorithm*.mp.
17	negative* predict* value*.mp.	15 (specificity or sensitivity).mp. or "sensitivity and specificity"/ or positive* predict* value*.mp. or negative* predict* value*.mp.
18	sensitivity.mp.	16 area under the curve/ or area* under* curve*.mp.
19	specificity.mp.	17 roc curve/ or receiver operating characteristic/ or ROC curve*.mp.
20	area* under* curve*.mp. or Area Under Curve/	18 14 or 15 or 16 or 17
21	ROC Curve/ or ROC curve*.mp.	19 code*.mp. or "Read code"/
22	14 or 15 or 16 or 17 or 18 or 19 or 20 or 21	20 ICD-10.mp. or "International Classification of Diseases"/ or ICD-10/ or disease classification/ or ICD-9.mp. or ICD-9/
23	code*.mp.	21 19 or 20
24	(ICD-9 or ICD-10).mp. or "International Classification of Diseases"/ or Clinical Coding/ or read code*.mp.	22 18 and 21
25	23 or 24	23 1 and 22
26	22 and 25	24 validation study/ or validation process/ or validat*.mp. or validity/ or predictive validity/ or validity.mp.
27	1 and 26	25 electronic medical record/ or medical record/ or medical record*.mp. or medical case note*.mp.
28	(validat* or validity).mp. or Validation Studies/	26 Registries.mp. or register/
29	Medical Records/ or medical record*.mp. or medical case note*.mp.	27 25 or 26
30	electronic health records.mp. or Medical Records Systems, Computerized/ or Electronic Health Records/	28 24 and 27
31	Registries.mp. or Registries/	29 1 and 28
32	29 or 30 or 31	30 7 or 13 or 23 or 29
33	28 and 32	31 limit 30 to yr="1975 -Current"
34	1 and 33	32 animal/ not human/
35	7 or 13 or 27 or 34	33 31 not 32
36	limit 35 to yr="1975 -Current"	
37	Animals/ not Humans/	
38	36 not 37	

Table 1: Search strategies for MEDLINE and EMBASE

- *Reference lists:* We will also identify studies meeting the inclusion criteria from the reference lists of included studies and relevant reviews identified through the electronic database searches.
- *Conference proceedings:* For conference abstracts that appear to meet the inclusion criteria but do not have a full-length article published, we will contact authors directly to request meta-data.

## Study Records



*Data management:* Literature search results will be uploaded onto Review Manager 5, an internet-based software program that facilitates collaboration among reviewers. Citation titles and abstracts will be uploaded. GM will then conduct an initial screen across individual studies to identify and exclude duplicate publications identified from the various electronic databases we will utilise by comparing author names, citation titles, sample sizes, outcomes used, and any other information held in abstract. All reviewer authors will have access to the systematic review process via the internet-based review software program and this will create an audit trail of studies included/excluded, data analysis steps, and subsequent manuscript revisions. All data will be within the management software and password-protected.

*Selection process:* Once duplicates have been excluded, two review authors (GM and KB) will independently screen the titles and abstracts yielded from the databases searches against the inclusion criteria. Where titles and abstracts indicate that a study may meet the inclusion criteria or where there is uncertainty about this, the full-length manuscripts will be downloaded and used to help decide. Where details in the manuscript are still insufficient for a decision to be made about eligibility, we will seek additional information from the study authors and automatically exclude studies where there is no response from authors after three weeks. We will record the reasons for excluding all excluded studies. The two review authors will compare their list of included and excluded trials and any disagreements will be resolved by mutual discussion and, where necessary, adjunction by a third reviewer (RC/SD/CS). Review authors will not be blind to the journal titles, study authors, or institutions.

*Data collection process and items:* Two review authors (GM and KB) will independently abstract data about the primary and secondary outcomes from included studies using the data collection tool shown below. The additional information extracted maps onto the items contained within the STARD Guidelines modified for epilepsy studies reporting diagnostic accuracy of administrative databases<sup>21</sup>. This will allow us to extract sufficient information to make a quality assessment of the completeness of included studies against the STARD Checklist<sup>21</sup>. Any disagreements in the contents of data abstraction will be resolved by mutual discussion and, where necessary, adjunction by a third reviewer (RC/SD/CS).

*Data collection tool*

- What is the study title?
- Who are the study authors?
- What is the year of study publication?
- What is the journal of publication?
- What country(s) was the study conducted in?
- Does the study explicitly identify as utilising “administrative data” (yes/no)?
  - o If not, how is this identified by the reviewer? – e.g. from descriptions given of the databases utilised and background knowledge about them or from correspondence with authors.
- Does the abstract provide a structured summary of study design, methods, results and conclusions (yes/no/unclear)?
- Does the introduction give a scientific and clinical background including intended use and clinical role of administrative data (yes/no/unclear)?
- Are the study objectives and hypotheses described (yes/no/unclear)?
  - o If so, what are they?
- What was the intended study sample size and how was it determined?
- What is the study design?
- Was a study/participant flow diagram used?

- What are the eligibility criteria for participation in the validation cohort (i.e. the cohort of patients to which the reference standard will be applied)?
- Where, when, and how were potentially eligible validation cohort participants identified? Include within this:
  - What is the name of the administrative database(s) on which the validation cohort was identified?
  - What are the setting and location of the administrative database(s) from which the validation cohort was identified? – e.g. is it primary care, secondary care, tertiary care, outpatient care, emergency care?
  - What are the names of any hospitals/organisations affiliated with or using the administrative database routinely?
  - What is the size of the administrative database(s) on which the validation cohort was identified? – i.e. how many people/records does it hold in total?
  - What were the epilepsy ICD-codes (or other disease classification system codes) used to identify the validation cohort within the administrative database(s)? – i.e. what are the diagnostic epilepsy codes that will be validated by the study?
  - What was the size of the validation cohort identified by the epilepsy codes? – i.e. give the number of participants identified by these diagnostic epilepsy codes.
  - Did the validation cohort include identifying a sample of people 1) without epilepsy? 2) with epilepsy “mimicker codes”?
    - If so, give details of the codes used and the number of participants for each of these groups.

Samples of people without epilepsy are often used to help calculate the specificity and NPV of an administrative database. “Mimicker codes” are often interrogated as the conditions may resemble epilepsy. These may include, for example, classical migraine (ICD-9 code 346.x, ICD-10 code G43.1), transient cerebral ischemia (TIA) (ICD-9 code 435, ICD-10 code G45), syncope (ICD-9 code 780.2, ICD-10 code R55), or convulsion (ICD-9 code 780.3, ICD-10 code R56.0 or R56.8); which are intended to be used for organic convulsions but not for epilepsy<sup>11</sup>.
  - What other information was obtained about an individual to help identify epilepsy on the administrative database? For example, describe if they linked an individual’s ICD epilepsy codes with the investigations they underwent (such as EEG) or the AED they were taking.
- What were the demographic and clinical characteristics of the validation cohort? – i.e. age, gender, type of epilepsy, comorbidities, and AEDs.
- Describe the reference standard. Include the following:
  - Name of the reference standard used;
  - What type of reference standard it was, e.g. clinical assessment, medical records, or validated disease registry;
  - Any rationale given for choosing this reference standard (if alternatives exist).
- What were the number, training and expertise of persons reading the reference standard?
  - If more than one person read the reference standard, what were the measures of consistency given? – e.g. kappa statistic.
- Describe any methods used to blind persons reading the reference standard to how the validation cohort were coded diagnostically on the administrative database, i.e. how a person reviewing the medical records diagnosis of an individual was made unaware of their administrative ICD diagnosis.

- Describe any methods used to blind persons reading the diagnostic administrative data codes to results of the reference standard diagnoses.
- What method was used to estimate the disease-identification accuracy of the administrative database? (i.e. sensitivity, specificity, PPV, NPV, or AUC).
  - o What were the results of this?
    - i.e. provide figures for these estimates and, where possible, also the individual true positive, false positive, true negative, and false negative figures;
    - Include the results of any cross-tabulation of the administrative data diagnoses results against the results of the reference standard diagnoses;
    - Describe the methods used and results of the measures used to estimate variability or precision of the diagnostic accuracy results (e.g. 95% confidence intervals).
- What method was used to compare the diagnostic accuracy of variables within the administrative database? – i.e. describe the method used to determine an optimal diagnostic algorithm.
  - o What were the results of this?
  - o Describe the methods used and results of the measures used to estimate variability or precision of the algorithm estimates (e.g. 95% confidence intervals)
- How were indeterminate or missing administrative database diagnoses or reference standard results handled?
- What were the time interval and any clinical interventions given between reference standard diagnosis and administrative database diagnosis?
- Describe any adverse events found from using the administrative database or reference standard.
- Summarise the study limitations described by the authors including any sources of potential bias, statistical uncertainty, generalisability limitation, and what they described as implications for practice.
- What were the study’s sources of funding?

Systematic review and meta-analysis outcomes

The primary outcome will be providing pooled disease-identification accuracy estimates of the included administrative databases using sensitivity, specificity, PPV, NPV, and AUC as the measures of accuracy. This will answer research question 1) How accurately do administrative data identify epilepsy within a population? We will provide a summary estimate of accuracy for ICD-9 and a summary estimate of accuracy for ICD-10 by pooling data across all studies using these classification systems provided there is no significant heterogeneity between studies as measured using the  $I^2$  statistic. The preferred estimators will be means with standard errors or medians with interquartile ranges, dependent on distribution. We will also quote the 95% confidence intervals for estimates. Where there is significant heterogeneity ( $I^2$  statistic > 50%), we will provide a descriptive analysis of the results and include ranges. Just as there may be heterogeneity introduced by making comparisons across different trial designs, we expect there to be heterogeneity introduced by making comparisons across different healthcare systems. This is because there are likely to be differences in the accuracy of administrative healthcare data owing to differences in coding practice and/or the quality of reference standards between different healthcare systems. Therefore, we will also conduct subgroup analysis in which diagnostic accuracy results are pooled together for studies that have used the same or similar healthcare systems, e.g. the National Health Service in the UK, Veterans Health Administration in the US, healthcare systems with geographical overlap, and private- versus state-funded healthcare systems. We will also create subgroups in which results are pooled together within the following study design groups: prospective cohorts; retrospective cohorts; case-control studies; primary care studies; secondary care studies; tertiary care studies; paediatric studies (age <18 years); adult studies; and studies from the same country. Differences in the results for each subgroup may

provide an important guide for future studies in the field, and they may also help to explain any statistical heterogeneity seen.

The secondary outcome will be the optimal administrative data algorithms for correctly identifying epilepsy within a population. For this, we will assign a dummy variable with a binary 0 = “no” or 1 = “yes” category to participants having the following:

- A. a reference standard diagnosis of epilepsy (yes/no);
- B. an administrative diagnosis code for epilepsy (yes/no);
- C. multiple administrative epilepsy diagnoses codes over time (yes/no)
- D. having previously had an EEG (yes/no);
- E. having previously had a CT/MRI brain (yes/no);
- F. having previously had epilepsy surgery (yes/no);
- G. being on an individual AED (yes/no);
- H. being on 2 or more AEDs (yes/no).

Multivariable logistic regression models with A as the outcome variable and B to H individually and in combinations as the independent variables will be used in order to demonstrate the algorithm(s) best fitting the data across the included studies and to assess the significance of each variable's contribution to the model. The results of the logistic models will be displayed as sensitivity, specificity, PPV, NPV and AUC where possible, with 95% confidence intervals and measures of inter-study heterogeneity provided using the  $I^2$  statistic.

#### Risk of bias analysis

We will utilise the Quality Assessment of Diagnostic Accuracy Studies 2 (QUADAS-2)<sup>27</sup> tool to assess risk of bias within and across studies, modified for studies validating administrative data. This is summarised in the table 2 and will be completed independently by two review authors (GM and KB) for each study with disagreements resolved by mutual discussion and, where necessary, adjunction by a third reviewer (RC/SD/CS). The tool consists of four key domains (see row 1) covering i) patient selection, ii) the administrative database, iii) the reference standard, and iv) flow of patients through the study and timing of the administrative database and reference standard. Each domain is assessed in terms of the risk of bias (graded as high, low, or unclear, see row 4) and the first three domains are also assessed in terms of concerns regarding applicability (see row 5). The description (see row 2) contains information used to support the risk of bias judgment. To help reach a judgement on the risk of bias, signalling questions are included (see row 3). These flag aspects of study design related to the potential for bias and aim to help reviewers make risk of bias judgements. If all signalling questions for a domain are answered “yes”, then risk of bias is judged “low”. If any signalling question is answered “no”, then risk of bias is judged “high”. If any signalling question is answered “unclear”, then risk of bias is judged “unclear”. Applicability sections are structured in a similar way to the bias sections, but do not include signalling questions. Review authors are asked to record the information on which the judgment of applicability is made and then to rate their concern that the study does not match the review question<sup>27 28</sup>.

DOMAIN	PATIENT SELECTION	ADMINISTRATIVE DATABASE	REFERENCE STANDARD	FLOW AND TIMING
Description	Describe methods of patient selection:  What is the review question?	Describe the administrative database and how it was used and interpreted:  Where available, include comment on how coding was done, by whom, and whether or not there was reimbursement for coding.	Describe the reference standard and how it was conducted and interpreted:  Where available, include comment on quality of the reference standard, including the level of experience of clinicians making the diagnosis, access to diagnostic tests such as electroencephalography and telemetry, and the thresholds/criteria used to make a diagnosis of epilepsy.	Describe any patients in the validation cohort who were not found within the reference standard or who were excluded from cross-tabulation of the administrative data diagnoses results against the results of the reference standard diagnoses:  Describe the time interval and any interventions between administrative database diagnosis and reference standard diagnosis:

Signalling questions (yes/no/unclear)	Was a consecutive or random sample of patients enrolled?	Were the administrative database diagnosis results interpreted without knowledge of the results of the reference standard diagnosis?	Is the reference standard likely to correctly classify the epilepsy?	Was there an appropriate interval between administrative database diagnosis and reference standard diagnosis?
	Was a case-control design avoided?	If a diagnostic threshold was used, was it pre-specified?	Were the reference standard results interpreted without knowledge of the results of the administrative database diagnosis?	Did all patients receive a reference standard?
	Did the study avoid inappropriate exclusions?			Did all patients receive the same reference standard?
				Were all patients included in the analysis?
Risk of bias: High/low/unclear	Could the selection of patients have introduced bias?	Could the conduct or interpretation of the administrative database have introduced bias?	Could the reference standard, its conduct, or its interpretation have introduced bias?	Could the patient flow have introduced bias?
Concerns regarding applicability: High/low/unclear	Are there concerns that the included patients do not match the review question?	Are there concerns that the administrative database, its conduct, or interpretation differ from the review question?	Are there concerns that epilepsy, as defined by the reference standard, does not match the review question?	

Table 2: Risk of bias and applicability judgements in QUADAS-2

Upon completing the QUADAS-2 table, we will provide a risk of bias and applicability concerns graph demonstrating the review authors' judgements about each domain presented as percentages across included studies. We will also provide a risk of bias and applicability concerns summary demonstrating review authors' judgements about each domain for each included study. We will use the Deek's test<sup>29</sup> to interrogate for publication bias. This test is specifically designed for detecting funnel plot asymmetry in reviews of diagnostic studies<sup>28</sup>.

Confidence in cumulative evidence

We will utilise the Grading of Recommendations Assessment, Development, and Evaluation (GRADE) approach to assess strength of the body of evidence<sup>30</sup>. The GRADE system classifies the quality of evidence into one of four grades:

- 1. *High*: further research is very unlikely to change our confidence in the estimate of effect;
- 2. *Moderate*: further research is likely to have an important impact on our confidence in the estimate of effect and may change the estimate;
- 3. *Low*: further research is very likely to have an important impact on our confidence in the estimate of effect and is likely to change the estimate;
- 4. *Very low*: any estimate of effect is very uncertain<sup>31</sup>.

A judgement is made on the individual studies used to provide the pooled effect estimates and the quality of evidence is then downgraded by the cumulative presence of i) bias (see risk of bias analysis), ii) inconsistency (i.e. heterogeneity present on I<sup>2</sup> statistic), iii) indirectness (i.e. high concerns regarding applicability, see table 2), iv) imprecision (small sample sizes, wide confidence intervals, inadequately powered studies), and v) publication bias (see Deek's test<sup>29</sup> comments)<sup>32</sup>. GRADE classifications will be independently conducted by two review authors (GM and KB) with any disagreements resolved by mutual discussion and, where necessary, adjunction by a third reviewer (RC/SD/CS).

We will rate the completeness of reporting for each study out of 30 using the STARD 2015 checklist<sup>21</sup>. A score of 0-10, 11-20, 21-30 will indicate a low, moderate, and high quality of completeness of reporting, respectively.

Patient and Public Involvement

Patients and the public were not involved in development of the research question and outcome measures, nor the study design. The study does not involve patient recruitment and patients were not involved in conduct of the study. We plan to liaise closely with patients, special interest groups, and charities in the dissemination of our results in printed and electronic media. Meta-data and information about the study will also be made available through our website ([www.muirmaxwellcentre.com](http://www.muirmaxwellcentre.com)).

Ethics and dissemination



Ethical approval is not required as primary data will not be collected. Results will be disseminated in peer-reviewed journals, conference presentations, and in press releases. Meta-data and information about the study will also be made available through our website ([www.muirmaxwellcentre.com](http://www.muirmaxwellcentre.com)) and via social media.

## References

1. Garratt E, Barnes H, Dibben C. Health administrative data: Exploring the potential for academic research. St Andrews: Administrative Data Liaison Service. 2010
2. Loke YK. Use of databases for clinical research. *Archives of Disease in Childhood* 2014;99(6):587-89. doi: 10.1136/archdischild-2013-304466
3. St Germaine-Smith C, Metcalfe A, Pringsheim T, et al. Recommendations for optimal ICD codes to study neurologic conditions: a systematic review. *Neurology* 2012;79(10):1049-55. doi: 10.1212/WNL.0b013e3182684707
4. Abraha I, Montedori A, Eusebi P, et al. The Current State of Validation of Administrative Healthcare Databases in Italy: A Systematic Review. *Pharmacoepidemiology and Drug Safety* 2012;21:400-00.
5. Traversa G, Bianchi C, Da Cas R, et al. Cohort study of hepatotoxicity associated with nimesulide and other non-steroidal anti-inflammatory drugs. *BMJ* 2003;327(7405):18-22. doi: 10.1136/bmj.327.7405.18
6. Hottes TS, Skowronski DM, Hiebert B, et al. Influenza vaccine effectiveness in the elderly based on administrative databases: change in immunization habit as a marker for bias. *PLoS One* 2011;6(7):e22618. doi: 10.1371/journal.pone.0022618
7. Lalmohamed A, Vestergaard P, Cooper C, et al. Timing of stroke in patients undergoing total hip replacement and matched controls: a nationwide cohort study. *Stroke* 2012;43(12):3225-9. doi: 10.1161/STROKEAHA.112.668509
8. Schneeweiss S, Avorn J. A review of uses of health care utilization databases for epidemiologic research on therapeutics. *J Clin Epidemiol* 2005;58(4):323-37. doi: 10.1016/j.jclinepi.2004.10.012
9. Krarup LH, Boysen G, Janjua H, et al. Validity of stroke diagnoses in a National Register of Patients. *Neuroepidemiology* 2007;28(3):150-4. doi: 10.1159/000102143
10. Kee VR, Gilchrist B, Granner MA, et al. A systematic review of validated methods for identifying seizures, convulsions, or epilepsy using administrative and claims data. *Pharmacoepidemiology and Drug Safety* 2012;21:183-93. doi: 10.1002/pds.2329
11. Jette N, Reid AY, Quan H, et al. How accurate is ICD coding for epilepsy? *Epilepsia* 2010;51(1):62-9. doi: 10.1111/j.1528-1167.2009.02201.x
12. Benson T. The history of the Read Codes: the inaugural James Read Memorial Lecture 2011. *Inform Prim Care* 2011;19(3):173-82.
13. Fonferko-Shadrach B, Lacey AS, White CP, et al. Validating epilepsy diagnoses in routinely collected data. *Seizure* 2017;52:195-98. doi: 10.1016/j.seizure.2017.10.008
14. Tu K, Wang M, Jaakkimainen RL, et al. Assessing the validity of using administrative data to identify patients with epilepsy. *Epilepsia* 2014;55(2):335-43. doi: 10.1111/epi.12506
15. Franchi C, Giussani G, Messina P, et al. Validation of healthcare administrative data for the diagnosis of epilepsy. *J Epidemiol Community Health* 2013;67(12):1019-24. doi: 10.1136/jech-2013-202528
16. Reid AY, St Germaine-Smith C, Liu M, et al. Development and validation of a case definition for epilepsy for use with administrative health data. *Epilepsy Res* 2012;102(3):173-9. doi: 10.1016/j.epilepsyres.2012.05.009
17. Parko K, Thurman DJ. Prevalence of epilepsy and seizures in the Navajo Nation 1998-2002. *Epilepsia* 2009;50(10):2180-5. doi: 10.1111/j.1528-1167.2009.02140.x
18. Pugh MJ, Van Cott AC, Cramer JA, et al. Trends in antiepileptic drug prescribing for older patients with new-onset epilepsy: 2000-2004. *Neurology* 2008;70(22 Pt 2):2171-8. doi: 10.1212/01.wnl.0000313157.15089.e6
19. Christensen J, Vestergaard M, Olsen J, et al. Validation of epilepsy diagnoses in the Danish National Hospital Register. *Epilepsy Res* 2007;75(2-3):162-70. doi: 10.1016/j.epilepsyres.2007.05.009

20. Tan M, Wilson I, Braganza V, et al. Development and validation of an epidemiologic case definition of epilepsy for use with routinely collected Australian health data. *Epilepsy Behav* 2015;51:65-72. doi: 10.1016/j.yebeh.2015.06.031

21. Bossuyt PM, Reitsma JB, Bruns DE, et al. STARD 2015: An Updated List of Essential Items for Reporting Diagnostic Accuracy Studies. *Clin Chem* 2015;61(12):1446-52. doi: 10.1373/clinchem.2015.246280

22. Moher D, Shamseer L, Clarke M, et al. Preferred reporting items for systematic review and meta-analysis protocols (PRISMA-P) 2015 statement. *Syst Rev* 2015;4:1. doi: 10.1186/2046-4053-4-1

23. Shamseer L, Moher D, Clarke M, et al. Preferred reporting items for systematic review and meta-analysis protocols (PRISMA-P) 2015: elaboration and explanation. *BMJ* 2015;349:g7647. doi: 10.1136/bmj.g7647

24. Moher D, Liberati A, Tetzlaff J, et al. Preferred reporting items for systematic reviews and meta-analyses: the PRISMA statement. *BMJ* 2009;339:b2535. doi: 10.1136/bmj.b2535

25. World Health Organisation. Classification of Diseases (ICD), 2016. [www.who.int/classifications/icd/en/](http://www.who.int/classifications/icd/en/).

26. Florkowski CM. Sensitivity, specificity, receiver-operating characteristic (ROC) curves and likelihood ratios: communicating the performance of diagnostic tests. *Clin Biochem Rev* 2008;29 Suppl 1:S83-7.

27. Whiting PF, Rutjes AW, Westwood ME, et al. QUADAS-2: a revised tool for the quality assessment of diagnostic accuracy studies. *Ann Intern Med* 2011;155(8):529-36. doi: 10.7326/0003-4819-155-8-201110180-00009

28. Leeflang MM, Deeks JJ, Takwoingi Y, et al. Cochrane diagnostic test accuracy reviews. *Syst Rev* 2013;2:82. doi: 10.1186/2046-4053-2-82

29. Deeks JJ, Macaskill P, Irwig L. The performance of tests of publication bias and other sample size effects in systematic reviews of diagnostic test accuracy was assessed. *J Clin Epidemiol* 2005;58(9):882-93. doi: 10.1016/j.jclinepi.2005.01.016

30. Schünemann H, Broek J, Oxman A. GRADE handbook for grading quality of evidence and strength of recommendation. 2009; 3.2. [www.ccims.net/gradepr](http://www.ccims.net/gradepr).

31. Mbizvo GK, Dixon P, Hutton JL, et al. Levetiracetam add-on for drug-resistant focal epilepsy: an updated Cochrane Review. *Cochrane Database Syst Rev* 2012;9:CD001901. doi: 10.1002/14651858.CD001901.pub2

32. Singh S, Chang SM, Matchar DB, et al. Chapter 7: grading a body of evidence on diagnostic tests. *J Gen Intern Med* 2012;27 Suppl 1:S47-55. doi: 10.1007/s11606-012-2021-9

Authors’ contributions

GKM and RFMC conceived the idea for the protocol and made the main contribution to planning and preparation of timelines for its completion. GKM and KB put together and tested various search strategies for the protocol and, after consultation with CRS, had these reviewed and approved by Marshall Dozier at the University of Edinburgh library, who’s support we acknowledge. GKM and RC planned the data extraction and statistical analysis, as well as of risk of bias, quality of evidence, and completeness of reporting assessments. GKM designed the tables and wrote the first draft of the manuscript, which was then reviewed and amended by KB, CRS, SD and RFMC. GKM, KB, CRS, SD and RFMC then approved the final written manuscript. RFMC is the guarantor for the work.

Funding statement

This work was supported by Epilepsy Research UK (R44007) and the Juliet Bergqvist Memorial Fund. The funders had no role in the design of the protocol, its preparation, analyses, interpretation of the data, manuscript preparation, or decision to submit.

Competing interests statement

The authors declare no competing interests.

Additional information



Any future amendments of the protocol will be listed in this section along with a date, description, and rational for each amendment.

For peer review only

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47

**PRISMA-P (Preferred Reporting Items for Systematic review and Meta-Analysis Protocols) 2015 checklist: recommended items to address in a systematic review protocol\***

Section and topic	Item No	Checklist item	Result	Page number where item can be found in manuscript Mbizvo <i>et al.</i> : <i>Manuscript ID bmjopen-2017-020824</i>
<b>ADMINISTRATIVE INFORMATION</b>				
Title:				
Identification	1a	Identify the report as a protocol of a systematic review	YES	Page 1 – Title
Update	1b	If the protocol is for an update of a previous systematic review, identify as such	NO	Not applicable (new review)
Registration	2	If registered, provide the name of the registry (such as PROSPERO) and registration number	YES	Page 2 - Abstract
Authors:				
Contact	3a	Provide name, institutional affiliation, e-mail address of all protocol authors; provide physical mailing address of corresponding author	YES	Page 1 – Title page
Contributions	3b	Describe contributions of protocol authors and identify the guarantor of the review	YES	Page 13 – Authors’ contributions
Amendments	4	If the protocol represents an amendment of a previously completed or published protocol, identify as such and list changes; otherwise, state plan for documenting important protocol amendments	YES	Page 13 – Additional information.
Support:				
Sources	5a	Indicate sources of financial or other support for the review	YES	Page 13 – Funding statement
Sponsor	5b	Provide name for the review funder and/or sponsor	YES	Page 13 – Funding statement
Role of sponsor or funder	5c	Describe roles of funder(s), sponsor(s), and/or institution(s), if any, in developing the protocol	YES	Page 13 – Funding statement
<b>INTRODUCTION</b>				
Rationale	6	Describe the rationale for the review in the context of what is already known	YES	Pages 2-3 – Introduction
Objectives	7	Provide an explicit statement of the question(s) the review will address with reference to participants, interventions, comparators, and outcomes (PICO)	YES	Pages 3-4 – Aims and objectives
<b>METHODS</b>				
Eligibility criteria	8	Specify the study characteristics (such as PICO, study design, setting, time frame) and report characteristics (such as years considered, language, publication status) to be used as criteria for eligibility for the review	YES	Pages 4-5 – Eligibility criteria
Information sources	9	Describe all intended information sources (such as electronic databases, contact with study authors, trial registers or other grey literature sources) with planned dates of coverage	YES	Pages 5-6 – Information sources
Search strategy	10	Present draft of search strategy to be used for at least one electronic database, including planned limits, such that it could be repeated	YES	Pages 5-6 – Information sources and table 1

Study records:				
Data management	11a	Describe the mechanism(s) that will be used to manage records and data throughout the review	YES	Pages 6-7 – Study records
Selection process	11b	State the process that will be used for selecting studies (such as two independent reviewers) through each phase of the review (that is, screening, eligibility and inclusion in meta-analysis)	YES	Page 7 – Selection process
Data collection process	11c	Describe planned method of extracting data from reports (such as piloting forms, done independently, in duplicate), any processes for obtaining and confirming data from investigators	YES	Page 7 – Data collection process and items
Data items	12	List and define all variables for which data will be sought (such as PICO items, funding sources), any pre-planned data assumptions and simplifications	YES	Page 7 – Data collection process and items, data collection tool
Outcomes and prioritization	13	List and define all outcomes for which data will be sought, including prioritization of main and additional outcomes, with rationale	YES	Page 7 – Data collection process and items, data collection tool
Risk of bias in individual studies	14	Describe anticipated methods for assessing risk of bias of individual studies, including whether this will be done at the outcome or study level, or both; state how this information will be used in data synthesis	YES	Page 10-11 – Risk of bias analysis
Data synthesis	15a	Describe criteria under which study data will be quantitatively synthesised	YES	Page 9 – Systematic review and meta-analysis outcomes
	15b	If data are appropriate for quantitative synthesis, describe planned summary measures, methods of handling data and methods of combining data from studies, including any planned exploration of consistency (such as $I^2$ , Kendall's $\tau$ )	YES	Page 9 – Systematic review and meta-analysis outcomes
	15c	Describe any proposed additional analyses (such as sensitivity or subgroup analyses, meta-regression)	YES	Page 9 – Systematic review and meta-analysis outcomes
	15d	If quantitative synthesis is not appropriate, describe the type of summary planned	YES	Page 9 – Systematic review and meta-analysis outcomes
Meta-bias(es)	16	Specify any planned assessment of meta-bias(es) (such as publication bias across studies, selective reporting within studies)	YES	Page 9-11 – Systematic review and meta-analysis outcomes, risk of bias analysis
Confidence in cumulative evidence	17	Describe how the strength of the body of evidence will be assessed (such as GRADE)	YES	Page 11 – Confidence in cumulative evidence

**\* It is strongly recommended that this checklist be read in conjunction with the PRISMA-P Explanation and Elaboration (cite when available) for important clarification on the items. Amendments to a review protocol should be tracked and dated. The copyright for PRISMA-P (including checklist) is held by the PRISMA-P Group and is distributed under a Creative Commons Attribution Licence 4.0.**

*From: Shamseer L, Moher D, Clarke M, Ghersi D, Liberati A, Petticrew M, Shekelle P, Stewart L, PRISMA-P Group. Preferred reporting items for systematic review and meta-analysis protocols (PRISMA-P) 2015: elaboration and explanation. BMJ. 2015 Jan 2;349(jan02 1):g7647.*