

**Supplementary Table 1** - Prevalence and risk of stillbirth associated with the 16 possible combinations of change in strength and frequency of movement in the last 2 weeks

Strength	Frequency	Stillbirths N=290 (missing=1)	Controls N=724 (missing=9)	Univariable OR (95%CI)
Increase	Increase	29 (10.0%)	232 (32.0%)	0.18 (0.11, 0.28)
Increase	Decrease	7 (2.4%)	27 (3.7%)	0.37 (0.16, 0.88)
Increase	Same	16 (5.5%)	190 (26.2%)	0.12 (0.07, 0.21)
Increase	Unknown	1 (0.3%)	6 (0.8%)	0.24 (0.03, 2.01)
Decrease	Increase	1 (0.3%)	2 (0.3%)	0.72 (0.06, 7.97)
Same	Increase	5 (1.7%)	13 (1.8%)	0.55 (0.19, 1.58)
Unknown	Increase	2 (0.7%)	6 (0.8%)	0.48 (0.10, 2.40)
Decrease	Decrease	51 (17.5%)	23 (3.2%)	3.17 (1.84, 5.46)
Same	Decrease	22 (7.6%)	10 (1.4%)	3.15 (1.44, 6.88)
Unknown	Decrease	5 (1.7%)	3 (0.4%)	2.38 (0.56, 10.16)
Decrease	Unknown	2 (0.7%)	2 (0.3%)	1.43 (0.20, 10.29)
Same	Unknown	5 (1.7%)	2 (0.3%)	3.57 (0.68, 18.73)
Unknown	Same	8 (2.7%)	6 (0.8%)	1.91 (0.65, 5.63)
Unknown	Unknown	7 (2.4%)	7 (1.0%)	1.43 (0.49, 4.18)
Decrease	Same	8 (2.7%)	23 (3.2%)	0.50 (0.22, 1.15)
Same	Same	121 (41.6%)	172 (23.5%)	Reference