

Supplementary Table: Demographic characteristics of residents admitted between 1 January 2008 and 26 February 2017 by study groups

	Internal department				Psychiatry department			
	Pre-incident	Post-incident			Pre-incident	Post-incident		
		Non-evacuees	Evacuees	New admittees		Non-evacuees	Evacuees	New admittees
Sex, number (%)								
Male	69 (35.6)	9 (21.4)	14 (30.4)	63 (35.8)	57 (67.1)	4 (36.4)	43 (68.3)	41 (64.1)
Female	125 (64.4)	33 (78.6)	32 (69.6)	113 (64.2)	28 (32.9)	7 (63.6)	20 (31.8)	23 (35.9)
Age at endpoint*, median (IQR)	85.0 (10.0)	85.0 (10.0)	87.0 (14.0)	86.0 (11.0)	57.0 (27.0)	47.0 (22.0)	63.0 (25.0)	63.5 (28.0)
Primary disease**								
Cardiovascular disease ¹	41 (21.1)	6 (14.3)	5 (10.9)	46 (26.1)	–	–	–	–
Lifestyle disease ²	19 (9.8)	0 (0.0)	6 (13.0)	16 (9.1)	–	–	–	–
Nervous disease ³	56 (28.9)	13 (31.0)	18 (39.1)	42 (23.9)	–	–	–	–
Mental illness	20 (10.3)	8 (19.1)	9 (19.6)	13 (7.4)	–	–	–	–
Other ⁴	58 (29.9)	15 (35.7)	8 (17.4)	59 (33.5)	–	–	–	–
ADL**, median (IQR)								
Bed morbidity	5 (3)	5 (3)	4 (2)	6 (1)	–	–	–	–
Toileting	6 (2)	6 (2)	4 (3)	6 (0)	–	–	–	–
Transfers	6 (3)	6 (3)	4 (3)	6 (0)	–	–	–	–
Eating	5 (4)	5 (4)	2 (4)	5 (1)	–	–	–	–
Total	23 (12)	23 (13)	12 (12)	23 (1)	–	–	–	–
Medical condition**, number (%)								
1	8 (4.1)	0 (0.0)	3 (6.5)	6 (3.4)	–	–	–	–
2	125 (64.4)	28 (66.7)	38 (82.6)	120 (68.2)	–	–	–	–
3	51 (26.3)	12 (28.6)	4 (8.7)	49 (27.8)	–	–	–	–
Missing	10 (5.2)	2 (4.8)	1 (2.2)	1 (0.6)	–	–	–	–

Multiple admission records were counted as different cases in this table. *endpoint with death/discharge or at the end of the study period (25 June 2017); **evaluated at admission; ADL: activities of daily living (on a scale from 0 to 24, with higher scores reflecting greater ability); IQR: interquartile range; SD: standard deviation; 1: stroke, coronary heart disease, etc.; 2: hypertension, diabetes, kidney disease, etc.; 3: Parkinson's disease, dementia, etc.; 4: digestive and respiratory diseases, cancer, etc.