BMJ Open Short implants (≤ 6 mm) versus longer implants with sinus floor elevation in atrophic posterior maxilla: a systematic review and meta-analysis

Qi Yan,^{1,2} Xinyu Wu,^{1,2} Meiying Su,² Fang Hua ,^{3,4} Bin Shi²

To cite: Yan Q, Wu X, Su M, *et al.* Short implants (≤ 6 mm) versus longer implants with sinus floor elevation in atrophic posterior maxilla: a systematic review and meta-analysis. *BMJ Open* 2019;**9**:e029826. doi:10.1136/bmjopen-2019-029826

► Prepublication history and additional material for this paper are available online. To view these files, please visit the journal online (<http://dx.doi.org/10.1136/bmjopen-2019-029826>).

QY and XW contributed equally.

Received 13 February 2019
Revised 06 September 2019
Accepted 26 September 2019

© Author(s) (or their employer(s)) 2019. Re-use permitted under CC BY-NC. No commercial re-use. See rights and permissions. Published by BMJ.

For numbered affiliations see end of article.

Correspondence to

Dr Fang Hua;
huafang@whu.edu.cn

Professor Bin Shi;
shibin_dentist@whu.edu.cn

ABSTRACT

Objectives To compare the use of short implants (≤ 6 mm) in atrophic posterior maxilla versus longer implants (≥ 10 mm) with sinus floor elevation.

Design A systematic review and meta-analysis based on randomised controlled trials (RCTs).

Data sources Electronic searches were conducted in PubMed, Embase and the Cochrane CENTRAL.

Retrospective and prospective hand searches were also performed.

Eligibility criteria RCTs comparing short implants (≤ 6 mm) and longer implants (≥ 10 mm) with sinus floor elevation were included. Outcome measures included implant survival (primary outcome), marginal bone loss (MBL), complications and patient satisfaction.

Data extraction and synthesis Risks of bias in and across studies were evaluated. Meta-analysis, subgroup analysis and sensitivity analysis were undertaken. Quality of evidence was assessed according to Grading of Recommendations Assessment, Development and Evaluation.

Results A total of seven RCTs involving 310 participants were included. No significant difference in survival rate was found for 1–3 years follow-up (RR 1.01, 95% CI 0.97 to 1.04, $p=0.74$, $I^2=0\%$, moderate-quality evidence) or for 3 years or longer follow-up (RR 1.00, 95% CI 0.97 to 1.04, $p=0.79$, $I^2=0\%$, moderate-quality evidence). However, short implants (≤ 6 mm) showed significantly less MBL in 1–3 years follow-up (MD = -0.13 mm, 95% CI -0.21 to 0.05 ; $p=0.001$, $I^2=87\%$, low-quality evidence) and in 3 years or longer follow-up (MD = -0.25 mm, 95% CI -0.40 to 0.10 ; $p=0.001$, $I^2=0\%$, moderate-quality evidence). In addition, short implant (≤ 6 mm) resulted in fewer postsurgery reaction (RR 0.11, 95% CI 0.14 to 0.31, $p<0.001$, $I^2=40\%$, moderate-quality evidence) and sinus perforation or infection (RR 0.11, 95% CI 0.02 to 0.63, $p=0.01$, $I^2=0\%$, moderate-quality evidence).

Conclusions For atrophic posterior maxilla, short implants (≤ 6 mm) are a promising alternative to sinus floor elevation, with comparable survival rate, less MBL and postsurgery reactions. Additional high-quality studies are needed to evaluate the long-term effectiveness of short implants (≤ 6 mm).

Trial registration number The protocol has been registered at PROSPERO (CRD42018103531).

Strengths and limitations of this study

- Only randomised controlled clinical trials were included.
- Participant-unit data were used for syntheses.
- Subgroup analyses by follow-up length and categories of complications were performed.
- Serious risks of bias were found within and across studies and the quality of evidence was only low to moderate.

INTRODUCTION

Dental implants supporting prosthesis are commonly considered a promising method for the rehabilitation of missing teeth.^{1–3} However, dental implantation in the posterior maxilla is usually challenging due to insufficient vertical bone volume, poor bone quality, limited visibility, reduced interarch space and sinus pneumatisation.^{4–5} These conditions are exacerbated if patients have a history of wearing removable dentures.⁶

To achieve sufficient vertical bone volume in the posterior maxilla, sinus floor elevation using the lateral window approach or the osteotomy technique has been introduced and widely used over the past 40 years.^{7,8} The lateral window approach is commonly used in dental implantation procedures.⁹ Using these techniques with or without bone grafting, conventional implants can be placed in the elevated sites. The implant success rate is typically greater than 90% in long-term evaluation.^{10–12} However, sinus floor elevation surgery is usually associated with higher cost, more complicated surgical procedures and a high prevalence of complications such as infection, sinus membrane perforation and graft failure.^{13–15} In addition, the clinical outcome of sinus floor elevation can also be restricted by extremely insufficient residual bone height, abnormal sinus anatomy,

thickening of the sinus membrane, stability of the grafted bone and the number of missing teeth.^{13 16–18}

Short implants with improved implant design and surface properties have been successfully applied as an alternative to sinus floor elevation surgery and have shown good results in posterior maxilla. Implants ≤ 10 mm,¹⁹ ≤ 8 mm,²⁰ ≤ 7 mm²¹ and 6–8 mm²² are reported to have survival rates comparable to those of longer implants. In addition, short implants ≤ 6 mm in length have been introduced as another alternative in atrophic posterior maxilla.^{6 23 24} Short implants require a less complicated surgical approach and could be used in cases when sinus floor elevation surgery is not applicable,^{25 26} especially in cases of maxillary sinusitis, maxillary cyst, large vessels and other cases involving abnormal sinus anatomy. Studies have explored the short-term and long-term survival rates of short implants (≤ 6 mm).^{26–30} Unfortunately, the evidence supporting the use of short implants (≤ 6 mm) in the posterior maxilla is weak, and no guideline statement is currently recommended.

The present systematic review aims to compare the effectiveness of short implants (≤ 6 mm) and longer implants (≥ 10 mm) with sinus floor elevation in atrophic posterior maxilla. Our null hypothesis was that the survival rate, patient satisfaction, marginal bone loss (MBL) and surgery-related complications of short implants (≤ 6 mm) were comparable to longer implants in combination with sinus floor elevation.

MATERIALS AND METHODS

Protocol and registration

This systematic review and meta-analysis was conducted in accordance with the Preferred Reporting Items for Systematic Reviews and Meta-Analyses guidelines.³¹

Eligible criteria

Randomised controlled trials (RCTs) meeting the following predetermined inclusion criteria (PICOS format) were included:

- ▶ Population: Partially edentulous patients in the premolar and molar regions of the maxilla, for whom the residual bone height in the atrophic posterior maxilla was sufficient for the insertion of a short implant (≤ 6 mm) but insufficient for the insertion of longer implants.
- ▶ Intervention: One or more short implants (≤ 6 mm) were placed in the posterior maxilla without sinus floor elevation in the short implant group.
- ▶ Comparison: One or more longer implants were placed in the posterior maxilla after sinus floor elevation by any technique in the elevation group.
- ▶ Outcomes: The primary (survival rate) and secondary (MBL, complications and patient satisfaction) outcomes of interest were measured, with a follow-up length of 1 year or longer postloading.

Information sources and search strategies

Two content experts (QY and XW) searched PubMed, Embase and the Cochrane CENTRAL (The Cochrane

Central Registration of Controlled Trials) for RCTs, independently and in duplicate. The last search was conducted on 31 May 2018. A methodologist (FH) was consulted to resolve any disagreements. Main search terms included: “dental implant”, “short implant”, “ultrashort”, “alveolar bone loss”, “atrophic maxilla”, “sinus lift”, and “sinus floor elevation”. No restriction was set regarding publication year, publication language or status. The detailed search strategies are listed in the online supplementary file. In addition, retrospective and prospective searches were conducted by checking the reference lists of key articles and studies citing these key articles, using Google Scholar.

Study selection and data collection

Two review authors (QY and XW) conducted the study selection independently and in duplicate. The titles and abstracts of all records were scanned. Full texts of studies were obtained in cases they appeared to meet the inclusion criteria or further information were needed to determine eligibility. Studies excluded at this or subsequent stages were recorded with the reasons for exclusion. All disagreements were resolved by discussion.

Two review authors (QY and XW) extracted the data independently and in duplicate using specifically designed data extraction forms. The extracted data included citation details (year of publication, country of origin, setting and source of funding), details on the participants (demographic characteristics, residual bone height and inclusion criteria), details of intervention (implant length, diameter, brand, surface structure, surgical method, follow-up time, prosthesis type), outcome assessment, sample size calculation and trial registration. Corresponding authors were contacted for missing data or information.

Risk of bias of included studies

Two authors (QY and XW) assessed the risk of bias of each included study independently and in duplicate using the Cochrane risk of bias assessment tool for RCTs.³² Disagreements were resolved through discussion. A third review author (FH) was consulted when necessary. Seven domains were assessed, including sequence generation (selection bias), allocation concealment (selection bias), blinding of participants and personnel (performance bias), blinding of outcome assessment (detection bias), incomplete outcome data (attrition bias), selective outcome reporting (reporting bias) and other bias (factors that had potential influence on outcomes but were not evenly distributed across groups or not clearly reported, such as the manufacturer or diameter of implants). Individual studies were categorised as having low, high or unclear risk of bias. The risk of bias across studies was determined according to the risk of bias in each included study.

Assessment of heterogeneity

Clinical heterogeneity among the included studies was assessed by comparing study design, participant

conditions (gender, age, residual bone height), intervention (implant length, diameter, surface structure, surgical method) and outcome measures. Statistical heterogeneity was evaluated using Cochrane's Q test and the I^2 statistic. In the Q test, a $p < 0.1$ was considered an indication of significant heterogeneity.

Assessment of publication bias

If at least 10 studies were included in a meta-analysis, We would have used a funnel plot and the Egger's test³³ asymmetry to assess the potential existence of publication bias if at least ten studies were included in a meta-analysis.

Synthesis of results

The unit of analysis was set as participant rather than implant.³⁴ RevMan V.5.3 software was used for data synthesis. Meta-analyses were undertaken only when at least two studies that made similar comparisons reported the same outcomes. The effect measures were risk ratio (RR) for dichotomous outcomes (implant survival and complications) and mean difference (MD) for continuous outcomes (MBL). RR was calculated through Mantel-Haenszel analysis and MD was calculated through inverse variance. $P < 0.05$ was considered statistically significant. The fixed-effect model was used when fewer than four studies were included in a meta-analysis, and the random-effects model was used when four or more studies were included.^{35–38}

Additional analysis

Subgroup analysis by length of follow-up was performed to control for the possibility that function time might influence implant survival.³⁹ In addition, subgroup analysis by categories of complications was performed. Complications were categorised according to into postsurgery reaction (bleeding, swelling and discomfort), biological complications (sinus perforation or infection, implant mobile, peri-implant mucositis and peri-implantitis) and technical complications (complications related to screws and crowns). If risks of bias in some studies were serious, we performed sensitivity analysis by excluding these studies. Considering only three studies were included in the MBL 3years or longer follow-up but the meta-analysis for MBL included more than three studies, a sensitivity analysis was conducted for MBL by using fixed-effect model.

Summary of findings

Grading of Recommendations Assessment, Development and Evaluation approach⁴⁰ was adopted to evaluate quality of evidence in this systematic review. A summary of findings table was made with an online tool (cebgrade.mcmaster.ca/gradepr.html). Outcomes were evaluated including survival rate and MBL of one to table was made with an online tool (cebgrade.mcmaster.ca/gradepr.html). Outcomes were evaluated including survival rate and MBL of 1–3years and 3years or longer follow-up, and complications. Five domains in quality of evidence were assessed: the overall risk of bias, directness

Figure 1 Flow diagram for study selection. RCT, randomised controlled trial.

of evidence, consistency of results, precision of estimates, as well as the risk of publication bias. The quality of the body of evidence was classified into four categories: high, moderate, low and very low.

Patient and public involvement

No patient or public was involved in this systematic review.

RESULTS

Study selection

Electronic searches identified a total of 879 titles and abstracts in PubMed and 251 in Embase. After removal of duplicates, the titles and abstracts of 1013 unique items were screened. We then retrieved the full texts of 25 potentially eligible articles, of which 20 were excluded for the reasons described in figure 1. Retrospective and prospective hand searches yielded two more articles. Finally, seven studies^{29 41–46} met our eligibility criteria and were included in this review (figure 1).

Study characteristics

The characteristics of the seven included studies are listed in table 1. One study was a split-mouth trial, and the rest were two-arm parallel RCTs. The length of follow-up ranged from 1 to 3years. For sinus floor elevation, either osteotomy-mediated sinus floor elevation or the lateral window technique was adopted. In two studies, single crowns were used as the rehabilitation method; in the

Table 1 Characteristics of the included studies

Study	Subjects		Short implant group					Elevation group				Rehabilitation methods	Outcome measures	
	Design	Gender (male/female)	Age	RBH (mm)	Follow-up period PL	Implant (n)	LEN (mm)	DIA (mm)	Implant (n)	LEN (mm)	DIA (mm)			Elevation methods
Bolle <i>et al</i> , 2018 ²⁹	RCT (TAP)	INT 7/13 CON 12/8	59.35 (47–73) 63.25 (46–72)	4–5	1 year	37	4	4 or 4.5	41	10, 11.5, 13	4 or 4.5	Osteotomy approach	SG or SP	NIF, MBL, COM
Gastaldi <i>et al</i> , 2018 ⁴¹	RCT (TAP)	INT 3/17 CON 7/13	58.6 (39–80) 52.8 (42–70)	4–6	3 years	20	5	5	20	10, 11.5, 13, 15	5	Lateral window technique	SG or SP	NIF, MBL, COM
Gastaldi <i>et al</i> , 2017 ⁴²	RCT (TAP)	INT 3/7 CON 5/5	53.4 (43–67) 58.6 (48–70)	5–7	3 years	16	5 or 6	5	18	10	5	Osteotomy approach	SG or SP	NIF, MBL, COM, PS
Guljé <i>et al</i> , 2014 ⁴³	RCT (TAP)	INT 7/14 CON 13/7	50 (30–71) 48 (29–72)	6–8	1 year	21	6	4	20	11	4	Lateral window technique	SG	NIF, MBL, COM, PS
Pohl <i>et al</i> , 2017 ⁴⁴	RCT (TAP)	49/52*	50.5 (20–75)*	5–7	3 years	67	6	4	70	11, 13, 15	4	Lateral window technique	SG	NIF, MBL, COM
Felice <i>et al</i> , 2018 ⁴⁵	RCT (SM)	11/9	57.6 (45–80)	5–7	3 years	39	6	4	44	10, 11.5, 13, 15	4	Lateral window technique	SG or SP	NIF, MBL, COM
Bechara <i>et al</i> , 2017 ⁴⁶	RCT (TAP)	INT 10/23 CON 9/11	47.5±16.2 49.2±13.4	≥4	3 years	45	6	4–8	45	10, 11.5, 13, 15	4–8	Lateral window technique	SG or SP	NIF, MBL, COM, PS

*Details for subject information in intervention and control group were not reported.

COM, complications; CON, control; DIA, implant diameter; INT, intervention; LEN, implant length; MBL, marginal bone loss; NIF, number of implant failures; NR, not reported; PL, postloading; PS, patient satisfaction; RBH, residual bone width under sinus floor; RCT, randomised controlled trial; SG, single crowns; SM, split mouth; SP, splinted prosthetics; TAP, two arm parallel.

Figure 2 Risk of bias in each included study.

remaining studies, single crowns or splinted prosthetics were used. The outcome measures used in these studies included implant failure, MBL, complications and patient satisfaction. Overall, 171 participants were included in the short implant groups, and 159 participants were included in the elevation groups.

Risk of bias assessment

The results of the risks of bias assessment are shown in figures 2 and 3. Selection bias and performance bias were assessed as low in all but one study by Bachara *et al*⁴⁶ due to inadequate description on random sequence generation and blinding of participants. For detection bias, most studies showed high risks because assessors could recognise sites that underwent sinus floor elevation. For attrition bias, three studies^{42 44 45} was assessed as high. Two studies^{44 46} showed high risk of reporting bias. Other risks of bias were considered high or unclear in three studies. Overall, all included studies were at high risk of bias for at least one domain (table 2).

Synthesis of results

Survival rate

Figure 4 shows the results of a meta-analysis for participant unit implant survival rate with a subgroup analysis based on length of follow-up. Five studies reported 100% survival of short implants (≤ 6 mm) within the study period. For this outcome, there was no evidence of a difference between the short implant group and the elevation group either 1–3 years postloading (RR 1.01, 95% CI 0.97 to 1.04, $p=0.74$, $I^2=0\%$, seven RCTs, 321 participants) or 3 years or longer postloading (RR 1.00, 95% CI 0.97 to 1.04, $p=0.79$,

Figure 3 Risk of bias across included studies.

$I^2=0\%$, five RCTs, 237 participants). Further details of the implant failures are summarised in table 3.

Marginal bone loss

The results of the meta-analysis and subgroup analysis regarding peri-implant MBL are shown in figure 5. A significant difference favouring the short implant group was found for both 1–3 years postloading (MD=−0.13, 95% CI −0.21 to 0.05; $p=0.001$, $I^2=87\%$, six RCTs, 249 participants) and 3 years or longer postloading (MD=−0.25, 95% CI −0.40 to 0.10; $p=0.001$, $I^2=0\%$, three RCTs, 88 participants). In sensitivity analysis by using fixed-effect model, results remained significant for both 1–3 years postloading (MD=−0.11, 95% CI −0.13 to 0.08; $p<0.001$, $I^2=87\%$) and 3 years or longer postloading (MD=−0.25, 95% CI −0.40 to 0.10; $p=0.001$, $I^2=0\%$).

Complications

Complications were categorised into postsurgery reaction, biological complications and technical complications (table 4). Short implant group was found with significantly less postsurgery reaction (RR 0.11, 95% CI 0.14 to 0.31, $p<0.001$, $I^2=40\%$, three RCTs, 184 participants) and sinus perforation or infection (RR 0.11, 95% CI 0.02 to 0.63, $p=0.01$, $I^2=0\%$). Only one study²⁹ reported implant migrating into sinus in 5% patients (1/19) in short implant group while in 10.5% patients (2/19) in sinus floor elevation group. No statistically significant difference was found in other complications between short implants (≤ 6 mm) and longer implants.

Patient satisfaction

Three studies^{42 43 46} reported patient satisfaction. Meta-analysis was not conducted because methods of evaluating patient satisfaction were different. Gullé *et al*⁴³ used a questionnaire to evaluate patient satisfaction before surgery and 1 year postloading. Both groups showed improvement of satisfaction after crown placement. Castaldi *et al*⁴² evaluated patient satisfaction in function and aesthetic aspects. All patients in the short implant group (10) were satisfied with both function and aesthetic aspects. However, three patients in elevation group (3/10) were partially satisfied with function. Bechara *et al*⁴⁶ used a questionnaire evaluating patient satisfaction in function, aesthetic, cleaning of the implant-supported restorations, satisfaction and cost. Significantly more patients in short implant group expressed satisfaction in cost. In the other four aspects, no significant difference was found between short implant group and sinus floor elevation group.

Quality of evidence

For survival rate, the quality of evidence in both subgroups was downgraded by one level (moderate quality evidence) due to serious risks of bias. For short-term MBL (1–3 years follow-up), quality of evidence was downgraded by two levels for serious risks of bias and inconsistency. For long-term MBL (3 years or longer follow-up), quality of evidence was downgraded by one level for serious risks

Table 2 Details on the risk of bias for each included study

Study	Random sequence generation	Allocation concealment	Blinding of patients/carers	Blinding of outcome assessment	Incomplete outcome data	Selective reporting	Other
Bolle <i>et al</i> , 2018 ²⁹	Low risk—quote: ‘a computer-generated restricted randomisation list’	Low risk—quote: ‘the information on how to treat each patient was enclosed in sequentially numbered, identical, opaque, sealed envelopes.’	Low risk—quote: ‘treatment allocation was concealed to the investigators in charge of enrolling and treating the patients.’	High risk—quote: ‘complications were dealt with directly and reported by the responsible clinicians, who were not blinded’; ‘augmented sites could be easily identified on radiographs due to the different implant lengths.’	Low risk—quote: ‘one patient from the short implant group and one from elevation group dropped out.’	Low risk—comment; All outcome measure in methods were reported in results	Unclear risk—comment: diameter of implants (4 mm or 4.5 mm) was not controlled
Gastaldi <i>et al</i> , 2018 ⁴¹	Low risk—quote: ‘a computer-generated restricted randomisation list’	Low risk—quote: ‘The randomised codes were enclosed in sequentially numbered, identical, opaque, sealed envelopes’	Low risk—quote: ‘treatment allocation was concealed to the investigators in charge of enrolling and treating the patients.’	High risk—quote: ‘augmented sites could be easily identified because of the different anatomy of the two sides after the augmentation procedure’	Low risk—comment: one patient dropped out of the short implant group (1/20), and two patients dropped out of the elevation group (2/20)	Low risk—comment; All outcome measures in methods were reported in results	Unclear risk—comment: information of short implants was not reported
Gastaldi <i>et al</i> , 2017 ⁴²	Low risk—quote: ‘a computer-generated restricted randomisation list’	Low risk—quote: ‘The randomised codes were enclosed in sequentially numbered, identical, opaque, sealed envelopes.’	Low risk—quote: ‘treatment allocation was concealed to the investigators in charge of enrolling and treating the patients’	High risk—quote: ‘sinus-lifted sites could be identified on radiographs because they appeared more radio-opaque and implants were longer.’	High risk—comment: no patients dropped out of the short implant group (0/10); two patients dropped out of the elevation group (2/10)	Low risk—comment; All outcome measures in methods were reported in results	Low risk
Guljé <i>et al</i> , 2014 ⁴³	Low risk—quote: ‘Randomisation was performed using a block randomization sequence to provide equal distribution of subjects.’	Low risk—quote: ‘A sealed envelope’	Low risk—quote: ‘A sealed envelope was opened by the surgical assistant at the beginning of the surgical procedure.’	High risk—quote: ‘blinding was possible in the clinical evaluation but not during analysis of the radiographs.’	Low risk—comment: no patient dropped out of the short implant group (0/21); one patient in the elevation group died (1/20)	Low risk—comment; All outcome measures in methods were reported in results	Low risk
Pohl <i>et al</i> , 2017 ⁴⁴	Low risk—quote: ‘A block randomization sequence was used to provide an equal distribution’	Low risk—quote: ‘A sealed envelope’	Low risk—quote: ‘After flap elevation, a sealed randomisation envelope was opened to allocate the subject to either one of the two treatment groups.’	Unclear risk—quote: ‘an independent examiner performed all the radiographic measurements.’ Other information was not reported.	High risk—comment: The reasons for incomplete reporting of MBL were not provided.	High risk—comment: MBL at 3-year follow-up was reported at the implant level rather than at the participant level	Low risk

Continued

Table 2 Continued

Study	Random sequence generation	Allocation concealment	Blinding of patients/carers	Blinding of outcome assessment	Incomplete outcome data	Selective reporting	Other
Felice <i>et al</i> , 2018 ⁴⁵	Low risk—quote: 'a computer-generated restricted randomisation list'	Low risk—quote: 'The information on how to treat site number one was enclosed in sequentially numbered, identical, opaque, sealed envelopes.'	Low risk—quote: 'Treatment allocation was concealed to the investigators in charge of enrolling and treating the patients.'	High risk—quote: 'augmented sites could be easily identified because of the different anatomy'	High risk—comment: it was a split-mouth design study, and two drop-outs (2/20) occurred	Low risk—comment: All outcome measures in methods were reported in results	Low risk
Bechara <i>et al</i> , 2017 ⁴⁶	Unclear risk—quote: 'Patients were randomly assigned'	Low risk—quote: 'a sequentially numbered sealed envelope'	Unclear risk—comment: not mentioned	Unclear risk—quote: 'At each annual inspection, an experienced, calibrated, independent examiner performed a careful clinical examination', but elevation site can be distinguished	Low risk—comment: one patient dropped out of the short implant group (1/33), and one patient dropped out of the elevation group (1/20)	High risk—comment: MBL was reported at the implant level rather than at the participant level	High risk—comment: diameter of implants was not controlled (4–8 mm)

MBL, marginal bone loss.

of bias. For complications, the quality of evidence in postsurgery reaction was moderate, downgrading by one level for serious risks of bias. The quality of evidence in other complications was low, downgrading by two levels for serious risks of bias and imprecision. Details are listed in table 5.

DISCUSSION

To the best of our knowledge, this is the first systematic review and meta-analysis to compare the clinical outcome of the use of short implants (≤ 6 mm) in atrophic posterior maxilla versus longer implants with sinus floor elevation. At 1 year or longer postloading, there

Figure 4 Forest plot for implant survival rate. M-H, Mantel-Haenszel; SI, short implant group; SFE, sinus floor elevation group.

Table 3 Details of implant failures reported in the included studies

Study	Short implant group				Elevation group			
	LEN (mm)	DIA (mm)	PAR/IMP (n)	Details	LEN (mm)	DIA (mm)	PAR/IMP (n)	Details
Bolle <i>et al</i> , 2018 ²⁹	4	4 or 4.5	2/3	PAR1. One implant was mobile 3 months after placement, and another implant migrated into the sinus 4 months after placement. PAR2. One implant was medially tilted 2 weeks after placement	10,11.5,13	4 or 4.5	4/6	PAR1. One implant was mobile 2 months after placement because of a perforation of the sinus lining at its detachment. Another implant was mobile 2 months later. PAR2. One implant migrated into the sinus 3 months after placement. PAR3. Two implants were mobile 3 months after placement because the patient insisted on wearing her removable denture. PAR4. One implant was mobile, and the patient experienced discomfort when chewing 5 months postloading.
Gastaldi <i>et al</i> , 2018 ⁴¹	5	5	1/1	PAR1. One implant failed 3 months postloading.	10,11.5,13,15	5	0	None
Felice <i>et al</i> , 2018 ⁴⁵	6	4	0	None	10,11.5,13,15	4	1/2	PAR1. Two implants failed due to peri-implantitis 2 years postloading.
Bechara <i>et al</i> , 2017 ⁴⁶	6	4–8	0	None	10,11.5,13,15	4–8	1/2	PAR1. Two implants were lost caused due to chronic sinus infection with loss of integration/implant stability 2 months after surgery.

DIA, implant diameter; LEN, implant length; PAR, participant; PAR/IMP, participant/implant.

is no significant difference in participant unit implant survival rate between the short implant group and the elevation group. The short implant group showed less

MBL than the elevation group for 1–3 years follow-up (low-quality evidence) and 3 years or longer follow-up (moderate-quality evidence). In addition, the short

Figure 5 Forest plot for marginal bone loss. SI, short implant group; SFE, sinus floor elevation group.

Table 4 Comparisons of complications

Outcome or subgroup titles	No of studies	No of participants	Statistical methods	Effect size
Postsurgery reaction	3	184	Risk ratio (Fixed, M-H, 95% CI)	0.11 (0.14 to 0.31)*
Biological complications				
Sinus perforation or infection	3	125	Risk ratio (Fixed, M-H, 95% CI)	0.11 (0.02 to 0.63)*
Implant mobile	2	132	Risk ratio (Fixed, M-H, 95% CI)	0.34 (0.06 to 2.06)
Peri-implant mucositis and peri-implantitis	2	54	Risk ratio (Fixed, M-H, 95% CI)	0.91 (0.14 to 5.79)
Technical complications			Risk ratio (Fixed, M-H, 95% CI)	
Screw loosening	3	169	Risk ratio (Fixed, M-H, 95% CI)	2.66 (0.93 to 7.60)
Crown loosening, decementation or chipping	5	223	Risk ratio (Random, M-H, 95% CI)	1.22 (0.33 to 4.49)

*Difference between the two groups was significant.
M-H, Mantel-Haenszel.

implant group showed fewer postsurgery reaction and sinus membrane perforation and infection.

The survival rate in this review was evaluated by participant unit as in a previous Cochrane review.³⁴ In this review, the overall survival rates for the short implant group and the elevation group were 98.21% and 96.08%, respectively, at 1–3 years follow-up and 99.20% and 98.23%, respectively, at longer than 3 years follow-up; no significant difference in survival rate was found. Other studies that assessed survival rate in implant unit had similar outcomes. A retrospective study⁴⁷ with a follow-up period of 17–48 months reported a 95.12% implant unit survival rate for 5–6 mm short implants. A prospective study⁶ of 2–3 years reported that 6 mm short implants with microrough surfaces achieved a 100% survival rate in posterior maxilla. Another retrospective study of 5–10 years⁴⁸ reported a 97% implant unit survival rate for 6 mm short implants supporting single crowns. All these results showed that short implants (≤ 6 mm) represent a promising rehabilitation method with respect to their short-term and long-term survival rates.

In this review, all of the failed short implants were 4 mm or 5 mm. Although the use of short implants (≤ 6 mm) could avoid complicated surgical procedures and related early failures, reduced implant length was still the major risk factor in survival rate. The authors of the included studies used wider implants (4–8 mm) to compensate for the short length of the implants. Finite element analyses showed that wider implants had increased functional surface area in cortical bone and decreased stress distribution on the implant neck; these qualities helped improve primary stability, produce a higher survival rate and reduce MBL.^{49–52} However, it was not determined whether implant length or diameter contributed more to implant failure. Another factor was implant surface structure. Studies^{53–56} have suggested that the implant surface influences bone-to-implant osseointegration, implant primary stability and MBL. In this review, implants 4 or

5 mm in length had novel surface structures, but they still presented a lower survival rate.

Significantly less MBL was found in the short implant group, and the difference was greater at the longer follow-up period. Additionally, in this review, 5 mm diameter implants tended to induce less MBL than 4 mm diameter implants. Implants ≤ 10 mm¹⁹ and ≤ 8 mm²⁰ were reported to induce MBL similar to that of longer implants, while implants ≤ 7 mm⁵⁷ showed less MBL. These results contradict a previous theory that short implants are more likely to have an extreme crown-to-implant ratio (C/I)⁵⁸ that induces more peri-implant bone loss and early implant failure.^{59–60} According to finite element analyses, inappropriate C/I results in adverse occlusal forces such as non-axial forces and overloading.⁶¹ Increased C/I was also correlated with more prosthesis complications such as screw loosening, implant or abutment fracture, chipping of the ceramic material and prosthesis fracture.^{62–65} However, the implants in the studies included in this systematic review had wider diameters (4–8 mm) and different surface structures. These two factors partially compensated for the complications of C/I and contributed to less MBL. Differences in implant diameter and surface structure also introduced heterogeneity among studies with respect to MBL. Short implants tolerated less MBL because of the limited implant length. As a result, less MBL was not necessarily correlated with better clinical outcome. MBL around short implants is still a challenging issue, and much effort should be made to resolve it.

With respect to complications, the use of short implants (≤ 6 mm) could decrease the incidence of postsurgery reactions and sinus membrane perforation and infection. Sinus membrane perforation was common in the elevation group.¹⁴ This was in accordance with a previous study⁶⁶ that reported more complications in cases involving longer implants with sinus floor elevation and that the surgical procedure made a major contribution to such

Table 5 Summary of findings**Short implant (≤ 6 mm) compared with longer implant (≥ 10 mm) with sinus floor elevation in atrophic posterior maxilla****Patient or population:** atrophic posterior maxilla**Intervention:** short implant (≤ 6 mm)**Comparison:** longer implant (≥ 10 mm) with sinus floor elevation

Outcomes	Anticipated absolute effects* (95% CI)		Relative effect (95% CI)	No of participants (studies)	Certainty of the evidence (GRADE)	Comments
	Assumed risk† (elevation group)	Corresponding risk (short implant group)				
Survival rate follow-up: range 1–3 years	961 per 1000	970 per 1000 (932 to 999)	RR 1.01 (0.97 to 1.04)	321 (7 RCTs)	⊕⊕⊕○ MODERATE‡	
Survival rate follow-up: range 3 years to longer years	982 per 1000	982 per 1000 (953 to 1000)	RR 1.00 (0.97 to 1.04)	237 (5 RCTs)	⊕⊕⊕○ MODERATE‡	
Marginal bone loss follow-up: range 1–3 years	The mean marginal bone loss ranged from 0.1 to 1.15 mm	The mean marginal bone loss in the intervention group was 0.13 mm lower (0.21 lower to 0.05 lower)	–	249 (6 RCTs)	⊕⊕○○ LOW§	
Marginal bone loss follow-up: range 3 years to longer years	The mean marginal bone loss ranged from 1.08 to 1.5 mm	The mean marginal bone loss in the intervention group was 0.25 mm lower (0.4 lower to 0.1 lower)	–	88 (3 RCTs)	⊕⊕⊕○ MODERATE‡	
Postsurgery reaction	307 per 1000	34 per 1000 (12 to 59)	RR 0.11 (0.04 to 0.31)	184 (3 RCTs)	⊕⊕⊕○ MODERATE‡	
Biological complications: sinus perforation or infection	197 per 1000	20 per 1000 (4 to 113)	RR 0.11 (0.02 to 0.63)	125 (3 RCTs)	⊕⊕○○ LOW¶	
Biological complications: implant mobile	59 per 1000	20 per 1000 (4 to 121)	RR 0.34 (0.06 to 2.06)	132 (2 RCTs)	⊕⊕○○ LOW¶	
Biological complications: peri-implant mucositis or peri-implantitis	200 per 1000	100 per 1000 (10 to 934)	RR 0.91 (0.14 to 5.79)	54 (2 RCTs)	⊕⊕○○ LOW¶	
Technical complications: screw loosening	81 per 1000	217 per 1000 (76 to 916)	RR 2.66 (0.93 to 7.60)	169 (3 RCTs)	⊕⊕○○ LOW¶	
Technical complications: crown loosening, decementation and chipping	27 per 1000	33 per 1000 (9 to 120)	RR 1.22 (0.33 to 4.49)	223 (5 RCTs)	⊕⊕○○ LOW¶	

*The risk in the intervention group (and its 95% CI) is based on the assumed risk in the comparison group and the **relative effect** of the intervention (and its 95% CI).

†Assumed risk is based on the overall event rate in the control groups of the included studies.

‡Downgraded one level due to serious risks of bias.

§Downgraded two levels due to serious risks of bias and serious inconsistency.

¶Downgraded two levels due to serious risks of bias and imprecision.

GRADE, Grading of Recommendations Assessment, Development, and Evaluation; RCTs, randomised controlled trials; RR, risk ratio.

complications. In this study, incidence of other biological and technical complications was similar between the two groups. Implant migration into the sinus, often with the co-occurrence of sinus infection, had a higher prevalence in the elevation group. When implant migration occurs, implants may be removed, thus leading to implant failure.⁶⁷ Technical complications, including screw loosening, crown loosening and chipping, were mainly associated with inappropriate loading, which could be resolved by improving supra rehabilitation structure. In addition, for short implants (≤ 6 mm), risks relating to reduced length could be partially alleviated by improving the design of the implants⁶⁸ or increasing their diameter. With respect to the prevalence and severity of adverse events, the use of short implants (≤ 6 mm) was acceptable and was a promising alternative to sinus floor elevation.

The present study has several strengths. First, we conducted a comprehensive literature search, and all included studies were RCTs. Second, participant was used as the unit of analysis to ensure logical statistical syntheses and relevant interpretations. Third, subgroup analysis by follow-up length and categories of complications was performed to reduce bias across studies. However, the evidence included in this systematic review was only of moderate or low quality. Serious risks of bias were found within and across studies. The number of participants and the follow-up period were limited. Due to limited data and methodological heterogeneity among studies, data synthesis for patient satisfaction was not performed. We suggest that researchers in this field carry out more well-designed, long-term and large-scale RCTs to provide

high-quality evidence regarding the effects of short implants (≤ 6 mm).

CONCLUSIONS

Within its limitations, the present review suggests that the survival rate of maxillary short implants (≤ 6 mm) was comparable to that of longer implants (≥ 10 mm) with sinus floor elevation. However, short implants (≤ 6 mm) show significantly less MBL and postsurgery reactions. Short implants (≤ 6 mm) are, therefore, the promising alternative to sinus floor elevation for posterior maxilla with insufficient bone volume. Additional high-quality studies are needed to evaluate the long-term effectiveness and safety of short implants (≤ 6 mm).

Author affiliations

¹Hubei-MOST KLOS & KLOBM, School & Hospital of Stomatology, Wuhan University, Wuhan, China

²Department of Oral Implantology, School & Hospital of Stomatology, Wuhan University, Wuhan, China

³Centre for Evidence-Based Stomatology, School & Hospital of Stomatology, Wuhan University, Wuhan, China

⁴Cochrane Oral Health, Division of Dentistry, The University of Manchester, Manchester, UK

Contributors Conception and design of the review: FH and BS. Methodology: FH. Drafting protocol, performing search strategies, literature searches, literature screening and data extraction: QY and XW. Original draft preparation: XW. Reviewing and editing draft: MS, QY, FH and BS. All authors critically reviewed and revised the manuscript and approved the final version for publication.

Funding This research was funded by the National Clinical Key Specialties Construction Project ((2013)544), the National Health Commission of China.

Competing interests None declared.

Patient consent for publication Not required.

Provenance and peer review Not commissioned; externally peer reviewed.

Data availability statement There are no data in this work.

Open access This is an open access article distributed in accordance with the Creative Commons Attribution Non Commercial (CC BY-NC 4.0) license, which permits others to distribute, remix, adapt, build upon this work non-commercially, and license their derivative works on different terms, provided the original work is properly cited, appropriate credit is given, any changes made indicated, and the use is non-commercial. See: <http://creativecommons.org/licenses/by-nc/4.0/>.

ORCID iD

Fang Hua <http://orcid.org/0000-0002-2438-5924>

REFERENCES

- Spitznagel FA, Horvath SD, Gierthmuehlen PC. Prosthetic protocols in implant-based oral rehabilitations: a systematic review on the clinical outcome of monolithic all-ceramic single- and multi-unit prostheses. *Eur J Oral Implantol* 2017;10 Suppl 1:89–99.
- Pjetursson BE, Thoma D, Jung R, et al. A systematic review of the survival and complication rates of implant-supported fixed dental prostheses (FDPs) after a mean observation period of at least 5 years. *Clin Oral Implants Res* 2012;23 Suppl 6:22–38.
- Davaranpanah M, Martinez H, Etienne D, et al. A prospective multicenter evaluation of 1,583 3i implants: 1- to 5-year data. *International Journal of Oral & Maxillofacial Implants* 2002;17:820–8.
- Esposito M, Felice P, Worthington HV. *Interventions for replacing missing teeth: augmentation procedures of the maxillary sinus*. John Wiley & Sons, Ltd, 2014.
- Morand M, Irinakakis T. The challenge of implant therapy in the posterior maxilla: providing a rationale for the use of short implants. *Journal of Oral Implantology* 2007;33:257–66.
- Malchiodi L, Caricasulo R, Cucchi A, et al. Evaluation of ultrashort and longer implants with Microrough surfaces: results of a 24- to 36-month prospective study. *Int J Oral Maxillofac Implants* 2017;32:171–9.
- Danesh-Sani SA, Loomer PM, Wallace SS. A comprehensive clinical review of maxillary sinus floor elevation: anatomy, techniques, biomaterials and complications. *Br J Oral Maxillofac Surg* 2016;54:724–30.
- Brånemark PI, Hansson BO, Adell R, et al. Osseointegrated implants in the treatment of the edentulous jaw. experience from a 10-year period. *Scand J Plast Reconstr Surg Suppl* 1977;16:1–132.
- Cha H-S, Kim J-W, Hwang J-H, et al. Frequency of bone graft in implant surgery. *Maxillofac Plast Reconstr Surg* 2016;38.
- Starch-Jensen T, Aludden H, Hallman M, et al. A systematic review and meta-analysis of long-term studies (five or more years) assessing maxillary sinus floor augmentation. *Int J Oral Maxillofac Surg* 2018;47:103–16.
- Starch-Jensen T, Schou S. Maxillary sinus membrane elevation with simultaneous installation of implants without the use of a graft material: a systematic review. *Implant Dent* 2017;26:621–33.
- Lee S-A, Lee C-T, Fu MM, et al. Systematic review and meta-analysis of randomized controlled trials for the management of limited vertical height in the posterior region: short implants (5 to 8 mm) vs longer implants (> 8 mm) in vertically augmented sites. *Int J Oral Maxillofac Implants* 2014;29:1085–97.
- Pjetursson BE, Tan WC, Zwahlen M, et al. A systematic review of the success of sinus floor elevation and survival of implants inserted in combination with sinus floor elevation. *J Clin Periodontol* 2008;35:216–40.
- Tan WC, Lang NP, Zwahlen M, et al. A systematic review of the success of sinus floor elevation and survival of implants inserted in combination with sinus floor elevation Part II: transalveolar technique. *J Clin Periodontol* 2008;35:241–54.
- Thoma DS, Zeltner M, Hüsler J, et al. EAO Supplement Working Group 4 - EAO CC 2015 Short implants versus sinus lifting with longer implants to restore the posterior maxilla: a systematic review. *Clinical Oral Implants Res* 2015;26:154–69.
- Lundgren S, Cricchio G, Hallman M, et al. Sinus floor elevation procedures to enable implant placement and integration: techniques, biological aspects and clinical outcomes. *Periodontology* 2000 2017;73:103–20.
- Huang H-L, Fuh L-J, Ko C-C, et al. Biomechanical effects of a maxillary implant in the augmented sinus: a three-dimensional finite element analysis. *Int J Oral Maxillofac Implants* 2009;24:455–62.
- Lin Y-H, Yang Y-C, Wen S-C, et al. The influence of sinus membrane thickness upon membrane perforation during lateral window sinus augmentation. *Clin Oral Implants Res* 2016;27:612–7.
- Monje A, Suarez F, Galindo-Moreno P, et al. A systematic review on marginal bone loss around short dental implants (<10 mm) for implant-supported fixed prostheses. *Clin Oral Implants Res* 2014;25:1119–24.
- Nielsen HB, Schou S, Isidor F, et al. Short implants (≤ 8 mm) compared to standard length implants (> 8 mm) in conjunction with maxillary sinus floor augmentation: a systematic review and meta-analysis. *Int J Oral Maxillofac Surg* 2019;48:239–49.
- Uehara PN, Matsubara VH, Igai F, et al. Short Dental Implants (≤ 7 mm) Versus Longer Implants in Augmented Bone Area: A Meta-Analysis of Randomized Controlled Trials. *Open Dent J* 2018;12:354–65.
- Al Amri MD, Abduljabbar TS, Al-Johany SS, et al. Comparison of clinical and radiographic parameters around short (6 to 8 mm in length) and long (11 mm in length) dental implants placed in patients with and without type 2 diabetes mellitus: 3-year follow-up results. *Clin Oral Implants Res* 2017;28:1182–7.
- Urdaneta RA, Daher S, Leary J, et al. The survival of ultrashort locking-taper implants. *Int J Oral Maxillofac Implants* 2012;27:644–54.
- Markose J, Eshwar S, Srinivas S, et al. Clinical outcomes of ultrashort sloping shoulder implant design: a survival analysis. *Clin Implant Dent Relat Res* 2018;20:646–52.
- Fan T, Li Y, Deng W-W, et al. Short Implants (5 to 8 mm) Versus Longer Implants (> 8 mm) with Sinus Lifting in Atrophic Posterior Maxilla: A Meta-Analysis of RCTs. *Clin Implant Dent Relat Res* 2017;19:207–15.
- Deporter D, Ogiso B, Sohn D-S, et al. Ultrashort sintered porous-surfaced dental implants used to replace posterior teeth. *J Periodontol* 2008;79:1280–6.
- Sahrmann P, Naenni N, Jung RE, et al. Success of 6-mm implants with Single-Tooth restorations: a 3-year randomized controlled clinical trial. *Journal of Dental Research* 2016;95:623–8.
- Felice P, Checchi L, Barausse C, et al. Posterior jaws rehabilitated with partial prostheses supported by 4.0 X 4.0 mm or by longer

- implants: one-year post-loading results from a multicenter randomised controlled trial. *Eur J Oral Implantol* 2016;9:35–45.
- 29 Bolle C, Felice P, Barausse C, et al. 4 MM long vs longer implants in augmented bone in posterior atrophic jaws: 1-year post-loading results from a multicentre randomised controlled trial. *Eur J Oral Implantol* 2018;11:31–47.
 - 30 Calvo-Guirado JL, López Torres JA, Dard M, et al. Evaluation of extrashort 4-mm implants in mandibular edentulous patients with reduced bone height in comparison with standard implants: a 12-month results. *Clin Oral Implants Res* 2016;27:867–74.
 - 31 Liberati A, Altman DG, Tetzlaff J, et al. The PRISMA statement for reporting systematic reviews and meta-analyses of studies that evaluate healthcare interventions: explanation and elaboration. *BMJ* 2009;339.
 - 32 Higgins JPT GS. Cochrane Handbook for systematic reviews of interventions version 5.1.0: the Cochrane collaboration, 2011. Available: www.cochrane-handbook.org
 - 33 Egger M, Davey SG, Schneider M, et al. Bias in meta-analysis detected by a simple, graphical test. *Bmj British Medical Journal* 1997;316:469–71.
 - 34 Esposito M, Ardebili Y, Worthington HV, et al. Interventions for replacing missing teeth: different types of dental implants. *Cochrane Database Syst Rev* 2014;26.
 - 35 Borenstein M, Hedges LV, Higgins JPT, et al. A basic introduction to fixed-effect and random-effects models for meta-analysis. *Res. Synth. Method* 2010;1:97–111.
 - 36 Esposito M, Grusovin MG, Maghaireh H, et al. Interventions for replacing missing teeth: different times for loading dental implants. *Cochrane Database of Systematic Reviews* 2013;16.
 - 37 Esposito M, Worthington HV, Cochrane Oral Health Group. Interventions for replacing missing teeth: hyperbaric oxygen therapy for irradiated patients who require dental implants. *Cochrane Database Syst Rev* 2013;43.
 - 38 Esposito M, Felice P, Worthington HV, et al. Interventions for replacing missing teeth: augmentation procedures of the maxillary sinus. *Cochrane Database of Systematic Reviews* 2014;2.
 - 39 Tolentino da Rosa de Souza P, Binhame Albini Martini M, Reis Azevedo-Alanis L. Do short implants have similar survival rates compared to standard implants in posterior single crown?: a systematic review and meta-analysis. *Clin Implant Dent Relat Res* 2018;20:890–901.
 - 40 Andrews J, Guyatt G, Oxman AD, et al. Grade guidelines: 14. going from evidence to recommendations: the significance and presentation of recommendations. *J Clin Epidemiol* 2013;66:719–25.
 - 41 Gastaldi G, Felice P, Pistilli V, et al. Posterior atrophic jaws rehabilitated with prostheses supported by 5 × 5 mm implants with a nanostructured calcium-incorporated titanium surface or by longer implants in augmented bone. 3-year results from a randomised controlled trial. *Eur J Oral Implantol*
 - 42 Gastaldi G, Felice P, Pistilli R, et al. Short implants as an alternative to crestal sinus lift: a 3-year multicentre randomised controlled trial. *Eur J Oral Implantol* 2017;10:391–400.
 - 43 Guljé FL, Raghoobar GM, Vissink A, et al. Single Crowns in the resorbed posterior maxilla supported by either 6-mm implants or by 11-mm implants combined with sinus floor elevation surgery: a 1-year randomised controlled trial. *Eur J Oral Implantol* 2014;7:247–55.
 - 44 Pohl V, Thoma DS, Sporniak-Tutak K, et al. Short dental implants (6 mm) versus long dental implants (11–15 mm) in combination with sinus floor elevation procedures: 3-year results from a multicentre, randomized, controlled clinical trial. *J Clin Periodontol* 2017;44:438–45.
 - 45 Felice P, Barausse C, Pistilli V, et al. Posterior atrophic jaws rehabilitated with prostheses supported by 6 MM long × 4 MM wide implants or by longer implants in augmented bone. 3-year post-loading results from a randomised controlled trial. *Eur J Oral Implantol* 2018;11:175–87.
 - 46 Bechara S, Kubilius R, Veronesi G, et al. Short (6-mm) dental implants versus sinus floor elevation and placement of longer (≥10-mm) dental implants: a randomized controlled trial with a 3-year follow-up. *Clin Oral Implants Res* 2017;28:1097–107.
 - 47 Lombardo G, Pighi J, Marincola M, et al. Cumulative success rate of short and ultrashort implants supporting single Crowns in the posterior maxilla: a 3-year retrospective study. *Int J Dent* 2017;2017:1–10.
 - 48 Lai H-C, Si M-S, Zhuang L-F, et al. Long-term outcomes of short dental implants supporting single crowns in posterior region: a clinical retrospective study of 5–10 years. *Clin Oral Implants Res* 2013;24:230–7.
 - 49 Himmlová L, Dostálová Tat'jana, Kácvský A, et al. Influence of implant length and diameter on stress distribution: a finite element analysis. *J Prosthet Dent* 2004;91:20–5.
 - 50 Moriwaki H, Yamaguchi S, Nakano T, et al. Influence of implant length and diameter, Bicortical anchorage, and sinus augmentation on bone stress distribution: three-dimensional finite element analysis. *Int J Oral Maxillofac Implants* 2016;31:e84–91.
 - 51 Kim S, Jung U-W, Cho K-S, et al. Retrospective radiographic observational study of 1692 Straumann tissue-level dental implants over 10 years: I. implant survival and loss pattern. *Clin Implant Dent Relat Res* 2018;20:860–6.
 - 52 Lin G, Ye S, Liu F, et al. A retrospective study of 30,959 implants: risk factors associated with early and late implant loss. *J Clin Periodontol* 2018;45:733–43.
 - 53 Tabassum A, Meijer GJ, Wolke JGC, et al. Influence of the surgical technique and surface roughness on the primary stability of an implant in artificial bone with a density equivalent to maxillary bone: a laboratory study. *Clin Oral Implants Res* 2009;20:327–32.
 - 54 Dos Santos MV, Elias CN, Cavalcanti Lima JH. The effects of superficial roughness and design on the primary stability of dental implants. *Clin Implant Dent Relat Res* 2011;13:215–23.
 - 55 Falco A, Berardini M, Trisi P. Correlation between implant geometry, implant surface, insertion torque, and primary stability: in vitro biomechanical analysis. *Int J Oral Maxillofac Implants* 2018;33:824–30.
 - 56 Jain N, Gulati M, Garg M, et al. Short implants: new horizon in implant dentistry. *J Clin Diagn Res* 2016;10:Ze14–17.
 - 57 Uehara PN, Matsubara VH, Igai F, et al. Short dental implants (≤7mm) versus longer implants in augmented bone area: a meta-analysis of randomized controlled trials. *Open Dent J* 2018;12:354–65.
 - 58 Anitua E, Alkhraist MH, Piñas L, et al. Implant survival and crestal bone loss around extra-short implants supporting a fixed denture: the effect of crown height space, crown-to-implant ratio, and offset placement of the prosthesis. *Int J Oral Maxillofac Implants* 2014;29:682–9.
 - 59 Quirynen M, Naert I, van Steenberghe D. Fixture design and overload influence marginal bone loss and future success in the BranemarkR system. *Clin Oral Implants Res* 1992;3:104–11.
 - 60 Blanes RJ. To what extent does the crown-implant ratio affect the survival and complications of implant-supported reconstructions? A systematic review. *Clin Oral Implants Res* 2009;20:67–72.
 - 61 Verri FR, Batista VEdeS, Santiago JF, et al. Effect of crown-to-implant ratio on peri-implant stress: a finite element analysis. *Materials Science and Engineering: C* 2014;45:234–40.
 - 62 Quaranta A, Piemontese M, Rappelli G, et al. Technical and biological complications related to crown to implant ratio. *Implant Dent* 2014;23:180–7.
 - 63 Urdaneta RA, Rodriguez S, McNeil DC, et al. The effect of increased crown-to-implant ratio on single-tooth locking-taper implants. *Int J Oral Maxillofac Implants* 2010;25:729–43.
 - 64 Tawil G, Aboujaoude N, Younan R. Influence of prosthetic parameters on the survival and complication rates of short implants. *International Journal of Oral & Maxillofacial Implants* 2006;21:275–82.
 - 65 Schneider D, Witt L, Hämmerle CHF. Influence of the crown-to-implant length ratio on the clinical performance of implants supporting single crown restorations: a cross-sectional retrospective 5-year investigation. *Clin Oral Implants Res* 2012;23:169–74.
 - 66 Thoma DS, Cha J-K, Jung U-W. Treatment concepts for the posterior maxilla and mandible: short implants versus long implants in augmented bone. *J Periodontol Implant Sci* 2017;47:2–12.
 - 67 Manor Y, Anavi Y, Gershonovitch R, et al. Complications and management of implants migrated into the maxillary sinus. *Int J Periodontics Restorative Dent* 2018;38:e112–8.
 - 68 Mangano F, Frezzato I, Frezzato A, et al. The effect of Crown-to-Implant ratio on the clinical performance of Extra-Short Locking-Taper implants. *J Craniofac Surg* 2016;27:675–81.