BMJ Open Knowledge and attitudes of syphilis and syphilis pre-exposure prophylaxis (PrEP) among men who have sex with men in Vancouver, Canada: a qualitative study

Ronita Nath ^{1,2,3} Troy Grennan,^{1,3} Robin Parry,¹ Fahmy Baharuddin,¹ James P Connell,¹ Jason Wong,^{1,4} Daniel Grace⁵

To cite: Nath R, Grennan T, Parry R, *et al.* Knowledge and attitudes of syphilis and syphilis pre-exposure prophylaxis (PrEP) among men who have sex with men in Vancouver, Canada: a qualitative study. *BMJ Open* 2019;**9**:e031239. doi:10.1136/bmjopen-2019-031239

► Prepublication history and additional material for this paper are available online. To view these files, please visit the journal online (<http://dx.doi.org/10.1136/bmjopen-2019-031239>).

Received 23 April 2019

Revised 17 October 2019

Accepted 04 November 2019

© Author(s) (or their employer(s)) 2019. Re-use permitted under CC BY-NC. No commercial re-use. See rights and permissions. Published by BMJ.

For numbered affiliations see end of article.

Correspondence to

Dr Daniel Grace;
daniel.grace@utoronto.ca

ABSTRACT

Objectives In British Columbia, Canada, syphilis is at record-high rates, with over 80% of cases in 2017 seen in gay, bisexual and other men who have sex with men (GBM). The syphilis epidemic is of particular concern for those living with HIV, since syphilis may lead to more serious complications in this population. We sought to explore syphilis-related knowledge and attitudes around biomedical prevention options for syphilis, with the goal of informing effective strategies to prevent syphilis.

Design We conducted a qualitative study consisting of in-depth, individual interviews from December 2016 to June 2017. Our interviews focused on participants' knowledge about syphilis and perceptions regarding syphilis pre-exposure prophylaxis (PrEP). Interviews were analysed using Grounded Theory.

Participants Twenty-five GBM were interviewed (64% white; median age: 43 years), including men living with HIV and/or with a history of syphilis.

Setting Vancouver, British Columbia.

Results Five interrelated themes emerged. First, GBM were aware of the local syphilis epidemic. Second, syphilis-related knowledge differed according to syphilis and HIV serostatus. Third, competing ideas emerged regarding men's concerns about syphilis. While our participants expressed concern about getting syphilis, they also described the importance of sexual pleasure. Fourth, many participants said that syphilis was not perceived to be alarming; preventing HIV infection remained a primary concern for many. Finally, while syphilis PrEP was appealing to those living with HIV or a prior syphilis diagnosis, others were concerned about antibiotic resistance, cost and side effects.

Conclusions Our participants organised their safer sex strategies around HIV, not syphilis. Although syphilis-related knowledge was relatively high among GBM living with HIV and those with a prior syphilis diagnosis, this knowledge did not appear to be related with safer sexual practices, such as increased condom use. This work highlights the importance of examining other potential prevention solutions, such as syphilis PrEP.

Strengths and limitations of this study

- This study included a diverse sample of gay, bisexual and other men who have sex with men who varied in age, ethnicity, income and sexually transmitted infections diagnosis.
- Interviewers were chosen from within the gay community and outside the community; similar findings were noted across interviewers, suggesting minimal interviewer bias.
- The level of health literacy among the participants was difficult to ascertain in the qualitative study because several participants provided vague descriptions of the infection, making the extent of their knowledge unclear.
- There may be a lack of diversity in the risk profile of the sample because recruitment was limited to online applications and clinics and did not include gay venues like bathhouses.

INTRODUCTION

Recent advances in biomedical prevention like HIV pre-exposure prophylaxis (PrEP) and treatment as prevention (TasP) have had dramatic impacts on HIV incidence.^{1 2} At the same time, there has been a resurging epidemic of bacterial sexually transmitted infections (STIs) and STI-related sequelae.^{3 4} In British Columbia (BC), Canada's western-most province, syphilis is at unprecedented levels, with a rate of 10.8 per 100 000 in 2017.⁵ Risk factors for syphilis include lack of condom use, unprotected oral sex and recreational drug use.^{6 7} Gay, bisexual and other men who have sex with men (GBM) continue to be disproportionately affected by the current infectious syphilis epidemic in BC, constituting 82.9% of all cases in 2017.⁸ The number of infectious syphilis cases (primary, secondary and early latent syphilis)

among GBM has been steadily increasing since 2011. Among GBM syphilis cases where HIV status was known, 40% were living with HIV.⁸ Forty-six per cent of HIV cases in BC are among GBM.⁹ The emergence of syphilis has particular implications for GBM living with HIV, who are often at risk for more serious complications from bacterial STIs, including more severe cutaneous lesions in early syphilis,¹⁰ higher likelihood of ocular syphilis¹¹ and infection of the lymphatic system known as lymphogranuloma venereum caused by the bacteria *Chlamydia trachomatis*.¹²

Interventions to address the syphilis epidemic have mainly focused on extending testing and treatment, enhancing partner notification and encouraging GBM to modify their sexual behaviours, such as using condoms.^{5,13} Despite these efforts, syphilis incidence continues to rise and novel approaches for responding to the syphilis epidemic are needed. Addressing the rates of syphilis among GBM requires a nuanced understanding of the knowledge and attitudes GBM have towards syphilis and syphilis prevention strategies. Although there has been a growing research base that qualitatively investigates the HIV knowledge, sexual practices and multifactorial impacts of biomedical advancements in HIV prevention among GBM living with HIV,^{14–16} there has been a dearth of qualitative work specifically examining syphilis and views towards syphilis prevention for GBM.

To date, limited qualitative studies have investigated syphilis knowledge and attitudes among GBM^{17,18}; Stahlman *et al* examined acceptable syphilis interventions among GBM with repeat syphilis infections.¹⁹ According to this research, there is a strong stigma associated with syphilis among many GBM.^{17,18} GBM perceive syphilis to be a ‘dirty’ disease,^{17,18} which hinders syphilis disclosure and partner notification,¹⁸ despite considerable knowledge about syphilis symptoms, transmission and consequences.¹⁷ Syphilis prevention strategies preferred by GBM include increased accessibility and testing strategies for syphilis, as well as prophylactic treatment, although participants expressed concerns about potential side effects with such treatment.¹⁹ This body of literature included only men who had been previously diagnosed with syphilis.

There is a critical need to better understand how syphilis is perceived among diverse GBM in Canada. BC is a particularly important context to study the syphilis epidemic due to the shifting knowledge around seroadaptive behaviours,^{15,16} its historical focus on early HIV detection and TasP²⁰ and the recent public funding of HIV PrEP.^{21,22} The high uptake of HIV PrEP, recent pilot studies demonstrating promise of using daily doxycycline for syphilis chemoprophylaxis (henceforth, syphilis PrEP)^{23,24} and the qualitative study in California that found that GBM are interested in syphilis PrEP as a syphilis prevention strategy,¹⁹ invite further exploration.

Our overarching objective was to inductively explore syphilis-related knowledge and attitudes around biomedical prevention options for syphilis, with the goal of informing effective strategies to prevent syphilis.

METHODS

Recruitment

We conducted one-on-one interviews with 25 GBM in the Greater Vancouver area between December 2016 and June 2017 as the initial phase of a sequential, mixed-methods study. A purposive sampling strategy²⁵ was used to recruit participants from clinic sites of the BC Centre for Disease Control and through the online sexual networking applications Scruff and Squirt. Posters in the clinics and advertisements on the online applications outlining the eligibility criteria were used to invite participants into the study. Individuals were eligible to participate if they identified as male, were 18 years of age or older, had sex with other men in the last 6 months, were fluent in English and were available to interview in-person.

Data collection and analysis

Data were collected in individual, semi-structured interviews that lasted between 45 min and 1 hour. Three trained interviewers conducted all the interviews. Interviews were audio-recorded and transcribed verbatim. The interview guide (online supplementary file) was developed in consultation with our Community Advisory Board. The guide sought to explore questions related to the social determinants of health, and covered the following domains: STI-related knowledge, beliefs and attitudes; perceptions of STI risk; the acceptability of PrEP for syphilis; experiences with healthcare providers; measures taken to prevent STIs; the role of substance use in sexual activity and decision-making; and the role of social networks. In this paper, we used a Grounded theory²⁶ approach to help us explicate what we learnt from our participants about their syphilis knowledge and attitudes, and their acceptability of syphilis PrEP. Our use of Grounded theory helped to ensure that our conceptualisation of these sexual health topics was formed from the participants’ interpretations and meaning.²⁷

Data were systematically analysed using QSR NVivo 10 according to the principles of Grounded theory. Grounded theory is an inductive methodology where codes are constructed from the data and grouped into categories. Categories are then integrated into a theoretical framework with coding occurring at three levels: open, axial and selective coding.²⁸ In open coding, we reviewed the transcripts several times and labelled key concepts. In axial coding, we identified the relationships among the codes and combined categories with subcategories. The selective-coding stage involved relating other categories to the core category and refining the theoretical explanation.

To ensure methodological rigour, the authors reviewed five transcripts independently and subsequently as a coding team to familiarise themselves with the data set, note initial trends and ideas and create a provisional codebook. These codes were developed from key themes that emerged during the review of transcripts. Codes were applied to the entire data set and refined as needed. Memos were recorded throughout the iterative process.

of coding to capture analytical insights about emerging themes and relationships across the narratives.

Participants received an honorarium of US\$30.

Patient and public involvement

A Community Advisory Board guided the research activities, including providing feedback on the development of the interview guide, the interviews, analysis and dissemination of the findings.

RESULTS

Characteristics of the 25 GBM we interviewed are provided in table 1. Most participants described being single and no participants had previously taken PrEP for HIV prevention. The majority of the participants described having active sexual lives. Some of the single participants reported having 'sex buddies', or partners they would have oral and/or anal sex with on a regular basis. The vast majority of the participants described using sexual networking applications to meet partners. Half of the participants also described visiting bathhouses recently. Most participants reported having condomless anal sex in the past 6 months.

Themes

Five overarching themes were identified in the analysis: (1) awareness of the local syphilis epidemiology, (2) syphilis-related health literacy, (3) competing ideas about concern for syphilis and the need to engage in pleasurable sex, (4) HIV infection as the most worrying health issue and (5) concerns, stigma and perceived benefits of syphilis PrEP. These interrelated domains are explored below.

Awareness of the local syphilis epidemiology

Most participants were aware of the recent surge in syphilis rates in the province, and they knew the epidemic was disproportionately impacting GBM. Some men talked about the apparent ubiquity or inevitability of syphilis if one was a sexually active gay man. A 31-year-old participant living with HIV who had been diagnosed with syphilis previously expressed this sentiment: 'Once you're gay, it's just like everyone has syphilis...'. In recent years, participants noted that they had seen in the media and in their doctors' offices that syphilis was 'hitting at all-time highs'. However, several participants said that it was not a topic discussed among their gay friends and sexual partners. One 57-year-old HIV-negative participant who had never been diagnosed with syphilis expressed that he did not 'get the sense that anyone's overly concerned'. He added, 'People don't really ask specifics about a specific STD online... No one seems to bring it up as a concern'. Many participants shared that they were not very worried about the epidemic because they had come to understand that syphilis could be treated with antibiotics. Another participant told us he found it hard to worry about the syphilis epidemic because he did not know how to prevent it. He

Table 1 Characteristics of participants

Variable	N (out of 25 participants)
Age	
18–29 years	7
30–40 years	5
41–51 years	7
52+	6
Sex at birth	
Male	24
Female	1
Gender	
Male	22
Female	0
Transgender	3
Ethnicity	
Asian	3
South Asian	1
White	16
Latino/Hispanic	1
Mixed	1
Black	1
Aboriginal	2
Sexual orientation	
Gay	19
Straight	1
Queer	1
Other	4
Bisexual	0
Highest level of education	
Did not complete high school	2
High school	5
College	15
Graduate	3
Annual income	
<\$20 000	5
\$20 000–\$39 000	8
\$40 000–\$59 000	7
\$60 000–\$79 000	3
\$80 000–\$99 000	1
Prefer not to answer	1
Prior syphilis diagnosis	
Yes	9
No	15
Missing	1
HIV status	
Negative	15

Continued

Table 1 Continued

Variable	N (out of 25 participants)
Positive	8
Unsure	2
Number of men that the participant had sex with in past 6 months	
1–5	11
6–10	8
11–15	2
16–20	1
20+	2
Missing	1
Number of participants who have ever taken non-occupational post exposure prophylaxis (nPEP)	2

went on to say that ‘It’s not like HIV where you can use a condom’.

The majority of participants explained that the rising rates in syphilis had not impacted their sexual behaviour or the way they met with prospective partners. However, a few participants did share that the epidemic had impacted their sex life. For example, a 44-year-old HIV-negative participant who had never been diagnosed with syphilis and previously frequented bathhouses, said that he stopped going several years ago when he found out that bathhouses ‘were the accused, or the guilty party or the cause of the (syphilis) epidemic in a way’. Another participant shared that he stopped having sex with strangers because of the syphilis epidemic, and another said he avoided having condomless anal sex with one of his partners ‘who slept around’ when syphilis rates started rising. One 51-year-old HIV-negative participant who had never been diagnosed with syphilis expressed his frustration that testing was not required for individuals at risk:

They (governments, healthcare organisations) need to be more aggressive, not just saying, oh, syphilis is on the rise and then leave it up to the person to get tested. Like if you’re sexually active, get tested, period; don’t just say oh, I will never get it, and the next thing you know, you’ve got it.

Syphilis-related health literacy

Perhaps not surprisingly, participants with a prior syphilis diagnosis or who were living with HIV, reported greater levels of syphilis-related knowledge than HIV-negative participants who had never had syphilis. Many in the former group discussed syphilis symptoms, transmission and treatment in the context of their own diagnosis. For example, a 31-year-old participant living with HIV explained how becoming infected with syphilis led him to

seek information about syphilis: ‘I didn’t know that there (were) four stages until I did some more research on it... I was in the hospital for a month, and I didn’t realise that I had third stage syphilis... So, with syphilis it’s passed like HIV or Hep C, it’s sexual. It’s passed through sex’.

Several participants with prior syphilis diagnoses made strong assertions of their high level of knowledge regarding syphilis, but some lacked accuracy on particular points. For instance, a few participants falsely believed that syphilis was a virus and was becoming resistant to antibiotics.

Many participants described syphilis as an ‘old’ infection, referring to an infection that has been around for a very long time. Among these participants, several understood that syphilis was a resurging problem as well, particularly in the gay community, as described in the accounts above. For example, a 30-year-old living with HIV and with a prior syphilis diagnosis explained, ‘It (syphilis) is very old (laughs)... (T)here’s been a huge increase especially in the gay community in people contracting syphilis. If it goes untreated, it can make you really sick’.

Most HIV-negative participants with no history of syphilis disclosed having very limited syphilis-related knowledge. For these participants, the differences between syphilis, chlamydia and gonorrhoea were blurry. For example, a 25-year-old HIV-negative participant who had never had syphilis told us that he did not know much about the infection, and that his ‘knowledge about syphilis, chlamydia and gonorrhoea were all the same; they burn when you pee (laughs), and you pretty much take the same meds to get rid of them’.

Competing ideas

Syphilis-related concerns emerged as a complex theme in men’s accounts. Although all of our participants expressed that they did not want to become infected with syphilis, heterogeneity existed around the level of concern for syphilis. One recurring theme in men’s interviews was the view that syphilis was a minor health problem that could be treated easily, leading few participants to have heightened concern about the infection. For example, a 25-year-old HIV-negative participant who had never been diagnosed with syphilis exclaimed, ‘(I am) not concerned. I don’t care. It’s like a cold. You take a couple of pills and it’s gone’. Similarly, a 44-year-old HIV-negative participant who had been diagnosed with syphilis previously said, ‘I just think you can get a shot and then you’re fine, with syphilis or any other... There’s no cure for HIV, so no one really talks about the stuff you can get cured from, because what’s the point?’

Among those who were concerned about getting syphilis, participants shared that it was difficult to change their sexual behaviour to prevent infection because sexual pleasure was also very important to them. For many, the need to engage in pleasurable sex outweighed concerns they had about syphilis during a sexual encounter. For instance, when asked whether his concern for syphilis impacted his decision to use condoms, or to be more

selective in choosing his sexual partners, a 57-year-old HIV-negative participant without a prior syphilis diagnosis explained: 'I'd say it probably doesn't affect that. Like I said, I have some concern about it, but if I'm looking to have sex with somebody it's not something that I'm thinking or worrying about at that time'.

The two men in the study who were living with HIV and had never been diagnosed with syphilis said that they were unconcerned about syphilis because their HIV status was a more significant issue. One of them, aged 54, put it this way: 'Well, what's the worst that's going to happen to me? Nothing really, so what's the big deal? I'm already really sick'.

Most participants with prior syphilis diagnoses were more concerned about a syphilis infection because of their traumatic experiences with the STI. Several such participants described feeling very sick when they had syphilis previously, and many reported that the treatment was extremely painful. They emphasised that they wanted to avoid a subsequent syphilis infection. Despite their negative experiences with syphilis, and their desire to avoid subsequent infections, these participants reported they did not feel inclined to make long-term behavioural changes, such as using condoms for all sexual encounters, to prevent further infections.

'It's HIV versus not HIV'

HIV surfaced as the health issue of primary concern to participants irrespective of HIV status or history of syphilis diagnosis. Men highlighted that HIV was of primary concern to them, and that the other STIs were relatively minor health problems compared with HIV. Some participants expressed that HIV would be the only STI they would take precautions against because unlike syphilis, HIV is a lifelong chronic health condition. For example, a 25-year-old HIV-negative participant who had never been diagnosed with syphilis said: 'HIV would be the only one that I would not want to get, for obvious reasons, but it wouldn't necessarily change my decision to have sex with someone. I would just use a condom'.

As discussed above, because syphilis and many of the other STIs can be cured, they were not considered to be very important health concerns. When asked how he felt about syphilis compared with other STIs, a 26-year-old HIV-negative participant without a prior syphilis diagnosis put it quite simply: 'It's HIV versus not HIV'. Some participants who had been diagnosed with syphilis expressed their frustration with these views. A 44-year-old HIV-negative participant exclaimed, 'People that haven't had syphilis don't realise how bad syphilis is, and it's not a wide talked about thing, right? It's always HIV status and that's pretty much where it ends'.

Syphilis PrEP: concerns, stigma and perceived benefits

While men living with HIV or with a prior diagnosis of syphilis were very interested in syphilis PrEP, many HIV-negative participants who did not have a history of syphilis

expressed several concerns about taking a daily antibiotic for syphilis prevention.

One of the major stated concerns was antibiotic resistance. A 23-year-old HIV-negative participant who had never been diagnosed with syphilis expressed his confusion with why someone would take a daily antibiotic to prevent syphilis: 'There are super bugs. There is antibiotic resistance. Why would you do that? I mean... antibiotics also takes out the bacteria in your system like your digestive bacteria and that's not healthy'. HIV-negative participants who did not have a prior syphilis diagnosis were also concerned about the unknown side effects to taking a daily antibiotic.

Both HIV-negative participants and participants living with HIV cited the cost of syphilis PrEP as being a potential barrier in accessing the prophylaxis. Participants believed that since the cost of HIV PrEP was exorbitant (at the time of data collection, the cost of HIV PrEP was not publicly funded), the cost of syphilis PrEP would be as well.

The idea of sex-related stigma emerged when discussing syphilis PrEP. Some participants thought that syphilis PrEP was only meant for those who were 'very promiscuous' and believed that they did not fall in that category: syphilis PrEP might be appropriate for 'other' GBM. For example, a 30-year-old HIV-negative participant who had never been diagnosed with syphilis exclaimed that he was not 'sexually active enough' for the drug and that it was only meant for 'deviant(s)'.

On the other hand, many of the participants living with HIV or with a prior diagnosis of syphilis were interested in syphilis PrEP. They explained that syphilis PrEP would afford them the convenience of not having to worry about syphilis. This was the sentiment of a 31-year-old man living with HIV who had several traumatic encounters with syphilis, one of which caused him to hallucinate. He was keen on the idea of taking a daily antibiotic to prevent future syphilis infections:

A daily antibiotic? Sign me up! I mean, anything to have one less thing that I have to worry about would be great. It sucks that I've had syphilis three times. The second time was the most dangerous time because there was the third stage, so it was in my spinal fluid... I was almost going crazy. Well, actually I was going crazy, because I was seeing things, hearing things.

DISCUSSION

Our study offers significant insights into the syphilis-related health literacy of GBM, how they organise their safer sexual strategies and their willingness to take a chemoprophylactic drug to prevent syphilis. Men living with HIV or with a prior diagnosis of syphilis possessed a high level of perceived knowledge on syphilis compared with their HIV-negative counterparts and those without a prior syphilis diagnosis, possibly as a result of having

more exposure to the healthcare system. Although men living with HIV or with a prior diagnosis of syphilis were health literate, there did not appear to be an association between men's health literacy and sexual behaviour. For instance, men's accounts depicted that they organised their safer sexual strategies around HIV, not syphilis, even though many of them understood that there was a resurgence of syphilis and they could be impacted by it. This was also a finding noted by Holt *et al* among GBM in Sydney, Australia.²⁹ Men were more worried about HIV; some used condoms when having sex with men living with HIV, or they engaged in seroadaptive behaviours such as selecting their partners based on their HIV status. Participants explained that they typically did not show the same consideration to avoid a syphilis infection. Overall, concern for syphilis was low among the participants, particularly among HIV-negative GBM who had never been diagnosed with the infection.

Despite a high level of concern for HIV, and a strong desire to avoid HIV infection, men did not seem to be aware that syphilis was associated with an increased risk of HIV acquisition.³⁰ These findings have implications for programme design. For instance, since participants are mostly worried about HIV infection, syphilis messaging in syphilis prevention programme should more explicitly address the link between syphilis and HIV. This message should be presented in a thoughtful and sex-positive framing.

Men's discussions regarding syphilis PrEP revealed that a daily antibiotic holds promise as an acceptable intervention for men living with HIV and those with a prior syphilis diagnosis. Our findings indicating greater acceptance of syphilis PrEP among men living with HIV compared with HIV-negative men are consistent with that of other findings in the literature.²³ HIV-negative participants and those without a prior syphilis diagnosis were wary about syphilis PrEP and cited several barriers to using a daily antibiotic including unknown side effects, high costs, antibiotic resistance and sex-related stigma associated with PrEP. Many of these concerns can be addressed. First, doxycycline, the antibiotic which has shown to be potentially effective in preventing syphilis in a previous small pilot study,²⁴ has been widely used for decades, often on a long-term basis for acne treatment and malaria prevention and is typically well-tolerated.^{31–34} Second, doxycycline is inexpensive. Third, a study is currently underway in Vancouver, BC, examining the impact of doxycycline on antibiotic resistance in HIV-negative GBM.³⁵ Although many of the concerns surrounding syphilis PrEP can be alleviated, there still lies the need to examine on a larger scale whether daily doxycycline is a feasible option as a syphilis prevention tool, which we will assess in the 2019 National *Sex Now* survey,³⁶ a community-based, gay men's health survey. It is important to note that none of the participants reported taking HIV PrEP at the time of the interviews, which took place between 2016 to 2017. It was not until January 2018 that HIV PrEP became publicly-funded in BC, and within 6 months had been prescribed

to over 2000 people.²⁰ With increasing acceptance of biomedical HIV and STI prevention, this new group of HIV PrEP-using GBM may be more receptive to syphilis PrEP.

Our study is subject to several limitations. One limitation may be the lack of diversity in the risk profile of our sample. Because recruitment was limited to online applications and clinics, and did not include gay venues like bathhouses, there was the potential of missing participants who did not come to clinics or use online applications. These participants may have a different risk profile than our sample and may therefore provide different insights into syphilis-related knowledge and attitudes. Face-to-face interviews may have led to a sampling bias by discouraging potential participants who may not have felt comfortable discussing their views on syphilis directly with the interviewer. Another limitation of this study is its inability to accurately ascertain the level of health literacy among the participants. To more robustly evaluate sexual health literacy among GBM and complement these qualitative findings, we have developed some health literacy questions which will be administered to a large sample of GBM participants in the province as part of the *Sex Now* survey.

Our findings reaffirm earlier qualitative research showing that GBM living with HIV or with a previous syphilis diagnosis possess a high level of perceived knowledge on syphilis,¹⁷ unlike their HIV-negative counterparts without prior syphilis. Furthermore, like Plant *et al*,¹⁷ we found that for most of the participants, 'the benefits of their current sexual behaviour outweigh(ed) the risks of...syphilis infection' and that participants were reluctant to make long-term behaviour changes, such as using condoms, to avoid syphilis infections, despite being knowledgeable and concerned about syphilis. Plant *et al* go on to say, 'Participants' general awareness of the risks of syphilis and unwillingness to change their behaviour to avoid possible future infections suggest that interventions aimed at reducing risk behaviours are unlikely to succeed with these men'. Because of these circumstances, a biomedical intervention such as syphilis PrEP may be an appropriate prevention strategy for the syphilis epidemic. Our findings suggest that syphilis PrEP would be most acceptable to those living with HIV or with a previous syphilis diagnosis, as also reported by Wilson *et al*²³ and Stahlman *et al*, respectively.¹⁹ However, despite some promising early data, syphilis PrEP is not currently a recommended intervention as some fundamental questions around acceptability, efficacy and its potential impact on antimicrobial resistance remain. The results presented herein contribute further feasibility data to the emerging literature on this novel syphilis prevention intervention, by highlighting important patient concerns around syphilis PrEP and identifying possible syphilis PrEP candidates.

Overall, there is a disjuncture between the concerns of public health actors and GBM. While health officials are becoming increasingly alarmed about the syphilis

epidemic, GBM continue to be moderately concerned about syphilis for several reasons, including a lack of self-efficacy or belief in their ability to overcome the epidemic, and their belief that syphilis may not be that serious because it can be treated. Because syphilis appears to be only moderately concerning to many GBM and balanced against the benefits of sexual pleasure, syphilis prevention is not prioritised among GBM. Syphilis PrEP may be a strategy that could empower some individuals to prevent syphilis and give them the agency to know they are doing something to prevent the infection, particularly in the context of a syphilis epidemic.

Author affiliations

¹Clinical Prevention Services, British Columbia Centre for Disease Control, Vancouver, British Columbia, Canada

²CIHR Canadian HIV Trials Network, Vancouver, British Columbia, Canada

³Department of Infectious diseases, University of British Columbia Faculty of Medicine, Vancouver, British Columbia, Canada

⁴School of Population and Public Health, University of British Columbia, Vancouver, British Columbia, Canada

⁵Dalla Lana School of Public Health, University of Toronto, Toronto, Ontario, Canada

Acknowledgements We would like to acknowledge all the men who participated and shared their time, knowledge and experiences with us. We are also thankful to the Community Advisory Board members. We are grateful to the staff at Qmunity for allowing us to use their space for recruitment, and to the nurses at the BC Centre for Disease Control who helped us with participant recruitment. Dr Ronita Nath is supported by a CTN James Kreppner Postdoctoral Fellowship Award. Dr Daniel Grace is supported by a Canada Research Chair in Sexual and Gender Minority Health.

Contributors DG and TG conceived the research question and design for the study. RN, RP and FB collected the data. RN analysed the data, with input from RP, FB and DG. RN drafted the manuscript. DG and TG supervised the data collection and analysis. DG, TG, JW and JC provided revision of the manuscript.

Funding This study was supported by the Vancouver Foundation and BC Centre for Disease Control Foundation for Public Health UNR150412.

Competing interests None declared.

Patient consent for publication Not required.

Ethics approval Research Ethics Board approval was obtained from the University of British Columbia Clinical Research Ethics Board (ethics approval number H16-00423). All participants gave informed written consent prior to taking part in this research.

Provenance and peer review Not commissioned; externally peer reviewed.

Data availability statement All data relevant to the study are included in the article or uploaded as supplementary information.

Open access This is an open access article distributed in accordance with the Creative Commons Attribution Non Commercial (CC BY-NC 4.0) license, which permits others to distribute, remix, adapt, build upon this work non-commercially, and license their derivative works on different terms, provided the original work is properly cited, appropriate credit is given, any changes made indicated, and the use is non-commercial. See: <http://creativecommons.org/licenses/by-nc/4.0/>.

ORCID iD

Ronita Nath <http://orcid.org/0000-0002-3630-7915>

REFERENCES

- Grant RM, Lama JR, Anderson PL, *et al*. Preexposure chemoprophylaxis for HIV prevention in men who have sex with men. *N Engl J Med* 2010;363:2587–99.
- Grant RM, Anderson PL, McMahan V, *et al*. Uptake of pre-exposure prophylaxis, sexual practices, and HIV incidence in men and transgender women who have sex with men: a cohort study. *Lancet Infect Dis* 2014;14:820–9.

- Tsuboi M, Nishijima T, Yashiro S, *et al*. Prognosis of ocular syphilis in patients infected with HIV in the antiretroviral therapy era. *Sex Transm Infect* 2016;92:605–10.
- Cranston RD. Anal cancer prevention: how we are failing men who have sex with men. *Sex Transm Infect* 2008;84:417–9.
- BC Centre for Disease Control. *Sti in British Columbia: annual surveillance report 2016, 2018*. Available: <http://www.bccdc.ca/search?k=sti%20annual%20report> [Accessed 10 Oct 2018].
- Colfax G, Vittinghoff E, Husnik MJ, *et al*. Substance use and sexual risk: a participant- and episode-level analysis among a cohort of men who have sex with men. *Am J Epidemiol* 2004;159:1002–12.
- He H, Wang M, Zaller N, *et al*. Prevalence of syphilis infection and associations with sexual risk behaviours among HIV-positive men who have sex with men in Shanghai, China. *Int J STD AIDS* 2014;25:410–9.
- BC Centre for Disease Control. *Sti in British Columbia: Quarterly report*. 2017, 2018.
- BC Centre for Disease Control. *Hiv in British Columbia: annual surveillance report 2016, 2018*. Available: <http://www.bccdc.ca/search?k=hiv%20annual%20report>
- Schöfer H, Imhof M, Thoma-Greber E, *et al*. Active syphilis in HIV infection: a multicentre retrospective survey. The German AIDS Study Group (GASG). *Sex Transm Infect* 1996;72:176–81.
- Hamze H, Ryan V, Cumming E, *et al*. Hiv seropositivity and early syphilis stage associated with ocular syphilis diagnosis: a case-control study in British Columbia, Canada, 2010–2018. *Clin Infect Dis* 2019. doi:10.1093/cid/ciz794. [Epub ahead of print: 17 Aug 2019].
- Ward H, Martin I, Macdonald N, *et al*. Lymphogranuloma venereum in the United Kingdom. *Clin Infect Dis* 2007;44:26–32.
- Murphy D, Holt M. Responses to syphilis outbreaks among gay and other men who have sex with men: case studies from the United Kingdom and the United States, 2009. Available: <http://www.researchgate.net/publication/237342830> [Accessed 21 Jun 2017].
- Nodin N, Carballo-Diéguez A, Ventuneac AM, *et al*. Knowledge and acceptability of alternative HIV prevention bio-medical products among MSM who bareback. *AIDS Care* 2008;20:106–15.
- Grace D, Chown SA, Jollimore J, *et al*. Hiv-Negative gay men's accounts of using context-dependent sero-adaptive strategies. *Cult Health Sex* 2014;16:316–30.
- Grace D, Chown SA, Kwag M, *et al*. Becoming "Undetectable": Longitudinal Narratives of Gay Men's Sex Lives After a Recent HIV Diagnosis. *AIDS Educ Prev* 2015;27:333–49.
- Plant A, Stahlman S, Javanbakht M, *et al*. Syphilis experiences and risk perceptions among repeatedly infected men who have sex with men. *Perspect Sex Reprod Health* 2015;47:181–6.
- Lambert NL, Imrie J, Fisher MJ, *et al*. Making sense of syphilis: beliefs, behaviours and disclosure among gay men recently diagnosed with infectious syphilis and the implications for prevention. *Sex Health* 2006;3:155–61.
- Stahlman S, Plant A, Javanbakht M, *et al*. Acceptable interventions to reduce syphilis transmission among high-risk men who have sex with men in Los Angeles. *Am J Public Health* 2015;105:e88–94.
- Woo A. B.C. the lone province to adopt HIV "treatment as prevention" strategy, 2013. Available: <http://www.theglobeandmail.com/news/british-columbia/bc-the-lone-province-to-adopt-hiv-treatment-as-prevention-strategy/article14920379/> [Accessed 3 Nov 2018].
- Suecharan C. "We've been waiting for this a long time": B.C. to fund HIV-prevention drug, 2017. Available: <https://www.cbc.ca/news/canada/british-columbia/province-announces-hiv-drug-coverage-1.4467003> [Accessed 8 Nov 2018].
- Bell N. "More than 2,000 British Columbians are now on HIV prevention pill.", 2018. Available: <https://www.dailyxtra.com/more-than-2000-british-columbians-are-now-on-hiv-prevention-pill-87809> [Accessed 21 Mar 2019].
- Wilson DP, Prestage GP, Gray RT, *et al*. Chemoprophylaxis is likely to be acceptable and could mitigate syphilis epidemics among populations of gay men. *Sex Transm Dis* 2011;38:573–9.
- Bolan RK, Beymer MR, Weiss RE, *et al*. Doxycycline prophylaxis to reduce incident syphilis among HIV-infected men who have sex with men who continue to engage in high-risk sex: a randomized, controlled pilot study. *Sex Transm Dis* 2015;42:98–103.
- Robinson OC. Sampling in Interview-Based qualitative research: a theoretical and practical guide. *Qual Res Psychol* 2014;11:25–41.
- Strauss A, Corbin JM. *Grounded theory in practice*. Sage 1997.
- Glaser B, Strauss A. *The Discovery of Grounded Theory: Strategies for Qualitative Research*. New York: Aldine De Gruyter, 1967.
- Strauss A, Corbin J. *Basics of qualitative research: techniques and procedures for developing grounded theory*. thousand oaks. CA: SAGE, 1998.

- 29 Holt M, Bernard D, Race K. Gay men's perceptions of sexually transmissible infections and their experiences of diagnosis: 'part of the way of life' to feeling 'dirty and ashamed'. *Sex Health* 2010;7:411–6.
- 30 Fleming DT, Wasserheit JN. From epidemiological synergy to public health policy and practice: the contribution of other sexually transmitted diseases to sexual transmission of HIV infection. *Sex Transm Infect* 1999;75:3–17.
- 31 Walker C, Preshaw PM, Novak J, *et al.* Long-Term treatment with sub-antimicrobial dose doxycycline has no antibacterial effect on intestinal flora. *J Clin Periodontol* 2005;32:1163–9.
- 32 Nguyen F, Starosta AL, Arenz S, *et al.* Tetracycline antibiotics and resistance mechanisms. *Biol Chem* 2014;395:559–75.
- 33 Webster GF, Graber EM. Antibiotic treatment for acne vulgaris. In: *Seminars in cutaneous medicine and surgery*. , 2008: 27, 183–7.
- 34 Moon SH, Roh HS, Kim YH, *et al.* Antibiotic resistance of microbial strains isolated from Korean acne patients. *J Dermatol* 2012;39:833–7.
- 35 U.S. National Laboratory of Medicine. Clinicaltrials.Gov. clinical trial unique identifier: NCT02844634. Available: <https://clinicaltrials.gov/>
- 36 Community-Based Research Centre for Gay Men's Health. Sex now survey, 2018. Available: <http://cbrc.net/sexnow/about> [Accessed 15 Dec 2018].