BMJ Open Patient safety in inpatient mental health settings: a systematic review

Bethan Thibaut,¹ Lindsay Helen Dewa ,¹ Sonny Christian Ramtale,¹ Danielle D'Lima,² Sheila Adam,¹ Hutan Ashrafian ,¹ Ara Darzi,¹ Stephanie Archer ^{1,3}

To cite: Thibaut B, Dewa LH, Ramtale SC, *et al*. Patient safety in inpatient mental health settings: a systematic review. *BMJ Open* 2019;**9**:e030230. doi:10.1136/bmjopen-2019-030230

► Prepublication history and additional material for this paper are available online. To view these files, please visit the journal online (<http://dx.doi.org/10.1136/bmjopen-2019-030230>).

BT and LHD are joint first authors.

Received 05 March 2019
Revised 11 October 2019
Accepted 20 November 2019

© Author(s) (or their employer(s)) 2019. Re-use permitted under CC BY. Published by BMJ.

¹NIHR Imperial Patient Safety Translational Research Centre, Department of Surgery and Cancer, Imperial College London, London, UK

²Centre for Behaviour Change, Department of Clinical, Educational and Health Psychology, University College London, London, UK

³Department of Public Health and Primary Care, University of Cambridge, Cambridge, UK

Correspondence to

Dr Stephanie Archer;
stephanie.archer@imperial.ac.uk

ABSTRACT

Objectives Patients in inpatient mental health settings face similar risks (eg, medication errors) to those in other areas of healthcare. In addition, some unsafe behaviours associated with serious mental health problems (eg, self-harm), and the measures taken to address these (eg, restraint), may result in further risks to patient safety. The objective of this review is to identify and synthesise the literature on patient safety within inpatient mental health settings using robust systematic methodology.

Design Systematic review and meta-synthesis. Embase, Cumulative Index to Nursing and Allied Health Literature, Health Management Information Consortium, MEDLINE, PsycINFO and Web of Science were systematically searched from 1999 to 2019. Search terms were related to 'mental health', 'patient safety', 'inpatient setting' and 'research'. Study quality was assessed using the Hawker checklist. Data were extracted and grouped based on study focus and outcome. Safety incidents were meta-analysed where possible using a random-effects model.

Results Of the 57 637 article titles and abstracts, 364 met inclusion criteria. Included publications came from 31 countries and included data from over 150 000 participants. Study quality varied and statistical heterogeneity was high. Ten research categories were identified: interpersonal violence, coercive interventions, safety culture, harm to self, safety of the physical environment, medication safety, unauthorised leave, clinical decision making, falls and infection prevention and control.

Conclusions Patient safety in inpatient mental health settings is under-researched in comparison to other non-mental health inpatient settings. Findings demonstrate that inpatient mental health settings pose unique challenges for patient safety, which require investment in research, policy development, and translation into clinical practice.

PROSPERO registration number CRD42016034057.

INTRODUCTION

Patient safety has been defined as the 'avoidance, prevention and amelioration of adverse outcomes or injuries stemming from the process of healthcare'.¹ Those receiving care in inpatient mental health settings face similar risks (eg, medication errors) to patients in other areas of healthcare. In addition, some of the unsafe behaviours associated with serious mental health problems

Strengths and limitations of this study

- This is the first review to examine patient safety within inpatient mental health settings that uses robust systematic methodology.
- The use of a robust patient safety taxonomy provides a comprehensive list of all incident types and resulted in a wide coverage of publications in terms of setting, country and population.
- This review only included peer-reviewed studies with primary data.
- The last systematic literature search was conducted on 27 June 2019, meaning that literature published since this date will not have been included.

(eg, self-harm), and the measures taken to address these (eg, restraint), may result in further risks to patient safety.²⁻⁶ There may also be a tension between maximising patient safety and maintaining patient autonomy. Inpatient services will often include patients who are experiencing high levels of mental distress and are therefore at greatest risk.

While mental health research has focused on components of quality of care, published research lacks focus on the science of patient safety⁷⁻⁹; the stigma and discrimination associated with mental health problems may contribute to this relative neglect.⁷ Only two reviews have examined patient safety in a mental health context and described factors that influence patient safety.^{7,10} These reviews highlighted the complexity of patient safety in mental health, including the importance of wider organisational safety culture. While these reviews offer important insights into this complex topic, only a small number of specific patient safety incidents and concepts were examined. As such, the current breadth and depth of patient safety research in inpatient mental health settings is unknown.

The review presented here is exploratory in nature; building on previous reviews, we aimed to report an overview of the existing research base on patient safety in inpatient

mental health settings. We also aimed to critically reflect on quality and methods used in included studies in the field.¹¹ In addition to our original protocol,¹¹ we aimed to collate, describe and construct the main research categories, allowing for an easily accessible reference index.

SEARCH STRATEGY AND SELECTION CRITERIA

A systematic search was developed in line with the Preferred Reporting Items for Systematic Reviews and Meta-Analyses (PRISMA) guidelines.¹² The protocol for this systematic review has been published elsewhere.¹¹

Six databases were searched: Embase, Cumulative Index to Nursing and Allied Health Literature (CINAHL), Health Management Information Consortium (HMIC), MEDLINE, PsycINFO and Web of Science. The search was originally conducted on 5 April 2016 and then updated on 27 June 2019 using a comprehensive list of search terms (n=343) related to 'mental health' (n=73), 'patient safety' (n=206), 'inpatient setting' (n=13) and 'research' (n=51); see online supplementary files 1 and 2 for full search criteria and terms. The search terms included in the 'patient safety' facet were based on the National Reporting and Learning System (NRLS) taxonomy for England and Wales¹³ to ensure all incident types were identified in the search. A Google Scholar search using the main search terms was also conducted; it was originally anticipated that the first 20 pages of Google scholar would need to be screened against criteria,¹¹ but screening stopped at five pages as no new publications were retrieved. Similarly, we had anticipated hand-searching references of all included papers within the review. However, due to the large number of papers included in the review, only the reference lists of the two existing systematic reviews were searched for additional references.

Five reviewers (BT, CR, LD, DD and SAr) screened all titles against the inclusion and exclusion criteria, with 10% independently screened by a second reviewer (split equally between BT, CR, LD, DD and SAr). Full definitions and descriptions of these criteria can be found in online supplementary file 1 and the protocol published elsewhere.¹¹ Inclusion and exclusion criteria were developed over several iterative rounds among the research team to ensure consistency between reviewers (online supplementary file 1). Any disagreements between reviewers were resolved through discussion and an overall consensus was obtained. Agreement between reviewers was calculated using Cohen's kappa,¹⁴ which is a widely accepted measure of inter-rater reliability.^{15 16} Full-text papers were assessed for inclusion by two reviewers from the research team (BT and one other from CR, LD and SAr); a third reviewer (DD) was consulted if necessary.

Inclusion criteria:

- ▶ Population: mental health inpatients;
- ▶ Intervention/outcomes: patient safety outcomes;
- ▶ Setting: inpatient setting;
- ▶ Comparators: no restriction;

- ▶ General inclusion criteria: empirical peer-reviewed studies with a clear aim or research question, that used primary data and written up in the English language between 1 January 1999 and 27 June 2019 (in line with the publication of the Institute of Medicine's report 'To Err is Human: Building a Safer Health System').¹⁷

Exclusion criteria:

- ▶ Population: centres on physical healthcare patients;
- ▶ Intervention/outcomes: patient safety was not the central aim, research question or outcome
- ▶ Setting: amalgamation of data from inpatient and outpatient settings (where inpatient sample cannot be separated out); primary care, outpatient mental health services, community or social care settings and risk assessment tool reliability/validity checks;
- ▶ Comparators: no restrictions;
- ▶ General exclusion criteria: secondary data, not in English language, protocols, editorials, commentaries/clinical case reviews/'snapshot' studies of a patient group, book chapters, conference abstracts, audits, dissertations, epidemiological studies and reviews.

QUALITY ASSESSMENT

Quality assessment was performed to give an overview of the methodological rigour of included studies and to support readers' interpretation of the literature. Publications were not excluded based on poor quality because the review was purposively exploratory and all-encompassing. Quality was assessed by four reviewers (BT, CR, LD and SAr) using the tool derived by Hawker *et al*,¹⁸ to allow appropriate assessment of the wide variety of studies included in this review. The checklist by Hawker *et al* evaluates nine domains: 1) abstract/title; 2) introduction and aims; 3) method and data; 4) sampling; 5) data analysis; 6) ethics and bias; 7) results; 8) transferability and generalisability and 9) implications and usefulness. For each study, the nine domains were assessed using one of four quality categories: very poor (10 points), poor (20 points), fair (30 points) and good (40 points). The scores for each study were then summed and divided by nine to get an average score.

DATA EXTRACTION

Data were extracted by five reviewers (BT, CR, LD, DD and SAr) using a standardised form that included study design information, participant characteristics, intervention description and patient safety outcomes. Extractions were compared within the research team to ensure reliability. Only published data were extracted; study authors were contacted only for confirmation or information clarity. If the contact attempt was unsuccessful, the article was assessed in its current form.

DATA SYNTHESIS

Studies were grouped into research categories through consensus. First, four research team members (BT, CR,

LD and SAr) individually re-read the included full-text publications and assigned each one based on the main topic area (eg, aggression). Second, each assigned topic area was checked by another team member to ensure reliability. Third, topic areas were grouped into broader research categories (eg, interpersonal violence) that best described the patient safety focus for easier navigation of the literature. Finally, these categories and the related subcategories (initially called topic areas) from the previous stage were finalised after group discussion and consensus was reached. This was to ensure mutual exclusivity and appropriate definition (table 1 and online supplementary file 3). Where data allowed, meta-analysis was performed applying a random-effects model, specifically calculating pooled prevalence considering both between-study and within-study variances that contributed to study weighting. Pooled values and 95% CIs were computed and represented on forest plots. Statistical heterogeneity was determined by the I^2 statistic, where <30% is low, 30%–60% is moderate and >60% is high. Analyses were performed using Stata V.15 (StataCorp, College Station, Texas, USA).

PATIENT AND PUBLIC INVOLVEMENT

Patients and the public were not involved in this study.

RESULTS

The search resulted in 79 672 records (figure 1) and reduced to 57 637 after de-duplication. Titles and abstracts were screened and excluded if they did not satisfy inclusion criteria (BT, CR, LD, DD and SAr). Ten per cent were then screened (n=5763) by a second independent reviewer (split equally between BT, CR, LD, DD and SAr), in line with guidance on improving decision making by including more than one person in this process¹⁹; good agreement was found between pairs of reviewers ($\kappa=0.72$). A total of 4758 publications were subjected to full-text review (BT, CR, LD and SAr). Two reviewers independently screened the full-text articles against inclusion criteria (BT, CR, LD and SAr). The third reviewer (DD) was consulted 59 times. Substantial agreement was reached ($\kappa=0.64$). From the full-text review, 4394 publications were excluded. Three hundred and sixty-four publications met the inclusion criteria and data were extracted (online supplementary file 4).

Study characteristics

Table 1 provides an overview of the study characteristics. The publications spanned 5 continents and 31 countries. The three countries contributing the greatest number of studies were the UK (n=102), the USA (n=55) and Australia (n=32). The included studies collected data from over 150 000 participants. Studies included staff (n=165; 45%), patients (n=120; 33%) and a mixture of staff, patients and/or carers (n=77; 21%). Only one study focused solely on patient family members (<1%). Most

studies were quantitative in nature (n=192; 53%), just over a third were qualitative (n=133; 37%) and a small proportion used mixed methodology (n=39; 11%). Studies were conducted in a variety of settings comprising: psychiatric inpatient wards/facilities (n=266; 73%), forensic inpatient facilities (n=50; 14%), long-term care/nursing homes (n=25; 7%), mixed inpatient settings (n=20; 5%), a learning disability unit (n=1; <1%), a health board (n=1; <1%) and a specialised research unit (n=1; <1%). More information about the study designs used is included in online supplementary file 4.

Quality assessment

Most research was assessed as 'fair' quality (n=251; 69%), 86 (24%) papers were assessed as 'good' quality and 26 (7%) were assessed as 'poor' quality. None was assessed as 'very poor' quality. Studies rated as 'poor' mainly did not discuss ethical considerations, potential biases or give sample or setting characteristics. For example, they did not consider recruitment strategies, sample demographics or provide detailed information on the research setting. All 'good' studies provided setting and sampling information to allow for replicability. In addition, 'good' studies provided detail on data analysis justification, more thorough literature reviews to place the study in context and had clear research aims/objectives. Online supplementary file 5 includes a table showing the breakdown of the quality domain scores for each paper.

Synthesis

Ten research categories were identified: interpersonal violence, coercive interventions, safety culture, harm to self, safety of the physical environment, medication safety, unauthorised leave, clinical decision making, falls and infection prevention and control. Within these categories 46 subcategories were identified (table 1).

Interpersonal violence

Interpersonal violence was the largest category (n=116; 32%). Studies were primarily concerned with the prevalence, management and prevention of violent and aggressive behaviours (n=75). The pooled prevalence for physical violence was 43.2% (95% CI 0.37 to 0.49) with high heterogeneity (I^2 100.0%) in 20 studies^{20–39} (online supplementary file 6). The pooled prevalence for verbal aggression was 57.4% (95% CI 0.34 to 0.81) with a high heterogeneity (I^2 100.0%) in 10 studies^{22–24 26 29 33–36 40} (online supplementary file 6).

One study examined the characteristics of aggressive incidents by ward type,⁴¹ and two studies identified correlates of violence.^{42 43} One study explored how patients described their aggressive behaviours.⁴⁴ Twenty-four studies evaluated intervention effectiveness (eg, staff training and medication use) to reduce violent and aggressive behaviours, with most finding significant improvements,^{45–65} two reporting negative outcomes^{66 67} and one reporting mixed findings.⁶⁸ The general management of violent and aggressive behaviours was explored

Table 1 Overview of study characteristic identified within each category

Category	Subcategory	Category definition	Number of studies	Countries	Number of studies using staff participants	Number of studies using patient participants	Total number of participants	Settings (number of studies conducted in each setting)
Interpersonal violence	Aggression	Behaviours or events that are considered hostile	115	UK-31	52	39	20 066 (excl. missing data)	Psychiatric inpatient wards/facilities- 73
	Violence	with the intent to cause harm, including violence, aggression and conflicts. This also encompasses sudden emergency incidents that require management.		USA-20	22 mixed	22 mixed		Forensic inpatient facilities-22
	Challenging behaviour			Australia-9		1 family member of patients		Long-term care/nursing homes-13
	Violence and aggression			Canada-7		1 N/A		Specialised research unit-1
	Critical incidents			The -6				Mixed-6
	Conflict			Sweden-7				
	Sexual Assault			Taiwan-4				
	Agitation			South Africa-2				
				Switzerland-2				
				India-3				
				Italy-2				
				Turkey-3				
				Europe-2				
				New Zealand-1				
				South Korea-1				
				Finland-3				
				Greece-1				
				Spain-1				
				Hong Kong-1				
				Israel-2				
				Nigeria-1				
				Norway-2				
				Denmark-1				
				Japan-1				
				Germany-1				
				Slovakia-1				
Coercive interventions	Restraint	Techniques for managing patient behaviour that are applied without consent, for the safety of the patient and others. These include seclusion, restraint and containment.	99	UK-31	36	34	59 732 (excl. missing data)	Psychiatric inpatient wards/facilities- 74
	Seclusion			Finland-7	29 mixed	29 mixed		Forensic inpatient facilities-13
	Attitudes to coercion			USA-8				Long-term care/nursing homes-3
	Containment			The Netherlands-5				Mixed-8
	Process of coercion			Australia-5				Health board-1
	Alternative interventions			Canada-7				
	Shielding			Norway-4				
	Conflict			Germany-2				
	Personal factors			Sweden-3				
				Japan-2				
				Mixed-5				
				New Zealand-2				
				Europe-1				
				China-1				
				Switzerland-3				
				South Korea-1				
				India-2				
				Brazil-1				
				Denmark-1				

Continued

Table 1 Continued

Category	Subcategory	Category definition	Number of studies	Countries	Number of studies using staff participants	Number of studies using patient participants	Total number of participants	Settings (number of studies conducted in each setting)
Safety culture	Process	The organisational attitudes, beliefs and values concerning safety. This encompasses the policies and procedures within the healthcare organisation in relation to safety.	49	UK-12	33	3	59 420	Psychiatric inpatient wards/facilities- 36
	Culture			Australia-10	13 mixed	13 mixed	(excl. missing data)	Forensic inpatient facilities-8
Harm to self	Policy	This encompasses the policies and procedures within the healthcare organisation in relation to safety.	36	USA-5	16	17	3631 (excl. missing data)	Long-term care/nursing homes-1
	Building therapeutic relationships			Sweden-4				Mixed-4
	Patient/family engagement			Finland-4				
				Canada-2				
				Ireland-2				
				The Netherlands-1				
				Greece-1				
				Italy-1				
				Germany-1				
				Belgium-2				
Safety of the physical environment	Self-harm	The ways in which the healthcare system attempts to prevent, mitigate or manage deliberate behaviours displayed by patients that are intended to cause harm or death to themselves.	21	Taiwan-1	8 mixed	7	3140 (excl. missing data)	Psychiatric inpatient wards/facilities- 17
	Self-neglect			Europe-1				Forensic inpatient facilities-1
				Iran-2				Long-term care/nursing homes-3
				USA-11				
				UK-8				
				Ireland-3				
				Norway-4				
				The Netherlands-2				
				Sweden-2				
				Taiwan-2				
Continued		The factors related to the physical environment of the healthcare setting that could impact on safety. This includes ligature points, staffing, security (door locking) and patient distribution.	6	Australia-1	8 mixed	7	3140 (excl. missing data)	Psychiatric inpatient wards/facilities- 17
				Japan-2				Forensic inpatient facilities-1
				Belgium-1				Long-term care/nursing homes-3
				UK-6				
				The Netherlands-3				
				USA-4				
				Australia-4				
				Germany-2				
				Mixed-1				
				Sweden-1				

Continued

Table 1 Continued

Category	Subcategory	Category definition	Number of studies	Countries	Number of studies using staff participants	Number of studies using patient participants	Total number of participants	Settings (number of studies conducted in each setting)
Medication safety	Adverse events	Mistakes made at any stage of the medication use process, from preparation, to administration, to management	17	UK-7	9	7	2396 (excl. missing data)	Psychiatric inpatient wards/facilities- 13
	Medication management	Medication dispensing		Turkey-1	1 mixed	1 mixed		Forensic inpatient facilities-2
	Adherence	Recording. This includes adverse drug events		Spain-2				Long-term care/nursing homes-1
	Substance use	(or injuries that are the result of a drug-related intervention) and issues surrounding drug/alcohol use.		The Netherlands-1				Mixed-1
Unauthorised leave	Absconding	The act of a patient leaving the healthcare setting without the knowledge or consent of staff/carers. This can be either with (absconding) or without intent (wandering) on the part of the patient.	11	UK-4	3	7	978 (excl. missing data)	Psychiatric inpatient wards/facilities- 10
	Wandering			Australia-3	1 mixed	1 mixed		Long-term care/nursing homes-1
Clinical decision making	Incident management	Incorrect diagnoses, risk assessments and other decision making processes of healthcare staff that impact on the safety of a patient.	9	USA-3	6	3	529	Psychiatric inpatient wards/facilities- 8
	Risk assessment			UK-3				Forensic inpatient facilities-1
Falls	Falls	Falling events that lead to the unintentional harm of an individual. This includes trips and injuries such as fractures.	6	Canada-1			180 (excl. missing data)	Psychiatric inpatient wards/facilities- 5
	Injuries			Greece-1				Long-term care/nursing homes-1
Infection prevention and control	Infection prevention and control	Preventing harm caused by infection to patients and health workers.	1	Taiwan-1	1	0	13	Psychiatric inpatient wards/facilities- 1

Figure 1 Flow chart of studies.

in 15 studies.^{22 25 29 30 69–79} Two studies explored the ways in which treatment can affect violence incidence.^{31 80}

Twenty-seven studies explored violent and aggressive incident experiences in staff,^{81–96} patients,^{97–99} mixed groups^{100–106} and patient family members.²⁷ Five studies explored the risk factors associated with verbal and physical aggression.^{35 37 107–109} Three studies explored mental health nurses' perspectives on the response to violent situations in high secure environments: one on the psychological impact of physical assault on staff,¹¹⁰ one on making violence risk assessments in imminent violent situations¹¹¹ and one on the decline of incident reports.¹¹² One study explored the link between aggressive behaviour and levels of burnout in staff¹¹³ and one study looked at the role of social support for staff following a violent incident.³²

Ten studies^{114–123} examined challenging behaviour and techniques, such as de-escalation and communication strategies, which could be used to manage this; seven studies found techniques that were effective.^{114–120} A further four studies investigated conflict behaviour management techniques employed by staff^{124–126} and patients¹²⁷; techniques used in the two intervention studies were effective in reducing conflict.^{126 127} Staff and patient attitudes towards critical incidents were the focus

of four qualitative studies^{128–131}; a further three studies focused on maintaining the psychological safety of patients who had experienced physical or sexual assault during an inpatient stay¹³² and outside of healthcare.^{133 134} Finally, one study explored an acupuncture intervention to reduce agitation, which was found to be effective.¹³⁵

Coercive interventions

Coercive interventions were the focus of 98 papers (27%). Most studies (n=42) reported on restraint and seclusion techniques. The pooled prevalence for coercive interventions was 47.8% (95% CI 0.38 to 0.57) with high heterogeneity (I^2 100.0%) in 12 studies^{136–147} (online supplementary file 6).

Studies explored staff,^{148–157} patient^{147 158–165} and mixed groups^{166–173} views and experiences of seclusion and restraint. Nine studies focused on the processes surrounding seclusion and restraint.^{136 137 174–180} A further 16 studies evaluated interventions to reduce seclusion and restraint, with 13 finding significant decreases in rates of use,^{146 181–192} one reporting an increase¹⁹³ and one reporting increased levels of knowledge about the topic area.¹⁹⁴ Four studies examined prevalence, trends and preventative factors^{138 195–197}; one found that 45% of patients were subjected to restraint,¹³⁸ and another found that restraint and seclusion declined over time.¹⁹⁷ One study explored the context in which seclusion and restraint had taken place.¹⁹⁸ Two studies found preventative factors of mechanical restraint to be staff education and increased patient involvement.^{195 196} The training of staff in techniques for seclusion and restraint were explored in two studies^{199 200} and one study examined adverse events resulting from restraint and seclusion.²⁰¹ Other studies explored staff and patient views of containment measures,^{202–205} Maori views of initiatives to reduce/prevent seclusion,²⁰⁶ the process of shielding (segregation under staff supervision),²⁰⁷ conflict management²⁰⁸ and alternative interventions.²⁰⁹

Thirty-two studies focused on coercion; one study examined prevalence of coercive measures¹⁴¹ and one study explored how the experience of staff might contribute to the use of restrictive practices.²¹⁰ The attitudes of staff,^{142 144 211–222} patients^{145 223–226} and mixed groups^{143 168 227–230} towards coercion were explored in 25 studies, and 5 studies examined the process of coercive interventions^{139 140 231 232} and rules of engagement in caring for aggressive patients.²³³

Safety culture

Safety culture included studies on processes, culture and policy across 49 papers (13%). Eighteen studies concerned safety-related organisational processes. Eleven of these investigated processes of treatment or care that healthcare staff undertake; processes included limit-setting and clothing restrictions,^{234–240} risk assessment^{241–243} and nursing handover.²⁴⁴ Two investigated errors and reporting^{245 246} and a further two studies explored staff and patient perceptions of safety when

involved in treatment processes.^{247 248} Two studies focused on change implementation.^{249 250} One study focused on the role of training.²⁵¹ Safety culture was featured in 18 publications relating to the management of serious incidents,^{252–254} stress and burnout,^{255–257} staff²⁵⁸ and patient perspectives of safety^{259–263} and communication²⁶⁴; there were also three papers that explored safety culture more generally.^{265–267} A further two evaluated the TeamSTEPPS (Team Strategies and Tools to Enhance Performance and Patient Safety) programme^{268 269} and both found significant clinical benefits in reducing seclusion and improving team functioning. One paper looked at the barriers and facilitators to implementing a Safewards intervention.²⁷⁰ With regard to policy, eight studies concerned safety policies related to: observation,^{271 272} risk assessment,^{273 274} treatment,²⁷⁵ safeguarding,²⁷⁶ security²⁷⁷ and ergonomic improvement.²⁷⁸ Two papers focused on the role of patient and family engagement in safety,^{279 280} and two papers focused on how to build better therapeutic relationships to improve patient safety.^{281 282}

Harm to self

Three subcategories centred on harmful behaviours: self-harm, suicidal behaviour and self-neglect (n=36; 10%). Half of the studies (n=18) focused on self-harm. One paper explored the prevalence of self-harm.²⁸³ Two studies explored risk factors for self-harm which included use of psychotropic medication.^{284 285} Eight papers explored staff attitudes and experiences of managing self-harm,^{286–293} and three explored patient experiences.^{294–296} Three intervention studies focused on training,²⁹⁷ therapy²⁹⁸ and observation²⁹⁹; all reported a reduction in self-harm behaviours and a further intervention focusing on training for staff resulted in positive attitude towards self-harm patients, greater closeness and improved self-efficacy.³⁰⁰ Of the 17 papers that centred on suicidal behaviours, five studies investigated the observance of risk factors^{301–305} and three intervention studies found significant reductions in suicide-related behaviours and cognitions.^{306–308} An additional eight papers explored staff,^{309–312} patient^{313 314} and both staff and patient^{315 316} views and attitudes towards suicidal behaviour. One study looked at the acceptability of an intervention to reduce suicide.³¹⁷ Finally, one study explored types of self-neglect behaviours in patients with dementia, including functional difficulties, serious hygiene problems and safety risks.³¹⁸

Safety of the physical environment

The safety of the physical environment category included 21 papers (6%). Seven studies investigated security measures (eg, door locking).^{319–325} Five studies investigated the effects of the physical environmental design on the safety of treatment settings.^{326–330} Three studies focused on safety during transitions of care,^{331–333} with most based in dementia care settings. Three studies examined how the location of patients within the hospital setting can impact on safety, focusing on topics such as:

privacy, female-only wards and the use of segregated or combined wards/units.^{334–336} The remaining three studies concerned staffing levels^{337 338} and ligature points.³³⁹

Medication safety

The medication safety category included 17 publications (5%). Five studies focused on adverse events, and examined: antipsychotics side effects,³⁴⁰ how best to manage the effect of psychotropics on long QT segments,³⁴¹ antidepressants³⁴² and medication error reporting.^{343 344} Three studies investigated errors occurring in broader medication management processes^{345–347} and a further five studies focused on medication administration specifically.^{348–352} The only intervention study aiming to reduce these errors found that a new medication dispensing system did not have any significant impact on patient safety.³⁵³ Two studies explored staff perceptions of illicit substance use.^{354 355} One further study described the development of a medication adherence intervention for patients who are prescribed mood-stabilising medication for bipolar disorder.³⁵⁶

Unauthorised leave

Unauthorised leave included 11 publications (3%). Three explored the patient experience of absconding, specifically relating to patient perspectives of treatment and involuntary commitment.^{357–359} One study explored staff perspectives of absconding management techniques,³⁶⁰ and two studies evaluated interventions to reduce absconding rates; both were found to be effective.^{361 362} Two studies focused on wandering behaviour in women with dementia, linking wandering to physical environment factors, such as light, sound, crowding³⁶³ and falls.³⁶⁴ The pooled prevalence of wandering behaviour was 50.2% (95% CI 0.49 to 0.52) with high heterogeneity (I^2 78.0%) in two studies^{363 364} (online supplementary file 6). The final three studies examined the consequences^{365 366} and security measures surrounding absconding.³⁶⁷

Clinical decision making

Clinical decision making accounted for 2% of the included publications (n=9). These publications covered the development of clinical judgements and decisions relating to incident management, risk assessment and diagnosis. Two studies explored the cultural differences considered by clinicians in the diagnosis of African-American patients.^{368 369} Clinical decisions on whether to engage in seclusion and/or restraint were explored in five studies^{370–374} and two studies explored the variation in assessment and prediction of violence between staff and settings.^{375 376}

Falls

Publications on falls formed the second smallest category within the review (n=6; 1%). Studies in this category focused on fall prevalence, falls experienced by older psychiatric inpatients with dementia and prevention/harm reduction techniques. A recurring risk factor for falling was found to be medication use.^{377–379} Two fall

prevention intervention studies did not identify significant benefits,^{380 381} and one study explored barriers and facilitators to such interventions.³⁸²

Infection prevention and control

One paper (<1%) focused on staff experiences of infection prevention and control in psychiatric clinical settings.³⁸³

DISCUSSION

Main findings

This is the first review to examine patient safety within inpatient mental health settings that uses robust systematic methodology. As a result, we have identified ten research categories: interpersonal violence, coercive interventions, safety culture, harm to self, safety of the physical environment, medication safety, unauthorised leave, clinical decision making, falls and infection prevention and control. In addition, we have been able to include a meta-analysis of incidence and prevalence of aggression (verbal and physical), coercive intervention and wandering behaviour as well as providing an easily accessible reference index of literature in the inpatient mental health and patient safety domain. Previous reviews on this topic had focused on collating the literature on a restricted number (n=8) of predefined patient safety incidents (eg, violence and aggression),⁷ or the concept of patient safety in inpatient mental health setting more broadly (eg, organisation management).¹⁰ As such, the findings presented here offer a contemporary view of the breadth and depth of patient safety research in inpatient mental health settings.

We were concerned to see that only 364 papers were identified as a result of our comprehensive search. Although this can be seen as a large number of publications for a systematic review, it is a relatively small number to cover the care of a wide range of patients in a variety of inpatient mental health settings over a 20-year period (around 18 papers per year across all countries). While important work not meeting our inclusion criteria (eg, quality improvement initiatives and studies using secondary analysis of data) may have focused on patient safety in mental health, the lack of prospective peer-reviewed publications adds to the ongoing discussion surrounding the disparity in research focusing on patient safety in physical and mental healthcare.³⁸⁴ In addition, there was a paucity of high-quality research in the area; just over two-thirds of the studies were considered to be 'fair', and only nine studies included in the meta-analysis were deemed 'good'. 'Poor' studies most frequently did not have clear research aims and objectives, study details were missing (eg, sample(s) and setting(s) used) and they failed to discuss issues related to ethical and researcher bias. Some qualitative studies explored both staff and patients' perspectives, an important aspect of research, particularly when safety in this context is a relatively new area of knowledge. However, there was limited intervention research,

particularly randomised controlled trials (RCTs). In the RCTs that were identified, sample sizes were mostly small.

The findings from the review also challenged our expectations in terms of breadth and depth of research. For example, we expected to find many publications on the prevention of suicide within inpatient settings due to the severity of harm. However, only one study that met inclusion criteria discussed suicide in relation to ligature points.³³⁹ A scoping review also found only this one study, suggesting a consistency of approach.³⁸⁵ This indicates that while the prevention of suicide is a well-established aspect of patient safety, it is now reviewed routinely, using pre-existing and secondary data, rather than through empirical research.

We also found little research focusing on the concepts required for system level reform,³⁸⁶ which was disappointing seeing as some improvements have been made in physical healthcare.³⁸⁷ For example, in line with research in the physical health domain,^{388 389} we were hoping to find several studies exploring how patient and family engagement in care can promote patient safety.³⁹⁰ However, only two studies identified in our review had patient/family engagement as their primary focus.^{279 280} Similarly, we were expecting to identify literature investigating the lack of integration between physical and mental healthcare and the impact it has on patient safety.³⁹¹ However, the need to prevent and manage co-existing physical ill health was not identified in the review. This is surprising as patients with serious mental illness are twice as likely to die prematurely and much more likely to develop long-term conditions or become disabled, as those without serious mental illness.³⁹² This patient group is also vulnerable to asphyxiation during restraint and rapid tranquilisation.³⁹³

Research on medication safety in inpatient mental health settings was also limited in this review. This was unexpected considering two-thirds of patients with mental health problems are prescribed medication and are therefore potentially at risk of experiencing a medication safety incident. Research pertaining to falls was also limited, contrasting with patient safety research within the physical health domain that includes a focus on slips, trips and falls.³⁹⁴

Strengths and limitations

We used a robust patient safety taxonomy to provide a comprehensive list of all incident types. This resulted in a wide coverage of publications in terms of setting, country and population. We systematically searched, screened, extracted and appraised data. As a result, our systematic review draws together all relevant literature concerning patient safety within inpatient mental health settings, simultaneously operating as an index resource for clinicians and researchers.

There were several limitations. We used the definition of patient safety given by Vincent¹ to guide this review. While this is more nuanced than the original Institute

of Medicine definition of safety ‘freedom from accidental injury’³⁹⁵ and is widely accepted within the patient safety movement, it may be that a more suitable definition reflects the specific challenges within the inpatient mental health setting.³⁹⁶ This review only included peer-reviewed studies with primary data. Therefore, literature using secondary data such as pre-existing datasets and data from internal audits was excluded as it did not fulfil the criteria of being a prospective research study with clear research aims.³⁹⁷ For example, data examined by the National Confidential Inquiry into Suicide and Homicide by People with Mental Illness is collected retrospectively from various sites across the country and would have been excluded from this review.³⁹⁸ Moreover, non-peer-reviewed quality improvement reports have also been excluded. The decision was made to only include peer-reviewed studies with primary data due to (i) the large number of potential publications in this area, (ii) the need to define the scope and focus of the review and (iii) the need for specificity as well as sensitivity. The investigation of patient safety in mental health inpatient settings using secondary data or in non-peer-reviewed formats is an avenue for additional systematic reviews.

The last systematic literature search was conducted on 27 June 2019, meaning that literature published since this date will not have been included. In order to further build on the review published here, a *living* systematic review (an ongoing updated summary of high-quality research)³⁹⁹ would continue to identify relevant literature in this area. In terms of the meta-analysis, there was expected statistical and methodological variability in studies, particularly for physical and verbal aggression. It is possible that this was due to the inclusion of different definitions of aggression, time periods and type of inpatient setting. In relation to the agreement between reviewers (including the use of recommended piloting of inclusion and exclusion criteria within the screening stage),⁴⁰⁰ inter-rater reliability calculations only achieved substantial agreement ($\kappa=0.61-0.80$) at both the title and full-text screening stages. Although higher kappas have been reported in other systematic reviews, a substantial agreement is classified as more than acceptable.⁴⁰¹

While the research spanned five continents, the UK, the USA and Australia contributed over 50% of the included studies, leading to a potential cultural bias in the body of research identified within the review. We recommend that, where possible, future systematic reviews incorporate manuscripts in languages other than English to establish greater insight into the global literature on patient safety in inpatient mental health settings, with a view to limiting any cultural bias. Similarly, while the removal of publications denoting non-inpatient setting restricted the conclusions to the inpatient setting, issues pertaining to this environment are likely to be different to that of community, primary or social care settings. Additionally, studies were excluded before 1999 to coincide with the release of the Institute of Medicine’s report ‘To Err is Human: Building a Safer

Health System’³⁹⁵; this may have narrowed the review scope as the historical context was minimised.

Clinical implications and future research

This review informs academics, clinicians and service providers about the evidence base in the patient safety field within inpatient mental health settings. The findings allow researchers and clinicians to be directed to literature relevant to a given patient safety topic area, a useful starting point when developing practice guidelines.⁴⁰² Similarly, the findings may influence clinical practice, with those implementing interventions or designing service changes being able to easily access the current scientific understanding.

Future research should be informed by patient safety science more broadly and focus on filling the knowledge gaps highlighted in this review, that is, studies that explore (i) systems level improvement, (ii) patient and carer engagement in safety, (iii) suicide prevention across different countries, (iv) the nature of medication safety in inpatient mental health settings and (v) the prevalence and impact of staff to patient violence. These findings support our previous expert consensus study where academic and service user experts agreed that patient-driven research studies were needed.⁴⁰³ The limited rigorous research surrounding patient safety within inpatient mental health settings necessitates future studies to: (i) include large inpatient samples relevant to the research design, (ii) perform appropriate intervention testing and (iii) examine safety from different perspectives. It should also focus on high-quality reporting of research, paying particular attention to the area of ethics, sampling and setting characteristics.

CONCLUSION

This is the first systematic review to comprehensively examine research on patient safety within inpatient mental health settings. It has drawn together the existing literature and shed light on the gaps in knowledge. Inpatient mental health settings may demonstrate unique patient safety challenges and more research is needed to achieve parity with physical health. Addressing this through a strong body of evidence, informed by patient safety science more broadly, will mean that mental health-care policy makers are in a better position to address safety issues, and implement robust and evidence-based interventions to improve care.

Twitter Lindsay Helen Dewa @dewalindsay

Acknowledgements The authors would like to thank the librarians at St Mary’s Library, Imperial College London for their support with the study.

Contributors BT, LD, SCR, SAr and DD contributed to the design, data searches, data extraction, synthesis and writing of the report. HA contributed to the design, data extraction, meta-analysis and writing of the report. SAd and AD contributed to the design and synthesis, as well as writing and critically reviewing the report.

Funding This work is supported by the National Institute for Health Research (NIHR) Imperial Patient Safety Translation Research Centre. Infrastructure support was provided by the NIHR Imperial Biomedical Research Centre.

Competing interests None declared.

Patient consent for publication Not required.

Provenance and peer review Not commissioned; externally peer reviewed.

Data availability statement All data relevant to the study are included in the article or uploaded as supplementary information.

Open access This is an open access article distributed in accordance with the Creative Commons Attribution 4.0 Unported (CC BY 4.0) license, which permits others to copy, redistribute, remix, transform and build upon this work for any purpose, provided the original work is properly cited, a link to the licence is given, and indication of whether changes were made. See: <https://creativecommons.org/licenses/by/4.0/>.

ORCID iDs

Lindsay Helen Dewa <http://orcid.org/0000-0001-8359-8834>

Hutan Ashrafian <http://orcid.org/0000-0003-1668-0672>

Stephanie Archer <http://orcid.org/0000-0003-1349-7178>

REFERENCES

- Vincent C. *Patient safety*. Edinburgh: Churchill Livingstone, 2006.
- Brickell TA, McLean C. Emerging issues and challenges for improving patient safety in mental health: a qualitative analysis of expert perspectives. *J Patient Saf* 2011;7:39–44.
- Powell J, Geddes J, Deeks J, et al. Suicide in psychiatric hospital in-patients: risk factors and their predictive power. *Br J Psychiatry* 2000;176:266–72.
- Roy A, Draper R. Suicide among psychiatric hospital in-patients. *Psychol Med* 1995;25:199–202.
- Gray NS, Hill C, McGleish A, et al. Prediction of violence and self-harm in mentally disordered offenders: a prospective study of the efficacy of HCR-20, PCL-R, and psychiatric symptomatology. *J Consult Clin Psychol* 2003;71:443–51.
- Goh SE, Salmons PH, Whittington RM. Hospital suicides: are there preventable factors? profile of the psychiatric hospital suicide. *Br J Psychiatry* 1989;154:247–9.
- Brickell TA, Nicholls TL, Procyshyn RM. *Patient safety in mental health*. Canadian Patient Safety Institute, 2000.
- Vincent GM. Psychopathy and violence risk assessment in youth. *Child Adolesc Psychiatr Clin N Am* 2006;15:407–28.
- Taylor TL, Killaspy H, Wright C, et al. A systematic review of the International published literature relating to quality of institutional care for people with longer term mental health problems. *BMC Psychiatry* 2009;9:55.
- Kanerva A, Lammintakanen J, Kivinen T. Patient safety in psychiatric inpatient care: a literature review. *J Psychiatr Ment Health Nurs* 2013;20:541–8.
- D'Lima D, Archer S, Thibaut BI, et al. A systematic review of patient safety in mental health: a protocol based on the inpatient setting. *Syst Rev* 2016;5:203.
- Moher D, Shamseer L, Clarke M, et al. Preferred reporting items for systematic review and meta-analysis protocols (PRISMA-P) 2015 statement. *Syst Rev* 2015;4:1.
- NHS Improvement. Learning from patient safety incidents, 2014. Available: <http://www.nrls.npsa.nhs.uk/patient-safety-data/>
- Cohen J. A coefficient of agreement for nominal scales. *Educ Psychol Meas* 1960;20:37–46.
- McHugh ML. Interrater reliability: the kappa statistic. *Biochemia Medica* 2012;22:276–82.
- Belur J, Tompson L, Thornton A, et al. Interrater reliability in systematic review methodology: exploring variation in coder decision-making. *Soc Med Res* 2018.
- Donaldson MS, Corrigan JM, Kohn LT. *To err is human: building a safer health system*. 6. National Academies Press, 2000.
- Hawker S, Payne S, Kerr C, et al. Appraising the evidence: reviewing disparate data systematically. *Qual Health Res* 2002;12:1284–99.
- Edwards P, Clarke M, DiGiuseppi C, et al. Identification of randomized controlled trials in systematic reviews: accuracy and reliability of screening records. *Stat Med* 2002;21:1635–40.
- Amoo G, Fatoye FO. Aggressive behaviour and mental illness: a study of in-patients at Aro neuropsychiatric Hospital, Abeokuta. *Niger J Clin Pract* 2010;13:351–5.
- Boström A-M, Squires JE, Mitchell A, et al. Workplace aggression experienced by frontline staff in dementia care. *J Clin Nurs* 2012;21:1453–65.
- Chen W-C, Hwu H-G, Wang J-D. Hospital staff responses to workplace violence in a psychiatric hospital in Taiwan. *Int J Occup Environ Health* 2009;15:173–9.
- Daffern M, Mayer M, Martin T. Staff gender ratio and aggression in a forensic psychiatric hospital. *Int J Ment Health Nurs* 2006;15:93–9.
- Ilkiw-Lavalle O, Grenyer BFS. Differences between patient and staff perceptions of aggression in mental health units. *Psychiatr Serv* 2003;54:389–93.
- Jacob P, Seshadri S, Girimaji SC, et al. Clinical characteristics of aggression in children and adolescents admitted to a tertiary care centre. *Asian J Psychiatry* 2013;6:556–9.
- Lehmann LS, McCormick RA, Kizer KW. A survey of assaultive behavior in veterans health administration facilities. *Psychiatr Serv* 1999;50:384–9.
- Raveendranathan D, Chandra PS, Chaturvedi SK. Violence among psychiatric inpatients: a victim's perspective. *East Asian Arch Psychiatry* 2012;22:141–5.
- Ryan EP, Hart VS, Messick DL, et al. A prospective study of assault against staff by youths in a state psychiatric hospital. *Psychiatr Serv* 2004;55:665–70.
- Schwartz TL, Park TL. Assaults by patients on psychiatric residents: a survey and training recommendations. *Psychiatr Serv* 1999;50:381–3.
- Sukhodolsky DG, Cardona L, Martin A. Characterizing aggressive and noncompliant behaviors in a children's psychiatric inpatient setting. *Child Psychiatry Hum Dev* 2005;36:177–93.
- Wystanski M. Assaultive behaviour in psychiatrically hospitalized elderly: a response to psychosocial stimulation and changes in pharmacotherapy. *Int J Geriatr Psychiatry* 2000;15:582–5.
- Kelly EL, Fenwick KM, Brekke JS, et al. Sources of social support after patient assault as related to staff well-being. *J Interpers Violence* 2017;8:088626051773877–79.
- Niu S-F, Kuo S-F, Tsai H-T, et al. Prevalence of workplace violent episodes experienced by nurses in acute psychiatric settings. *PLoS One* 2019;14:e0211183.
- Danivas V, Lepping P, Punitharani S, et al. Observational study of aggressive behaviour and coercion on an Indian acute ward. *Asian J Psychiatry* 2016;22:150–6.
- Podubinski T, Lee S, Hollander Y, et al. Patient characteristics associated with aggression in mental health units. *Psychiatry Res* 2017;250:141–5.
- van den Bogaard KJHM, Nijman HLI, Palmstierna T, et al. Characteristics of aggressive behavior in people with mild to borderline intellectual disability and co-occurring psychopathology. *J Ment Health Res Intellect Disabil* 2018;11:124–42.
- Pekurinen VM, Välimäki M, Virtanen M, et al. Organizational justice and collaboration among nurses as correlates of violent assaults by patients in psychiatric care. *Psychiatr Serv* 2017;68:490–6.
- Kelly EL, Fenwick K, Brekke JS, et al. Well-Being and safety among inpatient psychiatric staff: the impact of conflict, assault, and stress reactivity. *Adm Policy Ment Health* 2016;43:703–16.
- Bilici R, Sercan M, Izci F. Levels of the staff's exposure to violence at locked psychiatric clinics: a comparison by occupational groups. *Issues Ment Health Nurs* 2016;37:501–6.
- Daffern M, Ogloff J, Howells K. Aggression in an Australian forensic psychiatric hospital. *J British Forensic Pract* 2003;5:18–28.
- Sato M, Noda T, Sugiyama N, et al. Characteristics of aggression among psychiatric inpatients by ward type in Japan: Using the Staff Observation Aggression Scale - Revised (SOAS-R). *Int J Ment Health Nurs* 2017;26:602–11.
- Ben-Zeev D, Scherer EA, Brian RM, et al. Use of multimodal technology to identify digital correlates of violence among inpatients with serious mental illness: a pilot study. *Psychiatr Serv* 2017;68:1088–92.
- Bowers L, Allan T, Simpson A, et al. Identifying key factors associated with aggression on acute inpatient psychiatric wards. *Issues Ment Health Nurs* 2009;30:260–71.
- Selenius H, Leppänen Östman S, Strand S. Self-Harm as a risk factor for inpatient aggression among women admitted to forensic psychiatric care. *Nord J Psychiatry* 2016;70:554–60.
- van de Sande R, Nijman HLI, Noorthoorn EO, et al. Aggression and seclusion on acute psychiatric wards: effect of short-term risk assessment. *Br J Psychiatry* 2011;199:473–8.
- Park JS, Lee K. Modification of severe violent and aggressive behavior among psychiatric inpatients through the use of a short-term Token economy. *J Korean Acad Nurs* 2012;42:1062–9.
- Needham I, Abderhalden C, Meer R, et al. The effectiveness of two interventions in the management of patient violence in acute mental inpatient settings: report on a pilot study. *J Psychiatr Ment Health Nurs* 2004;11:595–601.

- 48 Murphy CJ, Siv AM. A one year study of mode deactivation therapy: adolescent residential patients with conduct and personality disorders. *Int J Behav Cogn Ther* 2007;3:327–41.
- 49 McLaughlin S, Bonner G, Mboche C, *et al*. A pilot study to test an intervention for dealing with verbal aggression. *Br J Nurs* 2010;19:489–94.
- 50 Killick S, Allen D. Training staff in an adolescent inpatient psychiatric unit in positive approaches to managing aggressive and harmful behaviour: does it improve confidence and knowledge? *Child Care Prac* 2005;11:323–39.
- 51 Giles GM, Wager J, Fong L, *et al*. Twenty-month effectiveness of a non-aversive, long-term, low-cost programme for persons with persisting neurobehavioural disability. *Brain Injury* 2005;19:753–64.
- 52 Daffern M, Howells K, Hamilton L, *et al*. The impact of structured risk assessments followed by management recommendations on aggression in patients with personality disorder. *J Forens Psychiatry Psychol* 2009;20:661–79.
- 53 C-Y Yip V, Gudjonsson GH, Perkins D, *et al*. A non-randomised controlled trial of the R&R2MHP cognitive skills program in high risk male offenders with severe mental illness. *BMC Psychiatry* 2013;13:267.
- 54 Lanza M, Ridenour M, Hendricks S, *et al*. The violence prevention community meeting: a multi-site study. *Arch Psychiatr Nurs* 2016;30:382–6.
- 55 Lanza ML, Rierdan J, Forester L, *et al*. Reducing violence against nurses: the violence prevention community meeting. *Issues Ment Health Nurs* 2009;30:745–50.
- 56 Calabro K, MacKey TA, Williams S. Evaluation of training designed to prevent and manage patient violence. *Issues Ment Health Nurs* 2002;23:3–15.
- 57 Björkdahl A, Hansebo G, Palmstierna T. The influence of staff training on the violence prevention and management climate in psychiatric inpatient units. *J Psychiatr Ment Health Nurs* 2013;20:396–404.
- 58 Arguvanli S, Çet al. Effect of aggression management training program on knowledge and attitudes of nurses working at psychiatric clinics/Agresyon yönetimi eğitim programının psikiyatri kliniklerinde çalışan hemşirelerin bilgi ve tutumlarına etkisi. *Anadolu Psikiyatri Dergisi* 2015;16.
- 59 Caspi N, Modai I, Barak P, *et al*. Pindolol augmentation in aggressive schizophrenic patients: a double-blind crossover randomized study. *Int Clin Psychopharmacol* 2001;16:111–5.
- 60 Davies BE, Lowe K, Morgan S, *et al*. An evaluation of the effectiveness of positive behavioural support within a medium secure mental health forensic service. *J Forens Psychiatry Psychol* 2019;30:38–52.
- 61 Isaak V, Vashdi D, Bar-Noy D, *et al*. Enhancing the safety climate and reducing violence against staff in closed hospital wards. *Workplace Health Saf* 2017;65:409–16.
- 62 Price O, Baker J, Bee P, *et al*. Patient perspectives on barriers and enablers to the use and effectiveness of de-escalation techniques for the management of violence and aggression in mental health settings. *J Adv Nurs* 2018;74:614–25.
- 63 Hvidhjelm J, Sestoft D, Skovgaard LT, *et al*. Aggression in psychiatric wards: effect of the use of a structured risk assessment. *Issues Ment Health Nurs* 2016;37:960–7.
- 64 Olsson H, Schön U-K. Reducing violence in forensic care - how does it resemble the domains of a recovery-oriented care? *J Ment Health* 2016;25:506–11.
- 65 Mistler LA, Ben-Zeev D, Carpenter-Song E, *et al*. Mobile mindfulness intervention on an acute psychiatric unit: feasibility and acceptability study. *JMIR Mental Health* 2017;4.
- 66 Needham I, Abderhalden C, Halfens RJG, *et al*. The effect of a training course in aggression management on mental health nurses' perceptions of aggression: a cluster randomised controlled trial. *Int J Nurs Stud* 2005;42:649–55.
- 67 Sjöström N, Eder DN, Malm U, *et al*. Violence and its prediction at a psychiatric hospital. *Eur Psychiatry* 2001;16:459–65.
- 68 Sival RC, Albronda T, Haffmans PMJ, *et al*. Is aggressive behaviour influenced by the use of a behaviour rating scale in patients in a psychogeriatric nursing home? *Int J Geriatr Psychiatry* 2000;15:108–11.
- 69 Skovdahl K, Kihlgren AL, Kihlgren M. Dementia and aggressiveness: video recorded morning care from different care units. *J Clin Nurs* 2003;12:888–98.
- 70 Stone T, McMillan M, Hazelton M, *et al*. Wounding words: swearing and verbal aggression in an inpatient setting. *Perspect Psychiatr Care* 2011;47:194–203.
- 71 Sutton D, Wilson M, Van Kessel K, *et al*. Optimizing arousal to manage aggression: a pilot study of sensory modulation. *Int J Ment Health Nurs* 2013;22:500–11.
- 72 Lipscomb JA, London M, Chen YM, *et al*. Safety climate and workplace violence prevention in state-run residential addiction treatment centers. *Work* 2012;42:47–56.
- 73 Wright S, Sayer J, Parr A-M, *et al*. Breakaway and physical restraint techniques in acute psychiatric nursing: results from a national survey of training and practice. *J Forens Psychiatry Psychol* 2005;16:380–98.
- 74 Berg J, Kaltiala-Heino R, Välimäki M. Management of aggressive behaviour among adolescents in forensic units: a four-country perspective. *J Psychiatr Ment Health Nurs* 2011;18:776–85.
- 75 Biancosino B, Delmonte S, Grassi L, *et al*. Violent behavior in acute psychiatric inpatient facilities: a national survey in Italy. *J Nerv Ment Dis* 2009;197:772–82.
- 76 Caspi E. Aggressive behaviors between residents with dementia in an assisted living residence. *Dementia* 2015;14:528–46.
- 77 Chaplin R, McGeorge M, Hinchliffe G, *et al*. Aggression on psychiatric inpatient units for older adults and adults of working age. *Int J Geriatr Psychiatry* 2008;23:874–6.
- 78 Delaney J, Cleary M, Jordan R, *et al*. An exploratory investigation into the nursing management of aggression in acute psychiatric settings. *J Psychiatr Ment Health Nurs* 2001;8:77–84.
- 79 Meaden A, Hacker D, Spencer K. Acute aggression risk: an early warning signs methodology. *The Journal of Forensic Practice* 2013;15:21–31.
- 80 Umut G, Altun Zeren Öztürk, Danişmant BS, *et al*. Relationship between treatment adherence, insight and violence among schizophrenia inpatients in a training Hospital sample. *Düşünen Adam* 2012;25:212–20.
- 81 Bahareethan M, Shah A. Aggressive behaviour, staff attitude and staff perception of patients on two continuing care psychogeriatric wards. *Aging Ment Health* 2000;4:66–71.
- 82 Berg J, Kaltiala-Heino R, Löyttyneemi V, *et al*. Staff's perception of adolescent aggressive behaviour in four European forensic units: a qualitative interview study. *Nord J Psychiatry* 2013;67:124–31.
- 83 Stevenson KN, Jack SM, O'Mara L, *et al*. Registered nurses' experiences of patient violence on acute care psychiatric inpatient units: an interpretive descriptive study. *BMC Nurs* 2015;14:35.
- 84 Spokes K, Bond K, Lowe T, *et al*. HOVIS - The Hertfordshire/Oxfordshire Violent Incident Study. *J Psychiatr Ment Health Nurs* 2002;9:199–209.
- 85 Zuzelo PR, Curran SS, Zeserman MA. Registered nurses' and behavior health associates' responses to violent inpatient interactions on behavioral health units. *J Am Psychiatr Nurses Assoc* 2012;18:112–26.
- 86 Camuccio CA, CHAMBERS M, VÄLIMÄKI M, *et al*. Managing distressed and disturbed patients: the thoughts and feelings experienced by Italian nurses. *J Psychiatr Ment Health Nurs* 2012;19:807–15.
- 87 McCann TV, Baird J, Muir-Cochrane E. Factors influencing clinicians' attitudes about aggression in Australian acute old age psychiatry inpatient units: a cross sectional survey design. *Issues Ment Health Nurs* 2014;35:542–50.
- 88 Tema TR, Poggenpoel M, Myburgh CPH. Experiences of psychiatric nurses exposed to hostility from patients in a forensic ward. *J Nurs Manag* 2011;19:915–24.
- 89 Nijman H, Bowers L, Oud N, *et al*. Psychiatric nurses' experiences with inpatient aggression. *Aggress Behav* 2005;31:217–27.
- 90 Chen Wen-Ching, Wang Jung-Der, Lew-Ting Chih-Yin, *et al*. Workplace violence on workers caring for Long-term institutionalized schizophrenic patients in Taiwan. *J Occup Health* 2007;49:311–6.
- 91 Cutcliffe JR. Qualified nurses' lived experience of violence perpetrated by individuals suffering from enduring mental health problems: a hermeneutic study. *Int J Nurs Stud* 1999;36:105–16.
- 92 Evans RE, Petter S. Identifying mitigating and challenging beliefs in dealing with threatening patients: an analysis of experiences of clinicians working in a psychiatric intensive care unit. *Journal of Psychiatric Intensive Care* 2012;8:113–9.
- 93 Lantta T, Daffern M, Kontio R, *et al*. Implementing the dynamic appraisal of situational aggression in mental health units. *Clinical Nurse Specialist* 2015;29:230–43.
- 94 Tomášková M, Bóriková I, Lepiešová M, *et al*. NURSES' EXPERIENCE AND ATTITUDES TOWARDS INPATIENT AGGRESSION ON PSYCHIATRIC WARDS. *CEJNM* 2016;7:462–9.
- 95 Hylén U, Engström I, Engström K, *et al*. Providing good care in the shadow of violence – an interview study with nursing staff and ward managers in psychiatric inpatient care in Sweden. *Issues Ment Health Nurs* 2019;40:148–57.
- 96 Lantta T, Anttila M, Kontio R, *et al*. Violent events, ward climate and ideas for violence prevention among nurses in psychiatric wards: a focus group study. *Int J Ment Health Syst* 2016;10.

- 97 Meehan T, McIntosh W, Bergen H. Aggressive behaviour in the high-secure forensic setting: the perceptions of patients. *J Psychiatr Ment Health Nurs* 2006;13:19–25.
- 98 Olsson H, Audulv Åsa, Strand S, et al. Reducing or increasing violence in forensic care: a qualitative study of inpatient experiences. *Arch Psychiatr Nurs* 2015;29:393–400.
- 99 Van Wijk E, Traut A, Julie H. Environmental and nursing-staff factors contributing to aggressive and violent behaviour of patients in mental health facilities. *Curationis* 2014;37:01–8.
- 100 Daffern M. Assessing the functions of aggression in psychiatric inpatients. *Behav Anal Today* 2007;8:43–51.
- 101 Duxbury J, Whittington R. Causes and management of patient aggression and violence: staff and patient perspectives. *J Adv Nurs* 2005;50:469–78.
- 102 Janicki N. Prosecuting inpatient violence: perceptions of staff, patients and others in a women's enhanced medium secure service. *The British Journal of Forensic Practice* 2009;11:27–38.
- 103 Lawn S, Pols R. Nicotine withdrawal: pathway to aggression and assault in the locked psychiatric ward? *Australasian Psychiatry* 2003;11:199–203.
- 104 Nolan KA, Shope CB, Citrome L, et al. Staff and patient views of the reasons for aggressive incidents: a prospective, incident-based study. *Psychiatr Q* 2009;80:167–72.
- 105 Dickens G, Piccirillo M, Alderman N. Causes and management of aggression and violence in a forensic mental health service: perspectives of nurses and patients. *Int J Ment Health Nurs* 2013;22:532–44.
- 106 Wright KM, Duxbury JA, Baker A, et al. A qualitative study into the attitudes of patients and staff towards violence and aggression in a high security hospital. *J Psychiatr Ment Health Nurs* 2014;21:184–8.
- 107 Lamanna D, Ninkovic D, Vijayarajnam V, et al. Aggression in psychiatric hospitalizations: a qualitative study of patient and provider perspectives. *J Ment Health* 2016;25:536–42.
- 108 Paschali M, Kamp D, Reichmann C, et al. A systematic evaluation of impulsive-aggressive behavior in psychogeriatric inpatients using the staff observation aggression scale-revision (SOAS-R). *Int Psychogeriatr* 2018;30:61–8.
- 109 Almvik R, Rasmussen K, Woods P. Challenging behaviour in the elderly—monitoring violent incidents. *Int J Geriatr Psychiatry* 2006;21:368–74.
- 110 Reininghaus U, Craig T, Gournay K, et al. The High Secure Psychiatric Hospitals' Nursing Staff Stress Survey 3: Identifying stress resistance resources in the stress process of physical assault. *Pers Individ Dif* 2007;42:397–408.
- 111 Trenoweth S. Perceiving risk in dangerous situations: risks of violence among mental health inpatients. *J Adv Nurs* 2003;42:278–87.
- 112 DE Niet GJ, Hutschemaekers GJM, Lendemeijer BHHG. Is the reducing effect of the staff observation aggression scale owing to a learning effect? an explorative study. *J Psychiatr Ment Health Nurs* 2005;12:687–94.
- 113 de Looft P, Nijman H, Didden R, et al. Burnout symptoms in forensic psychiatric nurses and their associations with personality, emotional intelligence and client aggression: a cross-sectional study. *J Psychiatr Ment Health Nurs* 2018;25:506–16.
- 114 Bharwani G, Parikh PJ, Lawhorne LW, et al. Individualized behavior management program for Alzheimer's/dementia residents using behavior-based ergonomic therapies. *Am J Alzheimers Dis Other Dement* 2012;27:188–95.
- 115 Carlson GA, Potegal M, Margulies D, et al. Liquid risperidone in the treatment of rages in psychiatrically hospitalized children with possible bipolar disorder. *Bipolar Disord* 2010;12:205–12.
- 116 Chan S, Fung MY, Tong CW, et al. The clinical effectiveness of a multisensory therapy on clients with developmental disability. *Res Dev Disabil* 2005;26:131–42.
- 117 DeYoung S, Just G, Harrison R. Decreasing aggressive, agitated, or disruptive behavior: participation in a behavior management unit. *J Gerontol Nurs* 2002;28:22–31.
- 118 Higuera Aet al. Effects of a humor-centered activity on disruptive behavior in patients in a general Hospital psychiatric ward. *International Journal of Clinical and Health Psychology* 2006;6:53–64.
- 119 Speziale J, Black E, Coatsworth-Puspoky R, et al. Moving forward: Evaluating a curriculum for managing responsive behaviors in a geriatric psychiatry inpatient population. *Gerontologist* 2009;49: :570–6. 10.1093/geront/gnp069
- 120 Zwijsen SA, Smalbrugge M, Eefsting JA, et al. Coming to grips with challenging behavior: a cluster randomized controlled trial on the effects of a multidisciplinary care program for challenging behavior in dementia. *J Am Med Dir Assoc* 2014;15:531.e1–531.e10. : p. e1-531.
- 121 Bennett R, Ramakrishna V, Maganty D. Management of disturbed behaviour in a psychiatric intensive care unit: views of staff on options for intervention. *Journal of Psychiatric Intensive Care* 2011;7:85–9.
- 122 Foley KL, Sudha S, Sloane PD, et al. Staff perceptions of successful management of severe behavioral problems in dementia special care units. *Dementia* 2003;2:105–24.
- 123 Hallett N, Dickens GL. De-escalation: a survey of clinical staff in a secure mental health inpatient service. *Int J Ment Health Nurs* 2015;24:324–33.
- 124 Bowers L. Association between staff factors and levels of conflict and containment on acute psychiatric wards in England. *Psychiatric Services* 2009;60:231–9.
- 125 Lowe T, Wellman N, Taylor R. Limit-setting and decision-making in the management of aggression. *J Adv Nurs* 2003;41:154–61.
- 126 Ireland JL, Sebalo I, McNeill K, et al. Impacting on factors promoting intra-group aggression in secure psychiatric settings. *Heliyon* 2019;5.
- 127 Quirk A, Lelliott P, Seale C. Risk management by patients on psychiatric wards in London: an ethnographic study. *Health Risk Soc* 2005;7:85–91.
- 128 Ireland CA, Halpin L, Sullivan C. Critical incidents in a forensic psychiatric population: an exploratory study of motivational factors. *J Forens Psychiatry Psychol* 2014;25:714–32.
- 129 Jeffs L, Rose D, Macrae C, et al. What near misses tell us about risk and safety in mental health care. *J Psychiatr Ment Health Nurs* 2012;19:430–7.
- 130 Koukia E, Mangoulia P, Stathopoulos T, et al. Greek mental health nurses' practices and attitudes in the management of acute cases. *Issues Ment Health Nurs* 2013;34:192–7.
- 131 Terkelsen TB, Larsen IB. Fear, danger and aggression in a Norwegian locked psychiatric ward: dialogue and ethics of care as contributions to combating difficult situations. *Nursing Ethics* 2016;23:308–17.
- 132 Cole M, Baldwin D, Thomas P. Sexual assault on wards: staff actions and reactions. *Int J Psychiatry Clin Pract* 2003;7:239–42.
- 133 Phillips L. Reflections on the education and training of mental health staff who work with women who have been sexually abused in childhood. *J Psychiatr Ment Health Nurs* 2011;18:696–705.
- 134 Gallop R, Engels S, DiNunzio R, et al. Abused women's concerns about safety and the therapeutic environment during psychiatric hospitalization. *Can J Nurs Res* 1999;31:53–70.
- 135 Yang M-H, Wu S-C, Lin J-G, et al. The efficacy of acupressure for decreasing agitated behaviour in dementia: a pilot study. *J Clin Nurs* 2007;16:308–15.
- 136 Cormac I, Russell I, Ferriter M. Review of seclusion policies in high secure hospitals and medium secure units in England, Scotland and Wales. *J Psychiatr Ment Health Nurs* 2005;12:380–2.
- 137 Georgieva I, Mulder CL, Noorthoorn E. Reducing seclusion through involuntary medication: a randomized clinical trial. *Psychiatry Res* 2013;205:48–53.
- 138 Kirkevold Øyvind, Engedal K. Prevalence of patients subjected to constraint in Norwegian nursing homes. *Scand J Caring Sci* 2004;18:281–6.
- 139 Raboch J, Kalisová L, Nawka A, et al. Use of coercive measures during involuntary hospitalization: findings from ten European countries. *Psychiatr Serv* 2010;61:1012–7.
- 140 Steinert T, Martin V, Baur M, et al. Diagnosis-Related frequency of compulsory measures in 10 German psychiatric hospitals and correlates with Hospital characteristics. *Soc Psychiatry Psychiatr Epidemiol* 2007;42:140–5.
- 141 Gowda GS, Lepping P, Noorthoorn EO, et al. Restraint prevalence and perceived coercion among psychiatric inpatients from South India: a prospective study. *Asian J Psychiatry* 2018;36:10–16.
- 142 Gowda GS, Lepping P, Ray S, et al. Clinician attitude and perspective on the use of coercive measures in clinical practice from tertiary care mental health establishment - A cross-sectional study. *Indian J Psychiatry* 2019;61:151–5.
- 143 Hotzy F, Jaeger M, Buehler E, et al. Attitudinal variance among patients, next of kin and health care professionals towards the use of containment measures in three psychiatric hospitals in Switzerland. *BMC Psychiatry* 2019;19:128.
- 144 Vedana KGG, da Silva DM, Ventura CAA, et al. Physical and mechanical restraint in psychiatric units: perceptions and experiences of nursing staff. *Arch Psychiatr Nurs* 2018;32:367–72.
- 145 Krieger E, Moritz S, Weil R, et al. Patients' attitudes towards and acceptance of coercion in psychiatry. *Psychiatry Res* 2018;260:478–85.
- 146 Reynolds EK, Grados MA, Pragowski N, et al. Use of modified positive behavioral interventions and supports in a psychiatric

- inpatient unit for high-risk youths. *Psychiatric Services* 2016;67:570–3.
- 147 Brady NS, Spittal MJ, Brophy LM, *et al.* Patients' Experiences of Restrictive Interventions in Australia: Findings From the 2010 Australian Survey of Psychosis. *Psychiatric Services* 2017;68:966–9.
 - 148 Bigwood S, Crowe M. 'It's part of the job, but it spoils the job': A phenomenological study of physical restraint. *Int J Ment Health Nurs* 2008;17:215–22.
 - 149 Lee S, GRAY R, GOURNAY K, *et al.* Views of nursing staff on the use of physical restraint. *J Psychiatr Ment Health Nurs* 2003;10:425–30.
 - 150 Perkins E, Prosser H, Riley D, *et al.* Physical restraint in a therapeutic setting; a necessary evil? *Int J Law Psychiatry* 2012;35:43–9.
 - 151 Sequeira H, Halstead S. The psychological effects on nursing staff of administering physical restraint in a secure psychiatric hospital: 'When I go home, it's then that I think about it'. *The British Journal of Forensic Practice* 2004;6:3–15.
 - 152 Exworthy T, Mohan D, Hindley N, *et al.* Seclusion: punitive or protective? *The Journal of Forensic Psychiatry* 2001;12:423–33.
 - 153 Kuosmanen L, Makkonen P, Lehtila H, *et al.* Seclusion experienced by mental health professionals. *J Psychiatr Ment Health Nurs* 2015;22:333–6.
 - 154 Muir-Cochrane EC, Baird J, McCann T. Nurses' experiences of restraint and seclusion use in short-stay acute old age psychiatry inpatient units: a qualitative study. *J Psychiatr Ment Health Nurs* 2015;22:109–15.
 - 155 Duxbury J, Thomson G, Scholes A, *et al.* Staff experiences and understandings of the restrain yourself initiative to minimize the use of physical restraint on mental health wards. *Int J Ment Health Nurs* 2019;28:845–56.
 - 156 Nielsen LD, Gildberg FA, Bech P, *et al.* Forensic mental health clinician's experiences with and assessment of alliance regarding the patient's readiness to be released from mechanical restraint. *Int J Ment Health Nurs* 2018;27:116–25.
 - 157 Kontio R, Välimäki M, Putkonen H, *et al.* Patient restrictions: are there ethical alternatives to seclusion and restraint? *Nurs Ethics* 2010;17:65–76.
 - 158 Chien W-T, Chan CWH, Lam L-W, *et al.* Psychiatric inpatients' perceptions of positive and negative aspects of physical restraint. *Patient Educ Couns* 2005;59:80–6.
 - 159 Knowles SF, Hearne J, Smith I. *Physical restraint and the therapeutic relationship.* *J Forens Psychiatry Psychol* 2015;26:461–75.
 - 160 Ezeobele IE, Malecha AT, Mock A, *et al.* Patients' lived seclusion experience in acute psychiatric hospital in the United States: a qualitative study. *J Psychiatr Ment Health Nurs* 2014;21:303–12.
 - 161 Faschingbauer KM, Peden-McAlpine C, Tempel W. Use of Seclusion: finding the voice of the patient to influence practice. *J Psychosoc Nurs Ment Health Serv* 2013;51:32–8.
 - 162 Holmes D, Kennedy SL, Perron A. The mentally ill and social exclusion: a critical examination of the use of seclusion from the patient's perspective. *Issues Ment Health Nurs* 2004;25:559–78.
 - 163 Bergk J, Einsiedler B, Flammer E, *et al.* A randomized controlled comparison of seclusion and mechanical restraint in inpatient settings. *Psychiatr Serv* 2011;62:1310–7.
 - 164 Kontio R, Joffe G, Putkonen H, *et al.* *Seclusion and restraint in psychiatry: patients' experiences and practical suggestions on how to improve practices and use alternatives.* *Perspect Psychiatr Care* 2012;48:16–24.
 - 165 Larue C, Dumais A, Boyer R, *et al.* The experience of seclusion and restraint in psychiatric settings: perspectives of patients. *Issues Ment Health Nurs* 2013;34:317–24.
 - 166 Bonner G, Lowe T, Rawcliffe D, *et al.* Trauma for all: a pilot study of the subjective experience of physical restraint for mental health inpatients and staff in the UK. *J Psychiatr Ment Health Nurs* 2002;9:465–73.
 - 167 Fish R, Hatton C. Gendered experiences of physical restraint on locked wards for women. *Disabil Soc* 2017;32:790–809.
 - 168 Wilson C, Rouse L, Rae S, *et al.* Mental health inpatients' and staff members' suggestions for reducing physical restraint: A qualitative study. *J Psychiatr Ment Health Nurs* 2018;25:188–200.
 - 169 Fish R. 'Behind This Wall' – Experiences of Seclusion on Locked Wards for Women. *Scandinavian Journal of Disability Research* 2018;20:139–51.
 - 170 Gowda GS, Kumar CN, Ray S, *et al.* Caregivers' attitude and perspective on coercion and restraint practices on psychiatric inpatients from South India. *J Neurosci Rural Pract* 2019;10:261–6.
 - 171 Larue C, Goulet M-H, Prevost M-J, *et al.* Identification and analysis of factors contributing to the reduction in Seclusion and restraint for a population with intellectual disability. *Journal of Applied Research in Intellectual Disabilities* 2018;31:e212–22.
 - 172 Wilson C, Rouse L, Rae S, *et al.* Is restraint a 'necessary evil' in mental health care? Mental health inpatients' and staff members' experience of physical restraint. *Int J Ment Health Nurs* 2017;26:500–12.
 - 173 Holmes D, Murray SJ, Knack N. Experiencing Seclusion in a forensic psychiatric setting: a phenomenological study. *J Forensic Nurs* 2015;11:200–13.
 - 174 Bonner G, Wellman N. *Postincident review of aggression and violence in mental health settings.* *J Psychosoc Nurs Ment Health Serv* 2010;48:35–40.
 - 175 Bowers L, Van Der Merwe M, Paterson B, *et al.* Manual restraint and shows of force: the City-128 study. *Int J Ment Health Nurs* 2012;21:30–40.
 - 176 Tompsett CJ, Domoff S, Boxer P. Prediction of restraints among youth in a psychiatric Hospital: application of translational action research. *J Clin Psychol* 2011;67:368–82.
 - 177 Boumans CE, Egger JIM, Souren PM, *et al.* Nurses' decision on seclusion: patient characteristics, contextual factors and reflexivity in teams. *J Psychiatr Ment Health Nurs* 2012;19:264–70.
 - 178 Mason T, Whitehead E. Some specific problems of secluding female patients. *Med Sci Law* 2001;41:315–24.
 - 179 Ryan R, Happell B. Learning from experience: using action research to discover consumer needs in post-seclusion Debriefing. *Int J Ment Health Nurs* 2009;18:100–7.
 - 180 Whitecross F, Seeary A, Lee S. Measuring the impacts of seclusion on psychiatry inpatients and the effectiveness of a pilot single-session post-seclusion counselling intervention. *Int J Ment Health Nurs* 2013;22:512–21.
 - 181 Bleijlevens MHC, Gulpers MJM, Capezuti E, *et al.* Process evaluation of a multicomponent intervention program (EXBELT) to reduce belt restraints in nursing homes. *J Am Med Dir Assoc* 2013;14:599–604.
 - 182 Pelfolk TJ-E, Gustafson Y, Bucht G, *et al.* Effects of a restraint minimization program on staff knowledge, attitudes, and practice: a cluster randomized trial. *J Am Geriatr Soc* 2010;58:62–9.
 - 183 Schreiner GM, Crafton CG, Sevin JA. Decreasing the use of mechanical restraints and locked seclusion. *Adm Policy Ment Health* 2004;31:449–63.
 - 184 Ching H, Daffern M, Martin T, *et al.* Reducing the use of seclusion in a forensic psychiatric Hospital: assessing the impact on aggression, therapeutic climate and staff confidence. *J Forens Psychiatry Psychol* 2010;21:737–60.
 - 185 Long CG, West R, Afford M, *et al.* Reducing the use of seclusion in a secure service for women. *Journal of Psychiatric Intensive Care* 2015;11:84–94.
 - 186 Smith S, Jones J. Use of a sensory room on an intensive care unit. *J Psychosoc Nurs Ment Health Serv* 2014;52:22–30.
 - 187 Espinosa L, Harris B, Frank J, *et al.* Milieu improvement in psychiatry using evidence-based practices: the long and winding road of culture change. *Arch Psychiatr Nurs* 2015;29:202–7.
 - 188 Kontio R, LAHTI M, PITKÄNEN A, *et al.* Impact of eLearning course on nurses' professional competence in seclusion and restraint practices: a randomized controlled study (ISRCTN32869544). *J Psychiatr Ment Health Nurs* 2011;18:813–21.
 - 189 Goulet M-H, Larue C, Lemieux AJ. A pilot study of "post-seclusion and/or restraint review" intervention with patients and staff in a mental health setting. *Perspect Psychiatr Care* 2018;54:212–20.
 - 190 Blair EW, Woolley S, Szarek BL, *et al.* Reduction of Seclusion and restraint in an inpatient psychiatric setting: a pilot study. *Psychiatr Q* 2017;88:1–7.
 - 191 Newman J, Paun O, Fogg L. Effects of a staff training intervention on Seclusion rates on an adult inpatient psychiatric unit. *J Psychosoc Nurs Ment Health Serv* 2018;56:23–30.
 - 192 Elzubeir K, Dye S. Can amount and duration of seclusion be reduced in psychiatric intensive care units by agreeing smart goals with patients? *Journal of Psychiatric Intensive Care* 2017;13:109–16.
 - 193 Huizing AR, Hamers JPH, Gulpers MJM, *et al.* Short-Term effects of an educational intervention on physical restraint use: a cluster randomized trial. *BMC Geriatr* 2006;6:17.
 - 194 Abdel-Hussein NH, Mohamed SH. Effectiveness of an Educational Program on Nurses' Knowledge toward Seclusion and Seclusion for inpatients at Psychiatric Teaching Hospitals. *Indian J Public Health Res Dev* 2018;9:1175–80.
 - 195 Bak J, Zoffmann V, Sestoft DM, *et al.* Mechanical restraint in psychiatry: preventive factors in theory and practice. A Danish-Norwegian association study. *Perspect Psychiatr Care* 2014;50:155–66.

- 196 Bak J, Zoffmann V, Sestoft DM, *et al.* Comparing the effect of non-medical mechanical restraint preventive factors between psychiatric units in Denmark and Norway. *Nord J Psychiatry* 2015;69:1715–25.
- 197 Keski-Valkama A, Sailas E, Eronen M, *et al.* A 15-year national follow-up: legislation is not enough to reduce the use of seclusion and restraint. *Soc Psychiatry Psychiatr Epidemiol* 2007;42:747–52.
- 198 Goulet M-H, Larue C, Dumais A. Evaluation of seclusion and restraint reduction programs in mental health: a systematic review. *Aggress Violent Behav* 2017;34:139–46.
- 199 Lee SW, Sayer J, Parr A-M, *et al.* Soo, physical restraint training for nurses in English and Welsh psychiatric intensive care and regional secure units. *J Ment Health* 2001;10:151–62.
- 200 Kontio R, Välimäki M, Putkonen H, *et al.* Nurses' and physicians' educational needs in seclusion and restraint practices. *Perspect Psychiatr Care* 2009;45:198–207.
- 201 Hatta K, Shibata N, Ota T, *et al.* Association between physical restraint and drug-induced liver injury. *Neuropsychobiology* 2007;56:180–4.
- 202 Bowers L, Alexander J, Simpson A, *et al.* Student psychiatric nurses' approval of containment measures: Relationship to perception of aggression and attitudes to personality disorder. *Int J Nurs Stud* 2007;44:349–56.
- 203 Braham LG, Heasley JF, Akiens S. An evaluation of night confinement in a high secure Hospital. *Mental Health Rev J* 2013;18:21–31.
- 204 Chu S, McNeill K, Wright KM, *et al.* The impact of a night confinement policy on patients in a UK high secure inpatient mental health service. *The Jnl of Forensic Practice* 2015;17:21–30.
- 205 Hottinen A, VÄLIMÄKI M, SAILAS E, *et al.* Attitudes towards different containment measures: a questionnaire survey in Finnish adolescent psychiatry. *J Psychiatr Ment Health Nurs* 2012;19:521–7.
- 206 Wharewera-Mika JP, Cooper EP, Wiki NRN, *et al.* Strategies to reduce the use of seclusion with tāngata whai i te ora (Māori mental health service users). *Int J Ment Health Nurs* 2016;25:258–65.
- 207 Haugom EW, Granerud A. Shielding in mental health hospitals: description and assessment by staff. *SAGE Open* 2016;6.
- 208 Papadopoulos C, Bowers L, Quirk A, *et al.* Events preceding changes in conflict and containment rates on acute psychiatric wards. *Psychiatr Serv* 2012;63:40–7.
- 209 Ejneborn Looi G-M, Engström Åsa, Sävénstedt S. A self-destructive care: self-reports of people who experienced coercive measures and their suggestions for alternatives. *Issues Ment Health Nurs* 2015;36:96–103.
- 210 Barr L, Wynaden D, Heslop K. Promoting positive and safe care in forensic mental health inpatient settings: evaluating critical factors that assist nurses to reduce the use of restrictive practices. *Int J Ment Health Nurs* 2019;28:888.
- 211 Bak J, Aggernæs H. Coercion within Danish psychiatry compared with 10 other European countries. *Nord J Psychiatry* 2012;66:297–302.
- 212 Jaeger M, Ketteler D, Rabenschlag F, *et al.* Informal coercion in acute inpatient setting—Knowledge and attitudes held by mental health professionals. *Psychiatry Res* 2014;220:1007–11.
- 213 Seo MK, Kim SH, Rhee M. Coercion in psychiatric care: can paternalism justify coercion? *Int J Soc Psychiatry* 2013;59:217–23.
- 214 Lovell A, Smith D, Johnson P. A qualitative investigation into nurses' perceptions of factors influencing staff injuries sustained during physical interventions employed in response to service user violence within one secure learning disability service. *J Clin Nurs* 2015;24:1926–35.
- 215 Tateno M, Sugiura K, Uehara K, *et al.* Attitude of young psychiatrists toward coercive measures in psychiatry: a case vignette study in Japan. *Int J Ment Health Syst* 2009;3:20.
- 216 Elmer T, Rabenschlag F, Schori D, *et al.* Informal coercion as a neglected form of communication in psychiatric settings in Germany and Switzerland. *Psychiatry Res* 2018;262:400–6.
- 217 Gustafsson N, Salzmänn-Erikson M. Effect of complex working conditions on nurses who exert coercive measures in forensic psychiatric care. *J Psychosoc Nurs Ment Health Serv* 2016;54:37–43.
- 218 Jalil R, Huber JW, Sixsmith J, *et al.* Mental health nurses' emotions, exposure to patient aggression, attitudes to and use of coercive measures: cross sectional questionnaire survey. *Int J Nurs Stud* 2017;75:130–8.
- 219 Martello M, Doronina O, Perillo A, *et al.* Nurses' Perceptions of Engaging With Patients to Reduce Restrictive Practices in an Inpatient Psychiatric Unit. *Health Care Manag* 2018;37:342–53.
- 220 McKeown M, Thomson G, Scholes A, *et al.* "Catching your tail and firefighting": The impact of staffing levels on restraint minimization efforts. *J Psychiatr Ment Health Nurs* 2019;26:131–41.
- 221 Molewijk B, Kok A, Husum T, *et al.* Staff's normative attitudes towards coercion: the role of moral doubt and professional context—a cross-sectional survey study. *BMC Med Ethics* 2017;18.
- 222 Raveesh BN, Pathare S, Noorthoorn EO, *et al.* Staff and caregiver attitude to coercion in India. *Indian J Psychiatry* 2016;58:221–9.
- 223 Georgieva I, Mulder CL, Wiersma A. Patients' preference and experiences of forced medication and seclusion. *Psychiatr Q* 2012;83:1–13.
- 224 Haw C, Stubbs J, Bickle A, *et al.* Coercive treatments in forensic psychiatry: a study of patients' experiences and preferences. *J Forens Psychiatry Psychol* 2011;22:564–85.
- 225 Sequeira H, Halstead S. "Is it meant to hurt, is it?" Management of violence in women with developmental disabilities. *Violence against women* 2001;7:462–76.
- 226 Sustere E, Tarpey E. Least restrictive practice: its role in patient independence and recovery. *J Forens Psychiatry Psychol* 2019;30:614–29.
- 227 Larsen IB, Terkelsen TB. Coercion in a locked psychiatric ward: perspectives of patients and staff. *Nurs Ethics* 2014;21:426–36.
- 228 Whittington R, Bowers L, Nolan P, *et al.* Approval ratings of inpatient coercive interventions in a national sample of mental health service users and staff in England. *Psychiatr Serv* 2009;60:792–8.
- 229 Reisch T, Beeri S, Klein G, *et al.* Comparing attitudes to containment measures of patients, health care professionals and next of kin. *Front Psychiatry* 2018;9.
- 230 Rippon D, Reid K, Kay G. Views on restrictive practices on young people in psychiatric wards. *Nursing Times* 2018;114:24–8.
- 231 Ryan CJ, Bowers L. Coercive manoeuvres in a psychiatric intensive care unit. *J Psychiatr Ment Health Nurs* 2005;12:695–702.
- 232 Johnston MS, Kilty JM. "It's for their own good": Techniques of neutralization and security guard violence against psychiatric patients. *Punishment & Society* 2016;18:177–97.
- 233 Mackay I, Paterson B, Cassells C. Constant or special observations of inpatients presenting a risk of aggression or violence: nurses' perceptions of the rules of engagement. *J Psychiatr Ment Health Nurs* 2005;12:464–71.
- 234 Delaney KR, Johnson ME. Keeping the unit safe: Mapping psychiatric nursing skills. *J Am Psychiatr Nurses Assoc* 2006;12:198–207.
- 235 Happell B, Koehn S. Seclusion as a necessary intervention: the relationship between burnout, job satisfaction and therapeutic optimism and Justification for the use of seclusion. *J Adv Nurs* 2011;67:1222–31.
- 236 Johnson ME, Delaney KR. Keeping the unit safe: a grounded theory study. *J Am Psychiatr Nurses Assoc* 2006;12:13–21.
- 237 Jonker EJ, Goossens PJJ, Steenhuis IHM, *et al.* Patient aggression in clinical psychiatry: perceptions of mental health nurses. *J Psychiatr Ment Health Nurs* 2008;15:492–9.
- 238 Langan C, McDonald C. Daytime night attire as a therapeutic intervention in an acute adult psychiatric in-patient unit. *Psychiatric Bulletin* 2008;32:221–4.
- 239 Maguire T, Daffern M, Martin T. Exploring nurses' and patients' perspectives of limit setting in a forensic mental health setting. *Int J Ment Health Nurs* 2014;23:153–60.
- 240 Salzmänn-Erikson M, Lützén K, Ivarsson A-B, *et al.* The core characteristics and nursing care activities in psychiatric intensive care units in Sweden. *Int J Ment Health Nurs* 2008;17:98–107.
- 241 Millar R, Sands N. 'He did what? Well, that wasn't handed over!' Communicating risk in mental health. *J Psychiatr Ment Health Nurs* 2013;20:345–54.
- 242 Sjöstrand M, Sandman L, Karlsson P, *et al.* Ethical deliberations about involuntary treatment: interviews with Swedish psychiatrists. *BMC Med Ethics* 2015;16:37.
- 243 Woods P. Risk assessment and management approaches on mental health units. *J Psychiatr Ment Health Nurs* 2013;20:807–13.
- 244 Cowan D, Brunero S, Luo X, *et al.* Developing a guideline for structured content and process in mental health nursing handover. *Int J Ment Health Nurs* 2018;27:429–39.
- 245 Cullen SW, Nath SB, Marcus SC. Toward understanding errors in inpatient psychiatry: a qualitative inquiry. *Psychiatr Q* 2010;81:197–205.
- 246 Gifford ML, Anderson JE. Barriers and motivating factors in reporting incidents of assault in mental health care. *J Am Psychiatr Nurses Assoc* 2010;16:288–98.
- 247 Martin T, Daffern M. Clinician perceptions of personal safety and confidence to manage inpatient aggression in a forensic psychiatric setting. *J Psychiatr Ment Health Nurs* 2006;13:90–9.
- 248 Mezey G, Hassell Y, Bartlett A. Safety of women in mixed-sex and single-sex medium secure units: staff and patient perceptions. *Br J Psychiatry* 2005;187:579–82.

- 249 Brennan G, Flood C, Bowers L. Constraints and blocks to change and improvement on acute psychiatric wards - lessons from the City Nurses project. *J Psychiatr Ment Health Nurs* 2006;13:475-82.
- 250 Kuosmanen A, Tiihonen J, Repo-Tiihonen E, *et al.* Changes in patient safety culture: a patient safety intervention for Finnish forensic psychiatric hospital staff. *J Nurs Manag* 2019;27:848-57.
- 251 Lavelle M, Attoe C, Tritschler C, *et al.* Managing medical emergencies in mental health settings using an interprofessional in-situ simulation training programme: a mixed methods evaluation study. *Nurse Educ Today* 2017;59:103-9.
- 252 Bowers L, Simpson A, Eyres S, *et al.* Serious untoward incidents and their aftermath in acute inpatient psychiatry: the Tompkins acute ward study. *Int J Ment Health Nurs* 2006;15:226-34.
- 253 Gabrielsson S, Looi G-ME, Zingmark K, *et al.* Knowledge of the patient as decision-making power: staff members' perceptions of interprofessional collaboration in challenging situations in psychiatric inpatient care. *Scand J Caring Sci* 2014;28:784-92.
- 254 Ward L. Ready, aim fire! mental health nurses under siege in acute inpatient facilities. *Issues Ment Health Nurs* 2013;34:281-7.
- 255 Ryan P, Anczewska M, Laijarvi H, *et al.* Demographic and situational variations in levels of burnout in European mental health services: a comparative study. *Diversity Health Soc Care* 2007;4:101-12.
- 256 Nathan R, Brown A, Redhead K, *et al.* Staff responses to the therapeutic environment: a prospective study comparing burnout among nurses working on male and female wards in a medium secure unit. *J Forens Psychiatry Psychol* 2007;18:342-52.
- 257 Vlayen A *et al.* A nationwide Hospital survey on patient safety culture in Belgian hospitals: setting priorities at the Launch of a 5-year patient safety plan. *BMJ Publishing Group Ltd* 2012.
- 258 Ajalli A *et al.* Explanation of patient safety provided by nurses in inpatient psychiatric wards in Iran: a qualitative study. *Iranian Journal of Psychiatry and Behavioral Sciences* 2018;12.
- 259 Wood D, Pistrang N. A safe place? service users' experiences of an acute mental health ward. *J Community Appl Soc Psychol* 2004;14:16-28.
- 260 Jones J, NOLAN P, BOWERS L, *et al.* Psychiatric wards: places of safety? *J Psychiatr Ment Health Nurs* 2010;17:124-30.
- 261 Ireland CA, Ireland JL, Jones NS, *et al.* Predicting security incidents in high secure male psychiatric care. *Int J Law Psychiatry* 2019;64:40-52.
- 262 Pelto-Piri V, Wallsten T, Hylén U, *et al.* Feeling safe or unsafe in psychiatric inpatient care, a hospital-based qualitative interview study with inpatients in Sweden. *Int J Ment Health Syst* 2019;13:23
- 263 Haines A, Brown A, McCabe R, *et al.* Factors impacting perceived safety among staff working on mental health wards. *BJPsych Open* 2017;3:204-11.
- 264 Kanerva A, Kivinen T, Lammintakanen J. Communication elements supporting patient safety in psychiatric inpatient care. *J Psychiatr Ment Health Nurs* 2015;22:298-305.
- 265 Kuosmanen A, Tiihonen J, Repo-Tiihonen E, *et al.* Patient safety culture in two Finnish state-run forensic psychiatric hospitals. *J Forensic Nurs* 2013;9:207-16.
- 266 Wu J-C, Tung T-H, Chen PY, *et al.* Determinants of workplace violence against clinical physicians in hospitals. *J Occup Health* 2015;57:540-7.
- 267 Kanerva A, Lammintakanen J, Kivinen T. Nursing staff's perceptions of patient safety in psychiatric inpatient care. *Perspect Psychiatr Care* 2016;52:25-31.
- 268 Stead K, Kumar S, Schultz TJ, *et al.* Teams communicating through STEPPS. *Medical Journal of Australia* 2009;190:S128-32.
- 269 Mahoney JS, Ellis TE, Garland G, *et al.* Supporting a psychiatric hospital culture of safety. *J Am Psychiatr Nurses Assoc* 2012;18:299-306.
- 270 Higgins N, Meehan T, Dart N, *et al.* Implementation of the Safewards model in public mental health facilities: a qualitative evaluation of staff perceptions. *Int J Nurs Stud* 2018;88:114-20.
- 271 Bowers L, Gournay K, Duffy D. Suicide and self-harm in inpatient psychiatric units: a national survey of observation policies. *J Adv Nurs* 2000;32:437-44.
- 272 O'Brien L, Cole R. Mental health nursing practice in acute psychiatric close-observation areas. *Int J Ment Health Nurs* 2004;13:89-99.
- 273 Stein WM. The use of discharge risk assessment tools in general psychiatric services in the UK. *J Psychiatr Ment Health Nurs* 2002;9:713-24.
- 274 Stübner S, GROß G, Nedopil N. Inpatient risk management with mentally ill offenders: results of a survey on clinical decision-making about easing restrictions. *Criminal Behaviour and Mental Health* 2006;16:111-23.
- 275 O'Neill C, Heffernan P, Goggins R, *et al.* Long-Stay forensic psychiatric inpatients in the Republic of Ireland: aggregated needs assessment. *Ir J Psychol Med* 2003;20:119-25.
- 276 Rees P, Manthorpe J. *Managers' and staff experiences of adult protection allegations in mental health and learning disability residential services: a qualitative study.* *Br J Soc Work* 2010;40:513-29.
- 277 Koukia E, Giannouli E, Gonis N, *et al.* Security rules and banned items in psychiatric acute admission wards in Athens, Greece. *Int J Ment Health Nurs* 2010;19:428-36.
- 278 Silvana S, Laura F, Ursula DF, *et al.* Ergonomics in the psychiatric ward towards workers or patients? *Work* 2012;41 Suppl 1:1832-5.
- 279 True G, Frasso R, Cullen SW, *et al.* Adverse events in Veterans Affairs inpatient psychiatric units: staff perspectives on contributing and protective factors. *Gen Hosp Psychiatry* 2017;48:65-71.
- 280 Vandewalle J, Malfait S, Eeckloo K, *et al.* Patient safety on psychiatric wards: a cross-sectional, multilevel study of factors influencing nurses' willingness to share power and responsibility with patients. *Int J Ment Health Nurs* 2018;27:877-90.
- 281 Gerace A, Muir-Cochrane E. Perceptions of nurses working with psychiatric consumers regarding the elimination of seclusion and restraint in psychiatric inpatient settings and emergency departments: an Australian survey. *Int J Ment Health Nurs* 2019;28:209-25.
- 282 Vahidi M, Ebrahimi H, Areshtanab HN, *et al.* Therapeutic relationships and safety of care in Iranian psychiatric inpatient units. *Issues Ment Health Nurs* 2018;39:967-76.
- 283 Holth F, Walby F, Røstbakken T, *et al.* Extreme challenges: psychiatric inpatients with severe self-harming behavior in Norway: a national screening investigation. *Nord J Psychiatry* 2018;72:605-12.
- 284 de Jonghe-Rouleau AP, Pot AM, de Jonghe JFM. Self-Injurious behaviour in nursing home residents with dementia. *Int J Geriatr Psychiatry* 2005;20:651-7.
- 285 Sansone RA, McLean JS, Wiederman MW. The relationship between medically self-sabotaging behaviors and borderline personality disorder among psychiatric inpatients. *Prim Care Companion J Clin Psychiatry* 2008;10:448-52.
- 286 Gough K, Hawkins A. Staff attitudes to Self-harm and its management in a forensic psychiatric service. *The British Journal of Forensic Practice* 2000;2:22-8.
- 287 O'Donovan A. Pragmatism rules: the intervention and prevention strategies used by psychiatric nurses working with non-suicidal self-harming individuals. *J Psychiatr Ment Health Nurs* 2007;14:64-71.
- 288 Tofthagen R, Talseth A-G, Fagerström L. Mental Health Nurses' Experiences of Caring for Patients Suffering from Self-Harm. *Nurs Res Pract* 2014;2014:1-10.
- 289 Lundegaard Mattson Åse, Binder P-E, Mattson L. A qualitative exploration of how health care workers in an inpatient setting in Norway experience working with patients who self-injure. *Nordic Psychology* 2012;64:272-90.
- 290 Thomas JB, Haslam CO. *How people who self-harm negotiate the inpatient environment: the mental healthcare workers perspective.* *Journal of Psychiatric & Mental Health Nursing* (John Wiley & Sons, Inc 2017;24:480-90.
- 291 Shaw DG, Sandy PT. *Mental health nurses' attitudes toward self-harm: Curricular implications.* *Health Sa Gesondheid* 2016;21.
- 292 Sandy PT. *The use of observation on patients who self-harm: Lessons from a learning disability service.* *Health Sa Gesondheid* 2016;21.
- 293 James K, Samuels I, Moran P, *et al.* Harm reduction as a strategy for supporting people who self-harm on mental health wards: the views and experiences of practitioners. *J Affect Disord* 2017;214:67-73.
- 294 Brown J, Beail N. Self-Harm among people with intellectual disabilities living in secure service provision: a qualitative exploration. *Journal of Applied Research in Intellectual Disabilities* 2009;22:503-13.
- 295 Lindgren B-M, Aminoff C, Hällgren Graneheim U. Features of everyday life in psychiatric inpatient care for self-harming: an observational study of six women. *Issues Ment Health Nurs* 2015;36:82-8.
- 296 Weber MT. Triggers for self-abuse: a qualitative study. *Arch Psychiatr Nurs* 2002;16:118-24.
- 297 Gibson J, Booth R, Davenport J, *et al.* Dialectical behaviour therapy-informed skills training for deliberate self-harm: a controlled trial with 3-month follow-up data. *Behav Res Ther* 2014;60:8-14.
- 298 Booth R, Keogh K, Doyle J, *et al.* Living through distress: a skills training group for reducing deliberate self-harm. *Behav Cogn Psychother* 2014;42:156-65.

- 299 Bowers L, Whittington R, Nolan P, *et al.* Relationship between service ecology, special observation and self-harm during acute in-patient care: City-128 study. *Br J Psychiatry* 2008;193:395–401.
- 300 Kool N, van Meijel B, Koekoek B, *et al.* Improving communication and practical skills in working with inpatients who self-harm: a pre-test/post-test study of the effects of a training programme. *BMC Psychiatry* 2014;14:64.
- 301 Drew BL. No-suicide contracts to prevent suicidal behavior in inpatient psychiatric settings. *Journal of the American Psychiatric Nurses Association* 1999;5:23–8.
- 302 Sjöström N, Hetta J, Waern M. Sense of coherence and suicidality in suicide attempters: a prospective study. *J Psychiatr Ment Health Nurs* 2012;19:62–9.
- 303 Swogger MT, Van Orden KA, Conner KR. The relationship of outwardly directed aggression to suicidal ideation and suicide attempts across two high-risk samples. *Psychol Violence* 2014;4:184–95.
- 304 Hill RM, Hatkevich CE, Kazimi I, *et al.* The Columbia-Suicide severity rating scale: associations between interrupted, aborted, and actual suicide attempts among adolescent inpatients. *Psychiatry Res* 2017;255:338–40.
- 305 Inoue K, Kawanishi C, Otsuka K, *et al.* A large-scale survey of inpatient suicides: comparison between medical and psychiatric settings. *Psychiatry Res* 2017;250:155–8.
- 306 Ellis TE, Green KL, Allen JG, *et al.* Collaborative assessment and management of suicidality in an inpatient setting: results of a pilot study. *Psychotherapy* 2012;49:72–80.
- 307 Ellis TE, Rufino KA, Allen JG, *et al.* Impact of a Suicide-Specific intervention within inpatient psychiatric care: the collaborative assessment and management of suicidality. *Suicide Life Threat Behav* 2015;45:556–66.
- 308 Awenat YF, Peters S, Gooding PA, *et al.* A qualitative analysis of suicidal psychiatric inpatients views and expectations of psychological therapy to counter suicidal thoughts, acts and deaths. *BMC Psychiatry* 2018;18:334.
- 309 Cleary M, Jordan R, Horsfall J, *et al.* Suicidal patients and special observation. *J Psychiatr Ment Health Nurs* 1999;6:461–7.
- 310 Takahashi C, Chida F, Nakamura H, *et al.* The impact of inpatient suicide on psychiatric nurses and their need for support. *BMC Psychiatry* 2011;11:38.
- 311 Vråle GB, Steen E. The dynamics between structure and flexibility in constant observation of psychiatric inpatients with suicidal ideation. *J Psychiatr Ment Health Nurs* 2005;12:513–8.
- 312 Vandewalle J, Beeckman D, Van Hecke A, *et al.* 'Promoting and preserving safety and a life-oriented perspective': A qualitative study of nurses' interactions with patients experiencing suicidal ideation. *Int J Ment Health Nurs* 2019;28:1122–34.
- 313 Davis SE, Williams IS, Hays LW. Psychiatric inpatients' perceptions of written no-suicide agreements: an exploratory study. *Suicide Life Threat Behav* 2002;32:51–66.
- 314 Esposito-Smythers C, McClung TJ, Fairlie AM. Adolescent perceptions of a suicide prevention group on an inpatient unit. *Archives of Suicide Research* 2006;10:265–75.
- 315 Sun F-K, Long A, Boore J, *et al.* Nursing people who are suicidal on psychiatric wards in Taiwan: action/interaction strategies. *J Psychiatr Ment Health Nurs* 2005;12:275–82.
- 316 Sun F-K, Long A, Boore J, *et al.* Patients and nurses' perceptions of ward environmental factors and support systems in the care of suicidal patients. *J Clin Nurs* 2006;15:83–92.
- 317 Pfeiffer PNet *et al.* Development and pilot study of a suicide prevention intervention delivered by peer support specialists. *Psychological Services* 2018.
- 318 Caspi E. Does self-neglect occur among older adults with dementia when unsupervised in assisted living? an exploratory, observational study. *J Elder Abuse Negl* 2014;26:123–49.
- 319 Bowers L, HAGLUND K, MUIR-COCHRANE E, *et al.* Locked doors: a survey of patients, staff and visitors. *J Psychiatr Ment Health Nurs* 2010;17:873–80.
- 320 Haglund K, von Essen L. Locked entrance doors at psychiatric wards – advantages and disadvantages according to voluntarily admitted patients. *Nord J Psychiatry* 2005;59:511–5.
- 321 Muir-Cochrane E, van der Merwe M, Nijman H, *et al.* Investigation into the acceptability of door locking to staff, patients, and visitors on acute psychiatric wards. *Int J Ment Health Nurs* 2012;21:41–9.
- 322 Cowman S, Bowers L. Safety and security in acute admission psychiatric wards in Ireland and London: a comparative study. *J Clin Nurs* 2009;18:1346–53.
- 323 Simpson A, Bowers L, Haglund K, *et al.* The relationship between substance use and exit security on psychiatric wards. *J Adv Nurs* 2011;67:519–30.
- 324 Kalagi J, Otte I, Vollmann J, *et al.* Requirements for the implementation of open door policies in acute psychiatry from a mental health professionals' and patients' view: a qualitative interview study. *BMC Psychiatry* 2018;18:304.
- 325 Fletcher J, Hamilton B, Kinner S, *et al.* Working towards least restrictive environments in acute mental health wards in the context of locked door policy and practice. *Int J Ment Health Nurs* 2019;28:538–50.
- 326 van der Schaaf PS, Dusseldorp E, Keuning FM, *et al.* Impact of the physical environment of psychiatric wards on the use of seclusion. *British Journal of Psychiatry* 2013;202:142–9.
- 327 Verbeek H, Zwakhalen SMG, van Rossum E, *et al.* Effects of small-scale, home-like facilities in dementia care on residents' behavior, and use of physical restraints and psychotropic drugs: a quasi-experimental study. *International Psychogeriatrics* 2014;26:657–68.
- 328 Curtis S, Gesler W, Wood V, *et al.* Compassionate containment? balancing technical safety and therapy in the design of psychiatric wards. *Soc Sci Med* 2013;97:201–9.
- 329 Dreyfus S, Phillipson L, Fleming R. Staff and family attitudes to fences as a means of detaining people with dementia in residential aged care settings: the tension between physical and emotional safety. *Australian Journal of Social Issues* 2018;53:107–22.
- 330 Bayramzadeh S. An assessment of levels of safety in psychiatric units. *HERD* 2017;10:66–80.
- 331 Bellantonio S, Kenny AM, Fortinsky RH, *et al.* Efficacy of a geriatrics team intervention for residents in Dementia-Specific assisted living facilities: effect on unanticipated transitions. *J Am Geriatr Soc* 2008;56:523–8.
- 332 Chandler G. From traditional inpatient to trauma-informed treatment: transferring control from staff to patient. *J Am Psychiatr Nurses Assoc* 2008;14:363–71.
- 333 Wilkes L, Fleming A, Wilkes BL, *et al.* Environmental approach to reducing agitation in older persons with dementia in a nursing home. *Australas J Ageing* 2005;24:141–5.
- 334 Gebhardt RP, Steinert T. Should severely disturbed psychiatric patients be distributed or concentrated in specialized wards? an empirical study on the effects of hospital organization on ward atmosphere, aggressive behavior, and sexual molestation. *Eur Psychiatry* 1999;14:291–7.
- 335 Stolker JJ, Nijman HLI, Zwanikken P-H. Are patients' views on seclusion associated with lack of privacy in the ward? *Arch Psychiatry Nurs* 2006;20:282–7.
- 336 Kulkarni J, Gavrilidis E, Lee S, *et al.* Establishing female-only areas in psychiatry wards to improve safety and quality of care for women. *Australasian Psychiatry* 2014;22:551–6.
- 337 Bowers L, Crowder M. Nursing staff numbers and their relationship to conflict and containment rates on psychiatric wards—A cross sectional time series poisson regression study. *Int J Nurs Stud* 2012;49:15–20.
- 338 Triplett P, Dearholt S, Cooper M, *et al.* The milieu manager: a nursing staffing strategy to reduce observer use in the acute psychiatric inpatient setting. *J Am Psychiatr Nurses Assoc* 2017;23:422–30.
- 339 Hunt IM, Windfuhr K, Shaw J, *et al.* Ligature points and ligature types used by psychiatric inpatients who die by hanging: a national study. *Crisis* 2012;33:87–94.
- 340 Ruzić K, Francisković T, Suković Z, *et al.* Aggressiveness in institutionalised schizophrenic patients and the selection of antipsychotics. *Coll Antropol* 2011;35:265–9.
- 341 Rodríguez-Leal CM, López-Lunar E, Carrascosa-Bernáldez JM, *et al.* Electrocardiographic surveillance in a psychiatric institution: avoiding iatrogenic cardiovascular death. *Int J Psychiatry Clin Pract* 2017;21:64–6.
- 342 Seemüller F, Riedel M, Obermeier M, *et al.* The controversial link between antidepressants and suicidality risks in adults: data from a naturalistic study on a large sample of in-patients with a major depressive episode. *Int J Neuropsychopharm* 2009;12:181–9.
- 343 Haw C, Stubbs J, Dickens GL. Barriers to the reporting of medication administration errors and near misses: an interview study of nurses at a psychiatric hospital. *J Psychiatr Ment Health Nurs* 2014;21:n/a–805.
- 344 Soerensen AL *et al.* The medication process in a psychiatric Hospital: are errors a potential threat to patient safety? *Risk management and healthcare policy* 2013;6:23.
- 345 Keers RN, Plácido M, Bennett K, *et al.* What causes medication administration errors in a mental health hospital? A qualitative study with nursing staff. *PLoS One* 2018;13:e0206233.
- 346 Bademli K, Buldukoglu K. Oral medication management in inpatient psychiatric care in turkey. *J Psychiatr Ment Health Nurs* 2009;16:355–62.

- 347 Prins MC, Drenth-van Maanen AC, Kok RM, *et al.* Use of a structured medication history to establish medication use at admission to an old age psychiatric clinic: a prospective observational study. *CNS Drugs* 2013;27:963–9.
- 348 Xie N, Kalia K, Strudwick G, *et al.* Understanding mental health nurses' perceptions of barcode medication administration: a qualitative descriptive study. *Issues Ment Health Nurs* 2019;40:326–34.
- 349 Strudwick G, Clark C, McBride B, *et al.* Thank you for asking: exploring patient perceptions of barcode medication administration identification practices in inpatient mental health settings. *Int J Med Inform* 2017;105:31–7.
- 350 Cottney A, Innes J. Medication-administration errors in an urban mental health Hospital: a direct observation study. *Int J Ment Health Nurs* 2015;24:65–74.
- 351 Dickens G, Stubbs J, Haw C, Raw C. Delegation of medication administration: an exploratory study. *Nursing Standard* 2008;22:35–40.
- 352 Haw C, Stubbs J, Dickens G. An observational study of medication administration errors in old-age psychiatric inpatients. *International Journal for Quality in Health Care* 2007;19:210–6.
- 353 Cottney A. Improving the safety and efficiency of nurse medication rounds through the introduction of an automated dispensing cabinet. *BMJ Open Quality* 2014;3.
- 354 Dolan M, Kirwan H. Survey of staff perceptions of illicit drug use among patients in a medium secure unit. *Psychiatric Bulletin* 2001;25:14–17.
- 355 Hughes E, Bressington D, Sharratt K, *et al.* Novel psychoactive substance use by mental health service consumers: an online survey of inpatient health professionals' views and experiences. *Adv Dual Diagn* 2018;11:30–9.
- 356 Gonzalez-Pinto A, De Azua SR. S.2.04 adherence to treatment in bipolar disorders. *European Neuropsychopharmacology* 2011;21:S108–9.
- 357 Meehan T, Morrison P, McDougall S. Absconding behaviour: an exploratory investigation in an acute inpatient unit. *Aust N Z J Psychiatry* 1999;33:533–7.
- 358 Muir-Cochrane E, Oster C, Grotto J, *et al.* The inpatient psychiatric unit as both a safe and unsafe place: implications for absconding. *Int J Ment Health Nurs* 2013;22:304–12.
- 359 Nurjannah I, FitzGerald M, Foster K. Patients' experiences of absconding from a psychiatric setting in Indonesia. *Int J Ment Health Nurs* 2009;18:326–35.
- 360 Grotto J, Gerace A, O'Kane D, *et al.* Risk assessment and absconding: perceptions, understandings and responses of mental health nurses. *J Clin Nurs* 2015;24:855–65.
- 361 Bowers L, Alexander J, Gaskell C. A trial of an anti-absconding intervention in acute psychiatric wards. *J Psychiatr Ment Health Nurs* 2003;10:410–6.
- 362 Simpson AIF, Penney SR, Fernane S, *et al.* The impact of structured decision making on absconding by forensic psychiatric patients: results from an A-B design study. *BMC Psychiatry* 2015;15:103.
- 363 Algase DL, Beattie ERA, Antonakos C, *et al.* Wandering and the physical environment. *Am J Alzheimers Dis Other Demen* 2010;25:340–6.
- 364 Colombo M, Vitali S, Cairati M, *et al.* Wanderers: features, findings, issues. *Arch Gerontol Geriatr* 2001;33:99–106.
- 365 Hunt IM, Clements C, Saini P, *et al.* Suicide after absconding from inpatient care in England: an exploration of mental health professionals' experiences. *Journal of Mental Health* 2016;25:245–53.
- 366 Hunt IM, Windfuhr K, Swinson N, *et al.* Suicide amongst psychiatric in-patients who abscond from the ward: a national clinical survey. *BMC Psychiatry* 2010;10:14.
- 367 Nijman Het al. Door locking and exit security measures on acute psychiatric admission wards. *Journal of Psychiatric and Mental Health Nursing* 2011;18): :614–21.
- 368 Whaley AL. A two-stage method for the study of cultural bias in the diagnosis of schizophrenia in African Americans. *Journal of Black Psychology* 2004;30:167–86.
- 369 Whaley AL. Cultural mistrust and the clinical diagnosis of paranoid schizophrenia in African American patients. *J Psychopathol Behav Assess* 2001;23:93–100.
- 370 Green R, Shelly C, Gibb J, *et al.* Implementing seclusion in forensic mental health care: a qualitative study of staff decision making. *Arch Psychiatr Nurs* 2018;32:764–8.
- 371 Koukia E, Madianos MG, Katostaras T. "On the spot" interventions by mental health nurses in inpatient psychiatric wards in Greece. *Issues Ment Health Nurs* 2009;30:327–36.
- 372 Lindsey PL. Psychiatric nurses' decision to restrain: the association between empowerment and individual factors. *Journal of psychosocial nursing and mental health services* 2009;47:41–9.
- 373 Mann-Poll PS, Smit A, de Vries WJ, *et al.* Factors contributing to mental health professionals' decision to use seclusion. *Psychiatric Services* 2011;62:498–503.
- 374 Marangos-Frost S, Wells D. Psychiatric nurses' thoughts and feelings about restraint use: a decision dilemma. *J Adv Nurs* 2000;31:362–9.
- 375 Brown B, Rakow T. Understanding clinicians' use of cues when assessing the future risk of violence: a clinical judgement analysis in the psychiatric setting. *Clin Psychol Psychother* 2016;23:125–41.
- 376 Fuller J, Cowan J. Risk assessment in a multi-disciplinary forensic setting: clinical judgement revisited. *The Journal of Forensic Psychiatry* 1999;10:276–89.
- 377 Abraham S. Managing patient falls in psychiatric inpatient units. *Health Care Manag* 2016;35:21–7.
- 378 Fonad E, Emami A, Wahlin T-BR, *et al.* Falls in somatic and dementia wards at community care units. *Scand J Caring Sci* 2009;23:2–10.
- 379 Tångman S, Eriksson S, Gustafson Y, *et al.* Precipitating factors for falls among patients with dementia on a psychogeriatric ward. *International Psychogeriatrics* 2010;22:641–9.
- 380 Garfinkel D, Radomislky Z, Jamal S, *et al.* High efficacy for hip protectors in the prevention of hip fractures among elderly people with dementia. *J Am Med Dir Assoc* 2008;9:313–8.
- 381 Holmes Det al. An evaluation of a monitoring system intervention: falls, injuries, and affect in nursing homes. *Clinical nursing research* 2007;16:317–35.
- 382 Powell-Cope G, Quigley P, Besterman-Dahan K, *et al.* A qualitative understanding of patient falls in inpatient mental health units. *J Am Psychiatr Nurses Assoc* 2014;20:328–39.
- 383 Li PHet al. Infection preventionists' challenges in psychiatric clinical settings. *American Journal of Infection Control* 2019;47:123–7.
- 384 D'Lima D, Crawford MJ, Darzi A, *et al.* Patient safety and quality of care in mental health: a world of its own? *BJPsych Bulletin* 2017;41:241–3.
- 385 Sakinofsky I. Preventing suicide among inpatients. *The Canadian Journal of Psychiatry* 2014;59:131–40.
- 386 Leape L, Berwick D, Clancy C, *et al.* Transforming healthcare: a safety imperative. *Qual Saf Health Care* 2009;18:424–8.
- 387 Gandhi TK, Kaplan GS, Leape L, *et al.* Transforming concepts in patient safety: a progress report. *BMJ Qual Saf* 2018;27:1019–26.
- 388 Schwappach DLB. Review: engaging patients as vigilant partners in safety: a systematic review. *Med Care Res Rev* 2010;67:119–48.
- 389 Davis RE, Sevdalis N, Vincent CA. Patient involvement in patient safety: how willing are patients to participate? *BMJ Qual Saf* 2011;20:108–14.
- 390 Institute, N.P.S.F.s.L.L.. *Safety is personal: partnering with patients and families for the safest care*. Moston: MA.: National Patient Safety Foundation, 2014.
- 391 Providers N. *Funding for mental health at a local level: Unpicking the variation*. NHS Providers 2016.
- 392 Prince M, Patel V, Saxena S, *et al.* No health without mental health. *The Lancet* 2007;370:859–77.
- 393 Unsworth J, McKeever M, Kelleher M. Recognition of physical deterioration in patients with mental health problems: the role of simulation in knowledge and skill development. *J Psychiatr Ment Health Nurs* 2012;19:536–45.
- 394 Cooper JB, Gaba DM, Liang B, *et al.* The National patient safety Foundation agenda for research and development in patient safety. *MedGenMed* 2000;2:E38.
- 395 Institute of Medicine Committee on Quality of Health Care in, A. *In To Err is Human: Building a Safer Health System*, L.T. Kohn, J.M. Corrigan, and M.S. Donaldson, Editors. 2000, National Academies Press (US) Copyright 2000 by the National Academy of Sciences. All rights reserved. Washington (DC).
- 396 Emanuel L, Berwick D, Conway J, *et al.* Advances in Patient Safety: New Directions and Alternative Approaches (Vol. 1: Assessment). In: *What exactly is patient safety?* Bethesda MD, USA: U.S. National Library of Medicine, 2008.
- 397 Hox JJ, Boeije HR. Data collection, primary versus secondary 2005.
- 398 Appleby L, Shaw J, Amos T. National confidential inquiry into suicide and homicide by people with mental illness. *Br J Psychiatry* 1997;170:101–2.
- 399 Elliott JH, Turner T, Clavisi O, *et al.* Living systematic reviews: an emerging opportunity to narrow the evidence-practice gap. *PLoS Med* 2014;11:e1001603.
- 400 Long L. Routine piloting in systematic reviews—a modified approach? *Syst Rev* 2014;3:77.

- 401 Landis JR, Koch GG. The measurement of observer agreement for categorical data. *Biometrics* 1977;33:159–74.
- 402 Gopalakrishnan S, Ganeshkumar P. Systematic reviews and meta-analysis: understanding the best evidence in primary healthcare. *J Family Med Prim Care* 2013;2.
- 403 Dewa LH, Murray K, Thibaut B, *et al.* Identifying research priorities for patient safety in mental health: an international expert Delphi study. *BMJ Open* 2018;8:e021361.