BMJ Open Cost-effectiveness of telephone coaching for physically inactive ambulatory care hospital patients: economic evaluation alongside the Healthy4U randomised controlled trial

Stephen Barrett ¹, Stephen Begg,² Paul O'Halloran,³ Michael Kingsley²

To cite: Barrett S, Begg S, O'Halloran P, *et al*. Cost-effectiveness of telephone coaching for physically inactive ambulatory care hospital patients: economic evaluation alongside the Healthy4U randomised controlled trial. *BMJ Open* 2019;9:e032500. doi:10.1136/bmjopen-2019-032500

► Prepublication history and additional material for this paper are available online. To view these files, please visit the journal online (<http://dx.doi.org/10.1136/bmjopen-2019-032500>).

Received 21 June 2019

Revised 31 October 2019

Accepted 25 November 2019

© Author(s) (or their employer(s)) 2019. Re-use permitted under CC BY-NC. No commercial re-use. See rights and permissions. Published by BMJ.

For numbered affiliations see end of article.

Correspondence to

Prof Michael Kingsley, Holsworth Research Initiative, La Trobe University, Bendigo, Victoria, Australia; M.Kingsley@latrobe.edu.au

ABSTRACT

Objective To assess whether telephone coaching is a cost-effective method for increasing physical activity and health-related quality of life for insufficiently active adults presenting to an ambulatory care clinic in a public hospital.

Design An economic evaluation was performed alongside a randomised controlled trial.

Setting Participants were recruited from an ambulatory care clinic in a public hospital in regional Australia.

Participants Seventy-two adults (aged 18–69) deemed insufficiently physically active via self-report.

Interventions Participants were randomised to either an intervention group that received an education session and eight sessions of telephone coaching over a 12-week period, or to a control group that received the education session only. The intervention used in the telephone coaching was integrated motivational interviewing and cognitive behavioural therapy.

Outcome measures The primary health outcome was change in moderate-to-vigorous physical activity (MVPA), objectively measured via accelerometry. The secondary outcome was the quality-adjusted life-year (QALY) determined by the 12-item Short Form Health Survey Questionnaire. Outcome data were measured at baseline, postintervention (3 months) and follow-up (6 months). Incremental cost-effectiveness ratios (ICERs) were calculated for each outcome. Non-parametric bootstrapping techniques and sensitivity analyses were performed to account for uncertainty.

Results The mean intervention cost was \$279±\$13 per person. At 6 months follow-up, relative to control, the intervention group undertook 18 more minutes of daily MVPA at an ICER of \$15/min for each additional minute of MVPA. With regard to QALYs, the intervention yielded an ICER of \$36 857 per QALY gained. Sensitivity analyses indicated that results were robust to varied assumptions.

Conclusion Telephone coaching was a low-cost strategy for increasing MVPA and QALYs in insufficiently physically active ambulatory care hospital patients. Additional research could explore the potential economic impact of the intervention from a broader healthcare perspective.

Trial registration number ANZCTR: ACTRN12616001331426.

Strengths and limitations of this study

- This study is the first to investigate the cost-effectiveness of telephone delivered motivational interviewing and cognitive behavioural therapy for insufficiently physically active secondary care hospital patients.
- Economic evaluations enable hospitals to estimate the costs of delivering preventive health interventions, which are important for addressing the burden of chronic disease.
- Changes in physical activity were measured using accelerometers, strengthening the confidence in the findings.
- Results are limited by a short time horizon and a narrow costing perspective.
- Further research is needed to explore the potential long-term economic impact of the intervention from a broader healthcare perspective.

INTRODUCTION

Insufficient physical activity (PA) is an established risk factor for the development of a number of chronic diseases, including cardiovascular disease, type 2 diabetes and obesity.¹ Despite the well-established benefits of PA,² more than half of the population does not attain sufficient levels of PA to derive such benefits.³ The estimated cost of insufficient PA in Australia is \$A805 million per annum.⁴ Addressing the prevalence of insufficient PA is a major public health priority, and necessitates that multiple sectors of the healthcare industry are actively engaged in PA promotion.⁵

Hospitals are important settings in which to offer health promotion interventions, for both admitted and ambulatory care. In the hospital setting, ambulatory care refers to non-admitted clinics that patients attend for specialist medical care. Patients attending

ambulatory care hospital clinics are more likely than the general population to have one or more chronic disease.^{6,7} Hospital patients can be motivated to engage with lifestyle behaviour change as their health is already compromised.^{8,9} Attending a hospital has been identified as a major life event,¹⁰ and a hospital visit has the potential to initiate health behaviour change.⁸ Ambulatory care settings provide an ideal opportunity for behaviour change interventions.^{8,10} Substantial efforts have been made to promote increased engagement in PA using individual and population-based approaches.¹¹ This has resulted in an increased use of behaviour change interventions to influence participation in PA.¹²

A number of studies suggest that telephone coaching results in improved clinical outcomes, self-efficacy and health status,^{13,14} as well as increases in PA.^{15,16} Additional work is required to embed telephone coaching within existing health services.^{13,15} To make the benefits of telephone coaching more broadly available for hospital outpatients, telephone coaching has been delivered in addition to standard ambulatory care.¹⁷ The addition of telephone coaching to standard care resulted in significant improvements in objectively measured PA and health-related outcomes;¹⁷ however, the cost to improve these outcomes has not been reported.

Few studies have employed any form of economic analyses on telephone coaching, and little is known about the relative cost-effectiveness of adding telephone coaching to routine care in ambulatory care hospital settings. The purpose of this study was to evaluate the cost-effectiveness of the Healthy4U programme for increasing measured PA and the number of quality-adjusted life-years (QALYs) experienced over a 6-month period from a hospital perspective.

METHODS

Study design

The Healthy4U study was a single-blind randomised controlled trial (RCT) reported in line with the CHEERS reporting guidelines¹⁸ (online supplementary file 1). The trial design, participants, sample size, intervention, outcomes and ethics approvals have been described in detail elsewhere.¹⁷ Briefly, between October 2016 and December 2017, 72 insufficiently physically active adults, aged 18–69 years, were recruited from ambulatory care clinics at a major hospital in a regional town in Victoria, Australia. The primary aim was to promote change in objectively measured PA during the trial.

Intervention

All enrolled participants attended a 30-min group education session. The education session was a facilitated learning session focused on self-management and lifestyle modification, and was carried out using a self-determination theory framework.¹⁹

The intervention group completed a telephone-based, integrated motivational interviewing and cognitive

behavioural therapy (MI-CBT) intervention, delivered in eight 30-min sessions over 12 weeks. The intervention was delivered by an experienced allied health clinician trained in MI-CBT. All participants enrolled into the control arm attended the education session. Apart from contact regarding follow-up outcome measures, participants in the control group received no further contact initiated by the research team.

Measurement of effects

Outcome measures were recorded at baseline, after 3 months of intervention (postintervention) and at 6 months (follow-up) by assessors blinded to the study group assignment. The primary outcome measure was change in moderate-to-vigorous physical activity (MVPA), objectively measured by accelerometry (wGT3X-BT; Actigraph, USA). Daily MVPA was determined using the manufacturers software (Actilife; Actigraph, USA) and the Freedson Adult (1998) cut point (vector magnitude >1961 cpm).²⁰ To be included in the analysis, a minimum wear time of ≥10 hour/day for 5 of the 7-day period was required, including at least one weekend day.²¹ Weekly PA totals were summed from the daily totals for persons with 7 valid days of monitoring, or estimated as seven times the average daily total for persons with 5–6 valid days of monitoring. Using the summed weekly totals, participants were classified as either meeting or not meeting the recommended PA guidelines.²²

A secondary outcome was a change in health-related quality of life (HrQoL) and QALYs, which was derived from the Medical Outcomes Study Short Form 12 Health Survey (SF-12) and the standard Brazier algorithm.^{23,24} The SF-12 scores were converted to utility scores on a scale of 0–1, with a higher score indicating a more favourable health state.²⁴ These utility estimates were converted to QALYs by calculating the ‘area under the curve’ utility estimates for the different follow-up time intervals for each participant, weighted by the length of follow-up at that time interval.

Measurement of costs

The cost analysis was designed and conducted from a hospital perspective, which allows healthcare organisations to gauge the approximate cost of offering this programme.²⁵ A bottom-up microcosting approach was used to calculate the intervention costs.²⁶ This approach involves the detailed collection of information regarding the quantities of resources consumed while implementing and executing the interventions, as well as their respective unit prices.²⁶ Only those costs involved in implementing the intervention (eg, training of individuals carrying out and undertaking the intervention) were included. Protocol-driven costs, namely the costs of gathering data as part of the clinical trial were considered to be sunk costs and were therefore excluded from the cost-effectiveness analysis.²⁵

The programme costs included group facilitator time, intervention time and supplies. The group facilitator’s

time was spent preparing for and facilitating group meetings. The intervention assistant's time was calculated as the time spent undertaking reminder phone calls to participants. The intervention costs were calculated as the time spent in one-to-one consultation with the participants. Both group facilitation and intervention costs were calculated using the annual salary of an experienced allied health clinician as they would most likely be used in delivering MI-CBT were the intervention to be implemented on a large scale (\$A82 924). Intervention assistant costs were based on the annual salary of an Allied Health Assistant (\$A45 338). Finally, the costs of supplies, including the programme manuals, were included in the programme cost.

The group facilitator's time per group meeting was estimated at 2.5 hours, which included 0.5 hours for the group meeting itself, 1.0 hour to set up before and clean up after the group meeting, and 1.0 hour to prepare for the group meeting (ie, reviewing meeting notes and presentation material). The group facilitator's cost per meeting per participant was calculated by dividing the facilitator's cost per meeting by the number of participants who attended each meeting. Due to the short time-frame in which costs and effects occurred, discounting was not necessary.²⁷ All programme costs were calculated in 2017 Australian dollars (AU\$).

Statistical analysis

Analyses of trial data have been reported elsewhere.¹⁷ In brief, mixed-model analysis of variances were used to assess the effects of the intervention on each of the outcome variables. The mean±SD for the overall cost and for the change in each outcome at 6 months was calculated. For each outcome, the incremental cost-effectiveness ratio (ICER) was calculated by dividing the difference in costs by the difference in effects between the intervention and control groups. The difference in effects between the two groups was calculated using a change from baseline approach to control for different baseline

utilities. Uncertainty in the ICER estimates was accounted for by generating 1000 bootstrap replicates of the data set, a widely used method in health economic evaluations. Probabilistic sensitivity analysis was completed by calculating the cost-effectiveness acceptability curve (CEAC) derived from the bootstrap replicates. CEAC indicates the probability that the intervention was cost effective at different values of willingness to pay for the additional improvement in the outcome.²⁸

In addition, four sensitivity analyses were used to examine how the results changed under different input assumptions. In the first two analyses, personnel wages and training costs were varied by 20% in either direction before recalculating the ICERs. The third sensitivity analysis used the summed weekly MVPA totals to estimate the ICER for each additional minute of MVPA per week. The fourth sensitivity analysis considered a different outcome measure for PA, using the summed weekly MVPA totals to estimate the ICER for changing one individual from insufficiently physically active to sufficiently physically active (≥150 min MVPA per week).²²

Patient and public involvement

Patients were not involved in the research question, study design or the conduction of the study.

RESULTS

Resource use and costs

Seventy-two participants were randomised; the group consisting of 54 females and 18 males, with an average age of 53±8 years (online supplementary file 2). A total of 72 participants completed their baseline and 3-month assessment, and 68 participants completed the 6-month assessment. For participants with missing data at 6-month follow-up (n = 2 in both groups), the last-observation-carried forward approach was adopted. The programme resources and cost per participant are described in table 1. Attendance at the education session

Table 1 Utilisation and cost of programme delivery for intervention and control groups

Item	Provider	Units	Time (hours)	Cost/hour \$A	Total \$A cost per participant
Intervention group					
Group sessions	Allied health professional	1	2.5	41.90	17.45
Phone call reminders	Intervention assistant	8	0.25	22.80	45.6
Intervention sessions	Allied health professional	8	0.5	41.90	167.6
Programme manual					3.5
Staff training					44.6
Total cost/participant					279
Control group					
Group sessions	Allied health professional	1	2.5	41.90	17.45
Programme manual					3.5
Total cost/participant					21

Table 2 Costs, changes in outcomes and incremental cost-effectiveness ratios at follow-up

Outcome	Cost\$/participant	Outcome	Incremental cost \$A	Incremental outcome	ICER
MVPA					
Control	21	23			
Intervention	279	41	258	18	\$A15/min MVPA per day
QALYs					
Control	21	-0.005			
Intervention	279	0.002	258	0.007	\$A36 857/QALY

ICER, incremental cost-effectiveness ratio; MVPA, moderate-to-vigorous physical activity; QALYs, quality-adjusted life-years.

was mandatory, with full attendance recorded for both groups. The mean group programme time was 34±6 min, and the mean total participant time spent in the intervention was 242±14 min. The mean cost per participant was \$279±\$10 for intervention versus \$21±\$3 for control. The main component costs of MI-CBT were intervention delivery and training.

Health outcomes

Table 2 presents the mean costs and the mean change in each outcome at follow-up for each group, and the corresponding ICERs. The ICER for MVPA was \$15 per each additional minute of MVPA per day. The difference in QALYs between intervention and control groups was 0.007 QALYs over the course of the follow-up period. The ICER for the intervention group in comparison with the control group was \$36 857 per QALY gained. Figures 1 and 2 illustrate the CEAC for each outcome derived from non-parametric bootstrapping replicates. For PA, given a willingness to pay of \$15 per additional minute of MVPA, the probability that the intervention was cost effective was 67% (figure 1). At a willingness to pay of \$37 000 per QALY gained, the probability that the intervention is cost effective was 52% (figure 2). If the decision maker was willing to pay \$40 000 per QALY, the probability of cost-effectiveness for the intervention was 70% (figure 2).

Figure 1 Cost-effectiveness acceptability curve showing the probability of the interventions being cost-effective in comparison to control for moderate-to-vigorous physical activity (MVPA).

In the sensitivity analyses, training and implementation costs were varied 20% in each direction, and the corresponding ICERs were recalculated (table 3). The varied ICERs for MVPA were found to range from \$11 to \$17 per each additional minute of MVPA per day, while the ICER for QALYs ranged from \$29 428 to AU\$44 285 per QALY gained. The third sensitivity analysis demonstrated an ICER of \$2.86 per additional minute of MVPA per week. The final sensitivity analysis found that the intervention group was 33% more likely than control to meet recommended PA guidelines at follow-up, with an ICER of \$781 per PA guideline attained.

DISCUSSION

This study examined the clinical and economic implications of a behaviour change intervention for changes in PA and QALYs for insufficiently physically active adults presenting to an ambulatory care clinic in a regional public hospital setting. Over the follow-up period, the MI-CBT intervention was significantly more effective than control in increasing PA and HrQoL. To the best of our knowledge, this is the first study to evaluate the cost-effectiveness and cost-utility of an integrated MI-CBT intervention for health-related behaviour change.

Telephone coaching can be cost effective for increasing PA with patients with or at risk of chronic disease.^{29 30}

Figure 2 Cost-effectiveness acceptability curve showing the probability of the interventions being cost-effective in comparison to control for quality-adjusted life-years (QALYs).

Table 3 Sensitivity analyses for costs, changes in outcomes and incremental cost-effectiveness ratios at follow-up

Outcome	Cost \$A/ participant	Outcome	Incremental cost \$A	Incremental outcome	ICER
MVPA+20% variation in cost					
Control	25	23			
Intervention	335	41	310	18	\$A17/min MVPA per day
QALY+20% variation in cost					
Control	25	-0.005			
Intervention	335	0.002	310	0.007	\$A44 285 QALY
MVPA-20% variation in cost					
Control	17	23			
Intervention	223	41	206	18	\$11/min MVPA per day
QALY-20% variation in cost					
Control	17	-0.005			
Intervention	223	0.002	206	0.007	\$A29 428 QALY
MVPA min per week					
Control	21	170			
Intervention	279	260	258	90	\$A2.86/min MVPA per week
PA guidelines attained					
Control	21	20/36 (55%)			
Intervention	279	31/36 (86%)	258	33%	\$A781/PA guideline achieved

ICER, incremental cost-effectiveness ratio; MVPA, moderate-to-vigorous physical activity; PA, physical activity; QALYs, quality-adjusted life-years.

However, little is known about the cost-effectiveness of adding preventive interventions to routine hospital care, where implementing an intervention requires an upfront investment of money. The total estimated cost of delivering the MI-CBT intervention was \$279 per person, resulting in the average attainment of 41±12 min of MVPA per day at follow-up. The per person cost was similar to the \$245 per person found in a recent primary care intervention for PA change,³¹ while both are considerably lower than costs of \$1756/person³² and \$1562/person³³ reported in other lifestyle interventions aiming at changes in PA.

The cost-effectiveness analysis for measured MVPA indicated a cost of \$15 per each additional minute of MVPA per day. Over the 6-month follow-up period, this can be translated to a total cost of \$9 per day, or \$63 per week to increase MVPA by 150 min. Sensitivity analysis indicated a one-time cost of \$2.86 per each additional minute of MVPA per week, or \$8.25 per week to increase MVPA to 150 min a week over the 6-month follow-up. The cost of \$8.25 per week to increase MVPA to 150 min is comparable to the \$4.99,³⁴ \$8.13³⁵ and \$10.19³⁶ per week found in other interventions aimed at increasing PA. Increases in PA result in decreased healthcare use, even in the short term, which result in net savings to society over time.³⁷ Determining the cost-effectiveness of integrating telephone coaching into routine care from a hospital perspective is dependent on the willingness-to-pay for each additional minute of MVPA. Interpreting the ICER of \$15

per additional minute of MVPA found here is difficult as there is no standard value for how much policy-makers are willing to pay per additional minute of MVPA.³⁴ The CEAC indicated a probability of 67% that \$15 per additional minute of MVPA was cost-effective (figure 1).

The intervention group was 33% more likely than the control group to undertake sufficient PA at follow-up. Sensitivity analysis demonstrated an ICER of \$781 for converting one insufficiently physically active adult to a sufficiently active state over the 6-month follow-up period. This value falls with the ranges of \$175 to \$1801,³⁸ and \$521 to \$5790³⁹ estimated in systematic reviews investigating the cost-effectiveness of PA interventions. Undertaking sufficient PA is strongly associated with decreased risk of chronic disease, morbidity and mortality,¹ as well as decreased healthcare expenditure over time.³⁷

In the cost-utility analysis, an incremental change in QALYs of 0.007 was demonstrated between the intervention and control groups, resulting in an ICER of \$36 857 per QALY gained. The ICER of \$36 857 per QALY gained is considerably smaller than the \$58 924 per QALY gained³² and the \$68 101 per QALY gained³³ found in similar intervention studies. Oksman *et al* recently reported an ICER of \$48 000 per QALY gained for a telephone-based health coaching intervention for chronic disease patients.⁴⁰ Direct comparison with the results from the aforementioned studies is challenging because different cost perspectives were considered in

these analyses. Nonetheless, the ICER of \$36 857 per QALY falls under the commonly used threshold of \$50 000 per QALY gained proposed for medical treatments and procedures.⁴¹ Sensitivity analysis demonstrated that costs for the Healthy4U study are only mildly sensitive to typical variation in the cost input values. A variation in hourly wage costs had the largest impact on the ICERs; however, the ICERs remained well below the threshold of \$50 000 per QALY gained at all imputed values. The CEAC provided a probability of 52% that the intervention was cost effective at a willingness to pay of \$37 000 per QALY gained (figure 2).

Behaviour change interventions are typically used to modify specific lifestyle factors known to predispose individuals to increased risk of chronic disease over the longer term.⁴² The long-term impact of such interventions on overall quality of life is less established. While the follow-up period in this study was too short for the mediating effect of PA on broader health outcomes to become fully apparent,⁴² the relatively small observed change in QALYs over 6 months was a combination of the -0.005 fall in the control group and the 0.002 increase in the intervention group. This finding supports the suggestion that an important impact of behavioural interventions on quality-of-life over the longer term might be to attenuate expected declines in HrQoL.⁴³

This study was unique in that we enrolled participants from an ambulatory care clinic in a public hospital, integrating preventative health into secondary care. It is important to note that this intervention was carried out in addition to standard care, not as a substitution, and as such an economic evaluation costed from a hospital perspective offers healthcare providers an estimate of the costs and effects of adding a preventative health intervention to clinical care. This study is one of the few economic evaluations of telephone coaching carried out in real-life settings using an RCT design. The addition of preventive health measures is likely to cost hospitals more; however, these preventive measures might be worthwhile due to the substantial health benefits that they confer, relative to their cost.³⁷ Nevertheless, implementation remains a challenge for hospitals,¹⁵ and it will be important to engage with key stakeholders, especially clinic leaders, to identify specific patients who can benefit from telephone coaching.⁴⁴ Health services with high implementation rates of telephone coaching have used multicomponent strategies to engage staff, as they were the most important source of referrals.⁴⁴ Investigating hospital clinicians' practice and beliefs around preventive health can also facilitate the development of pathways to increase preventive health practice in the hospital setting.⁴⁵

The greatest cost of the intervention was in the delivery, due to the fact that it was delivered individually by trained personnel. These costs could be reduced by decreasing the number of intervention sessions, with a recent meta-analysis indicating that five sessions of MI-CBT are significantly effective for PA change.⁴⁶ Costs could potentially be reduced by incorporating digital technology into the

intervention to decrease the time spent by trained professionals in 1:1 sessions. Innovations in digital technologies can assist individuals with health behaviour change,⁴⁷ which can potentially reduce healthcare expenditure.^{47 48} The long-term cost-effectiveness of digital health technologies has not been established.⁴⁷

The use of objectively measured PA at all time points was a considerable strength of the study. Objective measures offer more precise estimates of activity intensity while removing many of the issues associated with participant recall and response bias.⁴⁹ Individuals have been demonstrated to overestimate their PA levels via self-report.³⁸ Overestimation of PA can result in inaccurate estimations of both effectiveness and the cost-effectiveness of interventions.⁵⁰ Using objectively measured changes in PA and the collection of full cost data for all participants strengthens our findings.⁴⁹

This study has a number of limitations. A noteworthy limitation of this study is the restricted perspective used for the economic evaluation. Using a single hospital perspective might have led to the exclusion of important costs and benefits from a societal perspective, including healthcare utilisation and changes in productivity.⁵¹ Economic analyses from a societal perspective offer the most comprehensive evidence from which to base decisions.⁵¹ However, due to the relatively short follow-up time of this study, it was not feasible to undertake this method. Significant effects on overall healthcare utilisation or productivity loss were not expected over the time-frame of this study.⁵² As the intervention was delivered using the telephone, the intervention required relatively small amounts of participant time. With this in mind, the participant opportunity costs were expected to be small and therefore not included in the analyses. Additionally, the study participants included 54 females and 18 males, which might limit the generalisability of the current findings to different populations.

To the best of our knowledge, this is the first study comparing the cost-effectiveness and cost-utility of an integrated MI-CBT intervention for health behaviour change among community-dwelling adults presenting to a secondary care clinic in a regional public hospital setting. Considering the large group of people who might benefit from such an intervention (ie, approximately 50% of adults aged 18–69 years in Australia who are currently insufficiently physically active), the widespread adoption and implementation of MI-CBT to increase PA could have important economic implications. However, further research with a more comprehensive economic analysis is needed to investigate whether the long-term benefits of MI-CBT might justify this type of investment.

CONCLUSION

The Healthy4U programme is a relatively low-cost strategy for increasing PA among insufficiently physically active adults presenting to an ambulatory care hospital clinic. The intervention increased measured PA and quality

of life at low costs, with positive effects maintained out to 6 months. By increasing PA and quality of life at low costs, integrating telephone coaching into secondary hospital care offers a potentially cost-effective investment to produce better public health outcomes. The results are however grounded on a short-term follow-up and a restricted economic perspective, and more evidence is needed to explore the potential long-term economic impact of the intervention from a broader healthcare perspective.

Author affiliations

¹Health Promotion, Bendigo Health - Bendigo Hospital Campus, Bendigo, Victoria, Australia

²La Trobe Holsworth Research Initiative, La Trobe University, Bendigo, Victoria, Australia

³School of Psychology and Public Health, La Trobe University, Bundoora, Victoria, Australia

Twitter Stephen Barrett @barrettstephen1

Contributors SBA, MK, SBe and PO'H conceived the project and assisted with the protocol design. SBA managed the trial including recruitment and data collection, coordinated the intervention program, performed statistical analysis and drafted the manuscript. SBA and SBe interpreted the data and drafted the manuscript. All authors read, edited and approved the final manuscript as submitted.

Funding The authors have not declared a specific grant for this research from any funding agency in the public, commercial or not-for-profit sectors.

Competing interests None declared.

Patient consent for publication Not required.

Ethics approval The study was approved by the Research Ethics Committees of Bendigo Health Care group (approved 16 September 2016; reference number LNR/16/BHCG/42) and La Trobe University College of Science Health and Engineering Human Ethics Sub-Committee (approved 3 October 2016).

Provenance and peer review Not commissioned; externally peer reviewed.

Data availability statement Data are available upon reasonable request.

Open access This is an open access article distributed in accordance with the Creative Commons Attribution Non Commercial (CC BY-NC 4.0) license, which permits others to distribute, remix, adapt, build upon this work non-commercially, and license their derivative works on different terms, provided the original work is properly cited, appropriate credit is given, any changes made indicated, and the use is non-commercial. See: <http://creativecommons.org/licenses/by-nc/4.0/>.

ORCID iD

Stephen Barrett <http://orcid.org/0000-0003-2505-733X>

REFERENCES

- Warburton DER, Nicol CW, Bredin SS. Health benefits of physical activity: the evidence. *Can Med Assoc J* 2006;174:801-9.
- Khan KM, Thompson AM, Blair SN, et al. Sport and exercise as contributors to the health of nations. *Lancet* 2012;380:59-64.
- World Health Organization. *Global health risks: mortality and burden of disease attributable to selected major risks*. World Health Organization, 2009. <http://www.who.int/iris/handle/10665/44203>
- Ding D, Lawson KD, Kolbe-Alexander TL, et al. The economic burden of physical inactivity: a global analysis of major non-communicable diseases. *Lancet* 2016;388:1311-24.
- Sparling PBe et al. Promoting physical activity: the new imperative for public health. *Health Educ Res* 2000;15:367-76.
- Britt HC, Harrison CM, Miller GC, et al. Prevalence and patterns of multimorbidity in Australia. *Med J Aust* 2008;189:72-7.
- Steiner CA, Friedman B. Hospital utilization, costs, and mortality for adults with multiple chronic conditions, nationwide inpatient sample, 2009. *Prev Chronic Dis* 2013;10:E62.
- Gate L, Warren-Gash C, Clarke A, et al. Promoting lifestyle behaviour change and well-being in hospital patients: a pilot study of an evidence-based psychological intervention. *J Public Health* 2016;38:e292-300.
- Lawson PJ, Flocke SA. Teachable moments for health behavior change: a concept analysis. *Patient Educ Couns* 2009;76:25-30.
- Allender S, Hutchinson L, Foster C. Life-change events and participation in physical activity: a systematic review. *Health Promot Int* 2008;23:160-72.
- Foster C, Hillsdon M, Thorogood M, et al. Interventions for promoting physical activity. *Cochrane Database Syst Rev* 2005;18.
- Stephens J, Allen J. Mobile phone interventions to increase physical activity and reduce weight: a systematic review. *J Cardiovasc Nurs* 2013;28:320-9.
- Dennis SM, Harris M, Lloyd J, et al. Do people with existing chronic conditions benefit from telephone coaching? a rapid review. *Aust Health Rev* 2013;37:381-8.
- Kivelä K, Elo S, Kyngäs H, et al. The effects of health coaching on adult patients with chronic diseases: a systematic review. *Patient Educ Couns* 2014;97:147-57.
- Eakin EG, Lawler SP, Vandelanotte C, et al. Telephone interventions for physical activity and dietary behavior change. *Am J Prev Med* 2007;32:419-34.
- O'Hara BJ, Phongsavan P, Venugopal K, et al. Effectiveness of Australia's get healthy information and coaching Service®: translational research with population wide impact. *Prev Med* 2012;55:292-8.
- Barrett S, Begg S, O'Halloran P, et al. Integrated motivational interviewing and cognitive behaviour therapy can increase physical activity and improve health of adult ambulatory care patients in a regional Hospital: the Healthy4U randomised controlled trial. *BMC Public Health* 2018;18:1166.
- Husereau D, Drummond M, Petrou S, et al. Consolidated health economic evaluation reporting standards (cheers) statement. *BMJ* 2013;346.
- Vansteenkiste M, Sheldon KM. There's nothing more practical than a good theory: integrating motivational interviewing and self-determination theory. *Br J Clin Psychol* 2006;45:63-82.
- Freedson PS, Melanson E, Sirard J. Calibration of the computer science and applications, Inc. accelerometer. *Medicine & Science in Sports & Exercise* 1998;30:777-81.
- Olds T, Ferrar KE, Gomersall SR, et al. The elasticity of time. *Health Educ Behav* 2012;39:732-6.
- Nelson ME, Rejeski WJ, Blair SN, et al. Physical activity and public health in older adults: recommendation from the American College of sports medicine and the American heart association. *Circulation* 2007;116:1094-105.
- Ware Jr JE, Kosinski M, Keller SD. A 12-Item short-form health survey: construction of scales and preliminary tests of reliability and validity. *Med Care* 1996;34:220-33.
- Brazier J, Roberts J, Deverill M. The estimation of a preference-based measure of health from the SF-36. *J Health Econ* 2002;21:271-92.
- Drummond M, Sculpher M, Torrance G, et al. *Methods for the economic evaluation of health care programmes*. 3rd edn. New York: Oxford University Press, 2005.
- Tan SS, Rutten FFH, van Ineveld BM, et al. Comparing methodologies for the cost estimation of hospital services. *Eur J Health Econ* 2009;10:39-45.
- Glick HA, Doshi JA, Sonnad SS, et al. *Economic evaluation in clinical trials*. Oxford: Oxford University Press, 2014.
- Appleby J, Devlin N, Parkin D. NICE's cost effectiveness threshold. *BMJ* 2007;335:358-9.
- Hutchison AJ, Breckon JD. A review of telephone coaching services for people with long-term conditions. *J Telemed Telecare* 2011;17:451-8.
- Varney JE, Liew D, Weiland TJ, et al. The cost-effectiveness of hospital-based telephone coaching for people with type 2 diabetes: a 10 year modelling analysis. *BMC Health Serv Res* 2016;16:521.
- Ewald B, Stacey F, Johnson N, et al. Physical activity coaching by Australian Exercise Physiologists is cost effective for patients referred from general practice. *Aust N Z J Public Health* 2018;42:12-15.
- Groessl EJ, Kaplan RM, Castro Sweet CM, et al. Cost-effectiveness of the life physical activity intervention for older adults at increased risk for mobility disability. *GERONA* 2016;71:656-62.
- Groessl EJ, Kaplan RM, Blair SN, et al. A cost analysis of a physical activity intervention for older adults. *J Phys Act Health* 2009;6:767-74.
- Johnson TJ, E Schoeny M, Fogg L, et al. The cost of increasing physical activity and maintaining weight for midlife sedentary African American women. *Value Health* 2016;19:20-7.

- 35 Elley CR, Garrett S, Rose SB, *et al.* Cost-effectiveness of exercise on prescription with telephone support among women in general practice over 2 years. *Br J Sports Med* 2011;45:1223–9.
- 36 Seivick MA, Napolitano MA, Papandonatos GD, *et al.* Cost-effectiveness of alternative approaches for motivating activity in sedentary adults: results of Project STRIDE. *Prev Med* 2007;45:54–61.
- 37 Martinson BC, Crain AL, Pronk NP, *et al.* Changes in physical activity and short-term changes in health care charges: a prospective cohort study of older adults. *Prev Med* 2003;37:319–26.
- 38 GC V, Wilson ECF, Suhrcke M, *et al.* Are brief interventions to increase physical activity cost-effective? A systematic review. *Br J Sports Med* 2016;50:408–17.
- 39 Garrett S, Elley CR, Rose SB, *et al.* Are physical activity interventions in primary care and the community cost-effective? A systematic review of the evidence. *Br J Gen Pract* 2011;61:e125–33.
- 40 Oksman E, Linna M, Hörhammer I, *et al.* Cost-effectiveness analysis for a tele-based health coaching program for chronic disease in primary care. *BMC Health Serv Res* 2017;17:138.
- 41 Hirth RA, Chernew ME, Miller E, *et al.* Willingness to pay for a quality-adjusted life year: in search of a standard. *Med Decis Making* 2000;20:332–42.
- 42 Goyder E, Hind D, Breckon J, *et al.* A randomised controlled trial and cost-effectiveness evaluation of ‘booster’ interventions to sustain increases in physical activity in middle-aged adults in deprived urban neighbourhoods. *Health Technol Assess* 2014;18:1.
- 43 Cutler DM. Behavioral health interventions: what works and why. In: *Critical perspectives on racial and ethnic differences in health in late life*, 2004: 643.
- 44 Damschroder LJ, Reardon CM, Sperber N, *et al.* Implementation evaluation of the telephone lifestyle coaching (TLC) program: organizational factors associated with successful implementation. *Transl Behav Med* 2017;7:233–41.
- 45 Barrett S, Begg S, Sloane A, *et al.* Surgeons and preventive health: a mixed methods study of current practice, beliefs and attitudes influencing health promotion activities amongst public hospital surgeons. *BMC Health Serv Res* 2019;19:358.
- 46 Barrett S, Begg S, O'Halloran P, *et al.* Integrated motivational interviewing and cognitive behaviour therapy for lifestyle mediators of overweight and obesity in community-dwelling adults: a systematic review and meta-analyses. *BMC Public Health* 2018;18:1160.
- 47 Sanyal C, Stolee P, Juzwishin D, *et al.* Economic evaluations of eHealth technologies: a systematic review. *PLoS One* 2018;13:e0198112.
- 48 Jiang X, Ming W-K, You JHS. The cost-effectiveness of digital health interventions on the management of cardiovascular diseases: systematic review. *J Med Internet Res* 2019;21:e13166.
- 49 Prince SA, Adamo KB, Hamel M, *et al.* A comparison of direct versus self-report measures for assessing physical activity in adults: a systematic review. *Int J Behav Nutr Phys Act* 2008;5.
- 50 Bize R, Johnson JA, Plotnikoff RC. Physical activity level and health-related quality of life in the general adult population: a systematic review. *Prev Med* 2007;45:401–15.
- 51 Jönsson B. Ten arguments for a societal perspective in the economic evaluation of medical innovations. *Eur J Health Econ* 2009;10:357–9.
- 52 van Keulen HM, Bosmans JE, van Tulder MW, *et al.* Cost-effectiveness of tailored print communication, telephone motivational interviewing, and a combination of the two: results of an economic evaluation alongside the Vitalum randomized controlled trial. *Int J Behav Nutr Phys Act* 2010;7.