Correction: Predictors of physical activity among older adults in Germany: a nationwide cohort study

Manz K, Mensink GBM, Jordan S, *et al.* Predictors of physical activity among older adults in Germany: a nationwide cohort study *BMJ Open* 2018;8:e021940. doi: 10.1136/bmjopen-2018-021940

This article was previously published with an error in table 2.

The columns with variable categories 'yes'/'no' were swapped. Below is the updated table 2.

Bivariate associations between aerobic PA ≥ 1 day/week and potential predictor variables

		Aerobic PA ≥1 day/week				P value*
		No		Yes		
		%	95% CI	%	95% CI	
1	Sex					
	Women (n=622)	48.7	(44.7 to 52.7)	51.3	(47.3 to 55.3)	0.158
	Men (n=562)	44.7	(40.1 to 49.4)	55.3	(50.6 to 59.9)	
	Age group					
	50–60 years (n=713)	38.8	(35.1 to 42.8)	61.2	(57.2 to 64.9)	<0.001
	61–78 years (n=471)	58.8	(54.1 to 63.4)	41.2	(36.6 to 45.9)	
	Socioeconomic status					
	Low (n=151)	70.2	(62.0 to 77.3)	29.8	(22.7 to 38.0)	<0.001
	Middle (n=712)	47.9	(43.8 to 52.0)	52.1	(48.0 to 56.2)	
	High (n=307)	31.9	(27.0 to 37.3)	68.1	(62.7 to 73.0)	
	Chronic disease					
	No (n=869)	44.1	(40.5 to 47.7)	55.9	(52.3 to 59.5)	0.006
	Yes (n=312)	53.8	(47.6 to 60.0)	46.2	(40.0 to 52.4)	
	Obesity					
	Yes (n=297)	53.5	(47.4 to 59.5)	46.5	(40.5 to 52.6)	0.005
	No (n=884)	44.6	(41.2 to 48.0)	55.4	(52.0 to 58.8)	
2	Living with a spouse					
	No (n=216)	57.4	(51.1 to 63.4)	42.6	(36.6 to 48.9)	<0.001
	Yes (n=952)	44.1	(40.5 to 47.8)	55.9	(52.2 to 59.5)	
	Social support					
	Low (n=100)	65.0	(55.4 to 73.5)	35.0	(26.5 to 44.6)	<0.001
	High (n=1063)	44.9	(41.6 to 48.2)	55.1	(51.8 to 58.4)	
	Leisure time physical activity					
	Every week (n=784)	38.6	(34.8 to 42.6)	61.4	(57.4 to 65.2)	<0.001
	Not every week (n=374)	62.8	(57.6 to 67.8)	37.2	(32.2 to 42.4)	
3	At least one health programme					
	Yes (n=124)	34.7	(26.7 to 43.6)	65.3	(56.4 to 73.3)	0.005
	No (n=1060)	48.2	(44.8 to 51.7)	51.8	(48.3 to 55.2)	
	Smoking status					
	Smoker (n=188)	49.5	(41.7 to 57.3)	50.5	(42.7 to 58.3)	0.406

		Aerobic PA ≥1 day/week				P value*
		No		Yes		
		%	95% CI	%	95% CI	
4	Non-smoker (n=985)	46.0	(42.5 to 49.5)	54.0	(50.5 to 57.5)	0.873
	Residential area size					
	Rural (n=249)	48.6	(42.1 to 55.1)	51.4	(44.9 to 57.9)	
	Small-sized city (n=247)	48.2	(39.4 to 57.1)	51.8	(42.9 to 60.6)	
	Medium-sized city (n=318)	45.9	(39.6 to 52.4)	54.1	(47.6 to 60.4)	
	Metropolitan city (n=370)	45.4	(40.5 to 50.4)	54.6	(49.6 to 59.5)	
	Satisfaction with residential area				0.711	
	Not satisfied (n=308)	47.1	(42.0 to 52.2)	52.9	(47.8 to 58.0)	
	Satisfied (n=855)	46.0	(42.2 to 49.7)	54.0	(50.3 to 57.8)	

1: Individual factors; 2: Interpersonal factors; 3: Behavioural factors; 4: Environmental factors.

*Pearson's chi-squared test with Rao-Scott correction.

PA, physical activity.

Open access This is an open access article distributed in accordance with the Creative Commons Attribution Non Commercial (CC BY-NC 4.0) license, which permits others to distribute, remix, adapt, build upon this work non-commercially, and license their derivative works on different terms, provided the original work is properly cited, appropriate credit is given, any changes made indicated, and the use is non-commercial. See: <http://creativecommons.org/licenses/by-nc/4.0/>.

© Author(s) (or their employer(s)) 2019. Re-use permitted under CC BY-NC. No commercial re-use. See rights and permissions. Published by BMJ.

BMJ Open 2019;9:e021940corr1. doi:10.1136/bmjopen-2018-021940corr1