

Table 1. Odds for receiving an SSRI prescription each year between 2011 and 2015

| Variable | Coefficient | Standard Error | p-value | Odds ratio | 95% Confidence Interval | |
|--|--------------------|-----------------------|----------------|-------------------|--------------------------------|------|
| Sex (male = ref) | | | | | | |
| Female | 0.1699 | 0.0576 | 0.0032 | 1.19 | 1.06 | 1.33 |
| Age (65+ = ref) | | | | | | |
| 19–44 | -0.1077 | 0.0767 | 0.1600 | 0.90 | 0.77 | 1.04 |
| 45–64 | 0.2582 | 0.0700 | p < .001 | 1.29 | 1.13 | 1.49 |
| Disorder | | | | | | |
| Anxiety (no anxiety = ref) | 0.0333 | 0.0771 | 0.6656 | 1.03 | 0.89 | 1.20 |
| Depression (no depression = ref)) | -0.1207 | 0.0698 | 0.0838 | 0.89 | 0.77 | 1.02 |
| Somatic disorder (no somatic disorder = ref) | 0.1537 | 0.0849 | 0.0703 | 1.17 | 0.99 | 1.38 |
| | | | | | | |
| Practice variance | 4.711 | 0.626 | | | | |
| ICC | 0.59 | | | | | |
| Constant | -3.6397 | 0.2066 | | | | |

Table 2. Odds for receiving a TCA prescription during each year between 2011 and 2015

| Variable | Coefficient | Standard Error | p-value | Odds ratio | 95% Confidence Interval | |
|--|--------------------|-----------------------|----------------|-------------------|--------------------------------|------|
| Sex (male = ref) | | | | | | |
| Female | 0.2030 | 0.0917 | 0.0268 | 1.23 | 1.02 | 1.47 |
| Age (65+ = ref) | | | | | | |
| 19–44 | -0.6821 | 0.1300 | p < .001 | 0.51 | 0.39 | 0.65 |
| 45–64 | 0.0393 | 0.0915 | 0.6673 | 1.04 | 0.87 | 1.24 |
| Disorder | | | | | | |
| Anxiety (no anxiety = ref) | 0.2098 | 0.1387 | 0.1303 | 1.23 | 0.94 | 1.62 |
| Depression (no depression = ref) | 0.3797 | 0.1137 | p < .001 | 1.46 | 1.17 | 1.83 |
| Somatic disorder (no somatic disorder = ref) | -0.0631 | 0.1098 | 0.5654 | 0.94 | 0.76 | 1.16 |
| | | | | | | |
| Practice variance | 2.763 | 0.4156 | | | | |
| ICC | 0.46 | | | | | |
| Constant | -3.6022 | 0.1936 | | | | |