

# BMJ Open Impact of maternal smoking on early childhood health: a retrospective cohort linked dataset analysis of 697 003 children born in Scotland 1997–2009

Richard Lawder,<sup>1</sup> Bruce Whyte,<sup>2</sup> Rachael Wood,<sup>1,3</sup> Colin Fischbacher,<sup>1</sup> David Michael Tappin<sup>4</sup>

**To cite:** Lawder R, Whyte B, Wood R, *et al.* Impact of maternal smoking on early childhood health: a retrospective cohort linked dataset analysis of 697 003 children born in Scotland 1997–2009. *BMJ Open* 2019;**9**:e023213. doi:10.1136/bmjopen-2018-023213

► Prepublication history for this paper is available online. To view these files, please visit the journal online (<http://dx.doi.org/10.1136/bmjopen-2018-023213>).

Received 11 April 2018  
Revised 22 November 2018  
Accepted 23 November 2018



© Author(s) (or their employer(s)) 2019. Re-use permitted under CC BY-NC. No commercial re-use. See rights and permissions. Published by BMJ.

<sup>1</sup>Information Services Division, NHS National Services Scotland, Edinburgh, UK

<sup>2</sup>Glasgow Centre for Population Health, University of Glasgow, Glasgow, UK

<sup>3</sup>Child Life and Health, University of Edinburgh, Edinburgh, UK

<sup>4</sup>Scottish Cot Death Trust, 5th Floor, West Glasgow Ambulatory Care Hospital, University of Glasgow, Glasgow, UK

## Correspondence to

Dr David Michael Tappin;  
[david.tappin@glasgow.ac.uk](mailto:david.tappin@glasgow.ac.uk)

## ABSTRACT

**Objective** Smoking during pregnancy is associated with adverse health impacts on mother and child. We used a large linked Scottish dataset to produce contemporary estimates of the impact on child health, particularly hospitalisation.

**Design** Retrospective cohort study linking birth, death, maternity, infant health, child health surveillance and admission records. We examined the association between smoking status at maternity booking and pregnancy outcomes, hospital admission and death during the first 5 years of life. Models were adjusted for maternal age, socioeconomic status, infant feeding, country of birth, sex, parity and delivery mode. We calculated population attributable fraction (PAF) for each outcome.

**Setting** Scotland, UK.

**Participants** Singleton births between 1997 and 2009 (n=697 003) followed to March 2012.

**Results** 332 386 children had at least one admission by 31 March 2012. There were 56 588 born small for gestational age, 40 492 prematurely and 1074 postneonatal deaths. Within the first 5 years of life, 56 615 children had at least one admission for acute respiratory infections, 24 088 for bronchiolitis and 7549 for asthma. Maternal smoking significantly increased admission for acute respiratory infections (adjusted HR 1.29, 95% CI 1.25 to 1.34, PAF 6.7%) and bronchiolitis (HR 1.43, 95% CI 1.38 to 1.48 under 1 year, PAF 10.1%), asthma (HR 1.29, 95% CI 1.22 to 1.37 age 1–5 years, PAF 7.1%) and bacterial meningitis (HR 1.49, 95% CI 1.30 to 1.71, PAF 11.8%) age 0–5 years. Neonatal mortality (adjusted OR 1.32, 95% CI 1.17 to 1.49, PAF 6.7%), postneonatal mortality (OR 2.18, 95% CI 1.87 to 2.53, PAF 22.3%), small for gestational age (OR 2.67, 95% CI 2.62 to 2.73, PAF 27.5%) and prematurity (OR 1.41, 95% CI 1.37 to 1.44, PAF 8.8%) were higher among the offspring of smokers.

**Conclusion** Smoking during pregnancy causes significant ill health and death among children born in Scotland. These findings support continued investment to reduce smoking among women before, during and after pregnancy as 50% of women will go on to have further children.

## Strengths and limitations of this study

- Extensive dataset linkage allowed many outcomes to be assessed for each infant and inclusion of a control outcome to assess residual confounding.
- Use of smoking data collected only during pregnancy and lack of smoking data for 12% of the dataset.
- For 20% of records, maternal occupation was missing but balanced against this was the inclusion of three different measures of socioeconomic status (including maternal occupation, paternal occupation and deprivation Scottish Index of Multiple Deprivation score for area of maternal residence).
- Hospital admissions only were included and it is known that a large part of childhood morbidity is dealt with by Primary Care.

## INTRODUCTION

### Background

A total of 350 UK and 1100 US stillbirths each year are attributable to smoking during pregnancy<sup>1</sup> as are 200 UK and 900 US infant deaths.<sup>2–4</sup> Maternal smoking also increases a child's chance of being born prematurely with low birth weight,<sup>5</sup> and having asthma,<sup>6</sup> attention deficit disorder<sup>7</sup> and learning difficulties,<sup>8</sup> which if causally related all add substantial costs to healthcare.<sup>9</sup> Pregnancy is also an opportunity to help most women quit smoking permanently before their own health is irreversibly compromised. Cessation during pregnancy if permanent reduces the likelihood of offspring becoming smokers,<sup>10</sup> stopping mothers by example passing on their smoking habit to their children.

A fall in reported smoking during pregnancy has been reported in Scotland<sup>11</sup> (20.8% 2007, 15.5% 2015). In 2000, specific cessation services<sup>12</sup> for pregnant women were established, supported by a National Institute for Health and Care Excellence guideline.<sup>13</sup> However, self-reported smoking during

and after pregnancy remains common. Twenty-five per cent of biochemically verified smokers self-report as non-smokers,<sup>14</sup> and this proportion may be increasing as pressure to quit intensifies. Only 10% of pregnant smokers use cessation services in Scotland and as few as 3% quit during pregnancy.<sup>15</sup> A recent report from the UK is more encouraging with on average 50% cessation during pregnancy.<sup>16</sup>

### Objectives

We used a large linked Scottish dataset<sup>17</sup> to produce contemporary estimates of the impact of maternal smoking measured in early pregnancy on pregnancy, infant and early child health outcomes. We focused on infant deaths and admissions to hospital for a range of conditions to assess the proportion of events attributable to smoking during pregnancy.

## METHODS

### Study design, setting, participants, data sources

This was a retrospective cohort study of singleton births in Scotland between 1997 and 2009 using anonymised extracts of linked administrative data provided by the Information Services Division, National Health Service National Services Scotland. Approval for the project design and confidentiality of patient data was provided by the Privacy Advisory Committee of NHS National Services Scotland, a body set up to ensure the appropriate use of patient identifiable information.<sup>18</sup> Further ethical permission was not required. All 731 595 live birth records in Scotland between the calendar years 1997 and 2009 inclusive were linked in two phases by probabilistic matching techniques including the use of the Community Health Index (CHI), a unique identifier developed for health records in Scotland.<sup>19</sup> Phase 1 comprised linkage of births, deaths, migration, maternity, infant health and child health surveillance review records,<sup>20</sup> which was extended in phase 2 to include all hospital admissions (ie, emergency admission and elective) from birth until death or up to March 2012. Thus, follow-up information on each child in the cohort was available for at least 2.25 years (for all children) and up to 15 years (depending on their birth year). The original cohort of 731 595 births was reduced to 697 003 as a result of excluding multiple births, babies of non-Scottish residents and duplicate linkage records.

### Outcome variables and bias

The outcomes examined were neonatal and postneonatal mortality, pregnancy outcomes (infant small for gestational age and premature births) and hospital discharge up to the fifth birthday with a primary diagnosis from a list of conditions selected from a review of literature. Conditions of interest were asthma, bronchiolitis, bacterial meningitis, acute upper and lower respiratory infections, acute and chronic otitis media and mastoiditis and bronchopulmonary dysplasia. Emergency and planned

admissions (with day cases) were included. Admissions for long bone fracture, not thought to be associated with maternal smoking, were used as a control outcome to assess if confounding had been satisfactorily controlled in the multivariate analysis.

### Definition of smoking during pregnancy

Smoking behaviour in pregnancy is collected at a woman's first antenatal booking appointment which usually takes place within the first 3 months of pregnancy. These booking appointments take place either at hospital or in the community and are recorded on the Scottish Woman-Held Maternity Record. The smoking at booking information is subsequently transcribed onto the woman's Scottish Morbidity Record 02 which is submitted by the maternity units following delivery. The smoking behaviour is classified into either 'never smoked', 'current smoker', 'former smoker' or 'unknown/missing'.

### Statistical methods and quantitative variables

Descriptive and univariate analyses were used to identify variables associated with smoking during pregnancy and perinatal, mortality and hospitalisation outcomes. Cox regression was used to quantify the independent contribution of maternal smoking to hospitalisation outcomes, based on survival from birth to the earliest of first hospitalisation, death, migration, the fifth birthday or the end of the observation period (March 2012). Logistic regression was used for perinatal and mortality outcomes. The models included variables identified from directed acyclic graphs as potentially biasing (confounding) the association between smoking and the relevant outcomes. Adjustment was made for maternal age, maternal and paternal socioeconomic status, infant feeding at 6–8 weeks, country of birth, infant sex, parity, mode of delivery and quintile of the Scottish Index of Multiple Deprivation (SIMD). Each Cox regression model was tested for proportionality using inspection of log–log plots. Additional analysis was conducted to assess the risk of hospitalisation for long bone fractures (a condition not causally associated with maternal smoking), to test the adequacy of adjustment for socioeconomic and demographic confounders included in the linked dataset. Statistical modelling was carried out using Stata V.11; StataCorp LP.

Population attributable fractions (PAFs) were used to quantify the number of outcome events that would have been avoided if no women had smoked, using the formula:

$$PAF = \frac{P_c(RR-1)}{RR}$$

where RR is the relative risk and  $P_c$  is the proportion of cases exposed to the risk factor.

### Patient or public involvement

No patient or public involvement.

## RESULTS

### Participants

Of the 697003 singletons born between 1997 and 2009 included in the analysis (table 1), 63% were born by spontaneous vaginal delivery, 8% had teenage mothers, 16% were born to single parents or parents living apart, 44% were born to first-time mothers and 25% were residents in the most socioeconomically deprived quintile. Twenty-three per cent of infants had mothers who smoked (table 1). A total of 332386 (48%) had at least one hospital admission by 31 March 2012 (figure 1).

### Descriptive data

As previously reported,<sup>17</sup> there was a greater relative risk of hospital admission among infants resident in more deprived areas, of fathers and mothers with a semiroutine/routine occupation, of babies bottle fed or combination fed (bottle and breast), among infants with siblings, and those born by Caesarean delivery. Conversely, risk of infant admission decreased with increasing maternal age, among female infants and if a mother's country of birth was non-British.<sup>17</sup> During the study period, 87877 (12.6%) infants were hospitalised at least once for one of the selected conditions in the first 5 years (figure 1). At the first recorded hospital event for any of the selected conditions, 43% were under 1 year of age, and 57% were between first and fifth birthday. Most of these first hospital events were 'emergency admissions' (91%), especially among infants less than 1 year of age at admission (99%).

Younger unmarried pregnant women and women from deprived postcode areas and socioeconomic circumstances were more likely to be current smokers, as were British born compared with non-British born pregnant women (table 1). Smoking varied little by mode of delivery or number of previous children, however, babies who were bottle fed, or whose mother's religion was Christian or Jewish were more likely to have mothers who were current smokers at maternity booking (table 1).

### Outcome data

The crude rate of hospitalisation (any cause) per 1000 person-years was 141 among infants born to mothers who smoked, 108 for those who did not smoke, 119 for former smokers and 109 for those with missing smoking status (data not shown in tables or figure).

### Main results

#### Multivariate analyses

We start with the most devastating outcomes.

#### Perinatal and mortality outcomes and smoking status at first booking

Among babies whose mothers were current smokers at hospital booking, the odds of neonatal mortality (in first month after birth) were a third higher than among non-smokers after taking account of confounding variables (OR 1.32, 95% CI 1.17 to 1.49, PAF 6.7% table 2). The odds of postneonatal infant mortality (between 1 month and 1 year of age) were 2.18 times higher if the

mother smoked (PAF 22.3%). The odds of being born small for gestational age was 2.67 times higher for current smokers (PAF 27.5%) while the odds of premature delivery were also significantly increased (OR 1.41, 95% CI 1.37 to 1.44, PAF 8.8%).

#### Rates of hospitalisation for acute respiratory infections, bronchiolitis and asthma

Because of non-proportional hazards for these conditions over the whole period of follow-up, hazard ratios (HRs) are presented separately for the first year of the life and the subsequent 4 years (table 3). During the first year of life, the adjusted HR for hospital admission for acute respiratory infections was increased by 29% (PAF 6.7%), by 43% (PAF 10.1%) for bronchiolitis and more than doubled for asthma among infants of mothers who smoked. For the period from the first birthday to the age of 5, there were more modest increases in hazards for asthma (adjusted HR=1.29, PAF=7%) and for bronchiolitis (adjusted HR=1.16, PAF=4%) but no increase in hazard of acute respiratory infections.

#### Rates of hospitalisation for meningitis, ear infections and chronic lung disease

For children of mothers who were current smokers during pregnancy, the rate of hospitalisation in the first 5 years of life was substantially raised compared with non-smokers for bronchopulmonary dysplasia (50% increase, PAF 10.1%) and for bacterial meningitis (49% increase, PAF 11.8%, table 4). There was only a small increase in the rate of hospitalisation with acute and chronic otitis media and mastoiditis (7% increase, PAF 1.7%).

#### Other analyses

Hospital admissions related to long bone fractures were analysed as a control group. There were 5757 children admitted for long bone fractures from the cohort during the study period. After adjustment for other factors (maternal age, maternal and paternal socioeconomic status, infant feeding at 6–8 weeks, country of birth, infant sex, parity, mode of delivery and quintile of the SIMD), a mother being a smoker at maternity booking was not a significant predictor of hospital admission for long bone fractures.

## DISCUSSION

This study of Scottish births (1997–2009) provides more detailed evidence about the association between maternal smoking in pregnancy and childhood hospitalisation as well as birth conditions which can lead to lifelong ill health and devastating outcomes such as meningitis. This study also estimates the proportion of babies born prematurely, small for gestational age and infant deaths that were attributable to maternal smoking during pregnancy. These associations remain significant after adjustment for a range of socioeconomic factors.

**Table 1** Cohort characteristics of maternal smoking at maternity booking usually in early pregnancy: Scottish births in period 1997–2009

	Cohort		Smoking at first booking			
	N	Column %	Never smoked (row %)	Current smoker (row %)	Former smoker (row %)	Status unknown (row %)
All woman	697 003	100	56	23	9	12
Babies gender						
Male	347 399	50	58	24	9	10
Female	330 241	47	58	24	9	10
Missing	19 363	3	0	0	0	100
Maternal age (years)						
Less than 20	55 843	8	36	41	11	11
20–24	127 265	18	44	34	10	12
25–29	186 949	27	57	22	9	11
30–34	202 008	29	64	16	8	12
35–39	103 457	15	65	15	7	13
40+ years	18 848	3	64	16	6	14
Invalid	2 633	0	0	0	0	100
Area deprivation (quintiles)						
Most deprived	173 490	25	41	38	8	13
2	140 730	20	51	28	9	12
3	129 649	19	58	21	9	12
4	129 072	19	65	14	9	12
Least deprived	124 048	18	73	8	7	12
Missing	14	0	43	21	7	29
Mother's country of birth						
Non-British	57 334	8	69	9	6	17
British	639 647	92	55	24	9	12
Not Known	22	0	41	32	9	18
Mother's occupation						
Managerial/professional	187 513	27	71	9	7	13
Intermediate	156 848	23	63	15	9	12
Semiroutine/routine	200 774	29	46	33	10	11
Students	11 885	2	51	24	10	14
Other/not known	139 983	20	44	36	7	13
Father's occupation						
Managerial/professional	198 936	29	71	9	7	13
Intermediate	116 614	17	63	16	8	12
Semiroutine/routine	309 886	44	49	30	10	12
Students	7 112	1	59	17	7	17
Other/not known	64 455	9	34	44	8	14
Marital status						
Cohabiting	202 610	29	44	33	11	13
Single/parents living apart	113 254	16	34	44	9	13
Married	381 139	55	69	11	7	12
Parity						
No siblings/first child	309 769	44	57	22	10	11

Continued

Table 1 Continued

	Cohort		Smoking at first booking			
	N	Column %	Never smoked (row %)	Current smoker (row %)	Former smoker (row %)	Status unknown (row %)
One or more siblings	363 374	52	59	25	7	9
Other/unknown	23 860	3	12	4	2	82
Infant feeding at 6–8 weeks						
Both	46 877	7	65	13	8	14
Bottle	342 483	49	50	29	9	12
Breast	144 124	21	70	9	8	13
Invalid/unknown	163 519	23	55	25	9	12
Mode of delivery						
Normal/spontaneous	440 632	63	56	26	9	9
Instrumental/breech births	85 143	12	61	19	10	10
Caesarean (elective/emergency)	151 537	22	60	20	9	11
Other/unknown/no data recorded	19 691	3	1	0	0	98
Maternal religious background						
Buddhist	4 186	1	72	3	4	22
Christian	663 563	95	55	24	9	12
Hindu	1 848	0	75	5	2	18
Jewish	432	0	50	26	9	15
Muslim	18 057	3	79	4	2	15
Sikh	1 697	0	82	3	2	14
Non-applicable	7 220	1	58	18	7	17

## Key results

### Increased postneonatal mortality

More than 20% of postneonatal deaths, many of which will be defined as Sudden Unexpected Deaths in Infancy,<sup>21</sup> would probably have been avoided if mothers had not smoked during pregnancy. This result is consistent with the findings of a study of over 3 million births in the USA which reported a 40% higher infant mortality among the children of mothers who smoked, with a dose-dependent relationship with the number of cigarettes smoked.<sup>22</sup> An analysis of Swedish birth registry data which adjusted for a range of confounders found an increased risk of neonatal mortality.<sup>23</sup>

### Increased risk of small for gestational age

Births that are small for gestational age are associated with long-term ill health including hypertension and obesity, and 27.5% of these were attributable to maternal smoking. These findings are consistent with a number of large studies that have found associations between maternal smoking and intrauterine growth retardation.<sup>23 24</sup>


### Increased risk of bacterial meningitis

The rare but life-threatening condition—bacterial meningitis—was 49% more likely to occur among 0–5 years old

children born to mothers who smoked, making 11.8% of cases attributable to smoking during pregnancy. This increase in an infectious disease outcome has been described before for infections in the first year of life including meningitis.<sup>25</sup> Several studies have reported an increased risk of meningococcal disease including meningitis among the children of mothers who smoke.<sup>26–28</sup>

### Increased risk of admission for acute upper and lower respiratory infections, asthma and wheezing as well as bronchiolitis

The proportion of hospital admissions attributable to maternal smoking for common infant and child conditions including bronchiolitis (10.1%), and upper and lower respiratory infections (6.7%) in the first year of life, and asthma (7.1%) age 1–5 years are both statistically and, more importantly, clinically significant. These results are consistent with the findings from other studies that have reported an increased risk of hospital admission for bronchiolitis among the children of mothers who smoked during pregnancy.<sup>29</sup> Several reviews have found an association between maternal smoking and the subsequent development of wheezing or asthma among children.<sup>30</sup> Two large case-control studies reported by Metzger *et al* found that maternal smoking was associated with an



**Figure 1** Admissions to hospital for children born in Scotland between 1997 and 2009.

increased risk of hospitalisation for infectious disease, including a 69% increase in the odds of hospitalisation for respiratory infection.<sup>25</sup>

### Strengths and limitations

Other similar studies generally only looked at one or two outcomes,<sup>31</sup> for example, birth outcomes (small for gestational age and stillbirth) usually derived from a single dataset.<sup>32</sup> It is the linkage of routinely collected datasets that sets this study apart, allowing, in this case, hospitalisation outcomes to be gathered over time. This is possible in Scotland because the excellent routine data collection systems allow data linkage between separate datasets on all infants and their mothers.<sup>19</sup> Data linkage will only be improved by widespread use of the CHI, a unique identifier (<https://www.ndc.scot.nhs.uk/Dictionary-A-Z/Definitions/index.asp?ID=128&Title=CHI>), now given to every baby in Scotland which can be linked to the mothers' CHI number. Future linkage studies will provide a powerful tool with many research and surveillance applications to support improved healthcare and surveillance for the population of Scotland.

In addition, in our study, we modelled hospitalisations for long bone fractures—not causally associated with maternal smoking—to act as a control outcome. This analysis suggests that the adjustment for socioeconomic confounders was sufficient.

Smoking behaviour in pregnancy was collected as self-reported data at a woman's first antenatal booking appointment which usually takes place within the first 3 months of pregnancy. We know that 25% of current smokers at maternity booking do not admit to their smoking habit.<sup>14</sup> We also know that: 'Although many women quit smoking

during their pregnancy, relapse rates are high and most start smoking again within 6 months of giving birth'.<sup>13</sup> Both these issues would suggest that true smokers were in the non-smoker category in our analysis. This will have reduced our overall estimates of PAF. Therefore, the figures we have calculated are likely to be an underestimate of real effects of maternal smoking on outcomes. This study only looked at babies who were born alive and therefore does not include pregnancy outcomes leading to fetal death, that is, early miscarriage or stillbirth, both of which are devastating and are associated with maternal smoking.<sup>1 3 33 34</sup> This study focussed on hospitalisation of infants and children to 5 years of age, and therefore does not include older child illnesses or those only requiring support from primary care. This may explain the low PAF for otitis media, as children are treated in primary care and only rarely admitted to hospital. It also excluded short-term and long-term consequences of smoking for the mother herself where continued smoking will shorten life by on average 10 years.<sup>35</sup> Maternal smoking information was recorded as unknown or missing for 12% of individuals in the cohort. Pregnancy and mortality outcomes were significantly worse within this group, suggesting that perhaps unrecorded smoking or another unknown risk factor was important.

Although the coverage and completeness of variables using the routine datasets was relatively high, the study was limited by the availability of confounders on the linked dataset and uncertainty over the overall duration of maternal smoking. We do, however, know that few women who self-report as smokers at maternity booking quit during or probably after pregnancy.<sup>15</sup> In addition,

**Table 2** Likelihood of specific perinatal and mortality outcomes and smoking status at first booking

Outcome/smoking status at booking	Infant having perinatal or mortality outcome			All infants (N)	Crude OR HR (95% CI)	Adjusted* OR	
	N	Column %	Row %			HR (95% CI)	HR (95% CI)
<b>Neonatal mortality†</b>	<b>1843</b>	<b>100.0</b>	<b>0.3</b>	<b>697 003</b>			
Never smoked	789	42.8	0.2	391 861	1.00 (reference)	1.00 (reference)	6.7
Current smoker	503	27.3	0.3	159 819	1.56 (1.40 to 1.75)	1.32 (1.17 to 1.49)	
Former smoker	105	5.7	0.2	59 326	0.88 (0.72 to 1.08)	0.82 (0.66 to 1.00)	
Unknown/missing	446	24.2	0.5	85 997	2.58 (2.30 to 2.90)	2.01 (1.75 to 2.30)	
<b>Postneonatal mortality‡</b>	<b>1074</b>	<b>100.0</b>	<b>0.2</b>	<b>697 003</b>			
Never smoked	383	35.7	0.1	391 861	1.00 (reference)	1.00 (reference)	22.3
Current smoker	444	41.3	0.3	159 819	2.85 (2.48 to 3.27)	2.18 (1.87 to 2.53)	
Former smoker	72	6.7	0.1	59 326	1.24 (0.97 to 1.60)	1.16 (0.90 to 1.49)	
Unknown/missing	175	16.3	0.2	85 997	2.08 (1.74 to 2.49)	1.88 (1.54 to 2.29)	
<b>Small for gestational age§, ¶</b>	<b>56 588</b>	<b>100.0</b>	<b>8.4</b>	<b>676 095</b>			
Never smoked	22 535	39.8	5.8	391 068	1.00 (reference)	1.00 (reference)	27.5
Current smoker	24 863	43.9	15.6	159 433	3.02 (2.96 to 3.08)	2.67 (2.62 to 2.73)	
Former smoker	3826	6.8	6.5	59 203	1.13 (1.09 to 1.17)	1.03 (0.99 to 1.06)	
Unknown/missing	5364	9.5	8.1	66 391	1.44 (1.39 to 1.48)	1.31 (1.26 to 1.35)	
<b>Premature delivery¶, **</b>	<b>40 492</b>	<b>100.0</b>	<b>6.0</b>	<b>677 449</b>			
Never smoked	19 855	49.0	5.1	391 748	1.00 (reference)	1.00 (reference)	8.8
Current smoker	12 396	30.6	7.8	159 761	1.58 (1.54 to 1.61)	1.41 (1.37 to 1.44)	
Former smoker	3002	7.4	5.1	59 308	1.00 (0.96 to 1.04)	0.93 (0.89 to 0.97)	
Unknown/missing	5239	12.9	7.9	66 632	1.60 (1.55 to 1.65)	1.49 (1.44 to 1.54)	

\*Adjusted for: maternal age; infant gender, SIMD; mothers socioeconomic status; fathers socioeconomic status; parity; mode of delivery and country of birth.

†Neonatal mortality describes a baby that has died during the first 28 days of life.

‡Postneonatal mortality is defined as those infant deaths in which the infant is more than 28 days and less than 1 year of age.

§Under 10th centile—small for gestational age.

¶Unknown/missing data excluded from modelling.

\*\*Birth of a baby at fewer than 37 weeks gestational age.

PAF, population attributable fraction; SIMD, Scottish Index of Multiple Deprivation.

an important limitation of this study is that there was a high level of missing data for some of the variables used, particularly infant feeding method and maternal occupation. This is likely to relate to the phased introduction of the Child Health Surveillance Programme across Scotland during the study period which has been recognised for infant feeding data. The impact of missing data on maternal occupation may not have been significant because of the inclusion of the two other measures of socioeconomic level in the regression models: area deprivation and paternal occupation.

### Interpretation

Overall, it is likely that there is an underestimation of the association between maternal smoking and poor outcomes including hospitalisation as 25% of pregnant smokers deny their habit when asked at maternity booking.<sup>14</sup> This level of under-reporting may increase the attributable fraction for each condition by a third, making, for example, the proportion of small for gestational age babies attributable

to maternal smoking 36% rather than 27.5%. Alternatively, if those with missing smoking data were all smokers, the estimates we present would be reduced. However, our analysis of ‘small for gestational age’ (table 2)—the most well-known effect of maternal smoking during pregnancy—does not suggest that all those with missing smoking data are in fact current smokers. The OR for ‘unknown/missing’ smoking status is 1.31 (1.26, 1.35) and for ‘current smoker’ is 2.67 (2.62, 2.73).

This study adds to the evidence identifying key adverse childhood outcomes of maternal smoking.

Maternal smoking is still making a substantial contribution to poor pregnancy outcomes and morbidity in early childhood.

This study confirms previous findings that maternal smoking is associated with large increases in the risks of being small for gestational age, of postneonatal mortality and of premature delivery, but less evidence of impact on neonatal mortality.

**Table 3** Rate of hospitalisation for selected child outcomes and smoking status at first booking for respiratory infections, asthma and bronchiolitis for (A) the whole period of follow-up; and for two separate periods of follow-up: (B) from birth to first birthday and (C) from first birthday to fifth birthday

<b>(A) Follow-up from birth to fifth birthday</b>												
	Infants at risk (N)			Acute upper and lower respiratory infections			Asthma			Bronchiolitis		
	N	Column %	Row %	N	Column %	Row %	N	Column %	Row %	N	Column %	Row %
<b>Smoking status at first booking</b>	<b>697 003</b>	<b>100.0</b>	<b>8.1</b>	<b>7549</b>	<b>100.0</b>	<b>1.1</b>	<b>24 088</b>	<b>100.0</b>	<b>1.1</b>	<b>24 088</b>	<b>100.0</b>	<b>3.5</b>
Never smoked	391 861	54.0	7.8	3678	48.7	0.9	11 354	47.1	0.9	11 354	47.1	2.9
Current smoker	159 819	26.4	9.4	2412	32.0	1.5	7939	33.0	1.5	7939	33.0	5.0
Former smoker	59 326	8.8	8.4	654	8.7	1.1	1821	7.6	1.1	1821	7.6	3.1
Unknown/missing	85 997	10.9	7.2	805	10.7	0.9	2974	12.3	0.9	2974	12.3	3.5
<b>(B) Follow-up from birth to first birthday</b>												
	Admission for specified outcome			Infants at risk (N)			Crude HR	Adjusted* HR	PAF (%)			
	N	Column %	Row %	N	Column %	Row %						
<b>Acute upper and lower respiratory infections</b>	<b>18 541</b>	<b>100.0</b>	<b>2.7</b>	<b>697 003</b>	<b>100.0</b>	<b>2.7</b>	<b>HR (95% CI)</b>	<b>HR (95% CI)</b>	<b>PAF (%)</b>			
Never smoked	9573	51.6	2.4	391 861	1.00 (reference)	1.00 (reference)	1.00 (reference)	1.00 (reference)	6.7			
Current smoker	5537	29.9	3.5	159 819	1.43 (1.38 to 1.48)	1.43 (1.38 to 1.48)	1.29 (1.25 to 1.34)	1.29 (1.25 to 1.34)				
Former smoker	1484	8.0	2.5	59 326	1.02 (0.97 to 1.08)	1.02 (0.97 to 1.08)	0.97 (0.92 to 1.03)	0.97 (0.92 to 1.03)				
Unknown/missing	1947	10.5	2.3	85 997	0.93 (0.89 to 0.98)	0.93 (0.89 to 0.98)	0.89 (0.84 to 0.93)	0.89 (0.84 to 0.93)				
Asthma	606	100.0	0.1	697 003								
Never smoked	242	39.9	0.1	391 861	1.00 (reference)	1.00 (reference)	1.00 (reference)	1.00 (reference)	21.9			
Current smoker	248	40.9	0.2	159 819	2.51 (2.10 to 3.00)	2.51 (2.10 to 3.00)	2.15 (1.80 to 2.57)	2.15 (1.80 to 2.57)				
Former smoker	46	7.6	0.1	59 326	1.25 (0.91 to 1.72)	1.25 (0.91 to 1.72)	1.18 (0.86 to 1.62)	1.18 (0.86 to 1.62)				
Unknown/missing	70	11.6	0.1	85 997	1.33 (1.02 to 1.73)	1.33 (1.02 to 1.73)	1.29 (0.99 to 1.69)	1.29 (0.99 to 1.69)				
Bronchiolitis	2399	100.0	2.9	697 003								
Never smoked	4996	46.6	2.4	166	1.00 (reference)	1.00 (reference)	1.00 (reference)	1.00 (reference)	10.1			
Current smoker	6861	33.6	4.3	159 819	1.79 (1.73 to 1.84)	1.79 (1.73 to 1.84)	1.43 (1.38 to 1.48)	1.43 (1.38 to 1.48)				
Former smoker	1534	7.5	2.6	59 326	1.07 (1.01 to 1.13)	1.07 (1.01 to 1.13)	1.03 (0.97 to 1.08)	1.03 (0.97 to 1.08)				
Unknown/missing	2505	12.3	2.9	85 997	1.21 (1.16 to 1.27)	1.21 (1.16 to 1.27)	1.15 (1.09 to 1.21)	1.15 (1.09 to 1.21)				

Continued

Table 3 Continued

## (C) Follow-up from first birthday to fifth birthday

Acute upper and lower respiratory infections	Admission for specified outcome						Crude HR HR (95% CI)	Adjusted* HR HR (95% CI)	PAF (%)
	N		Row %		Infants at risk (N)†	662 454			
	38074	100.0	Column %	5.7					
Never smoked	20976	55.1	5.6	373 436	1.00 (reference)	1.00 (reference)	-0.2		
Current smoker	9408	24.7	6.2	151 010	1.09 (1.07 to 1.12)	0.99 (0.97 to 1.02)			
Former smoker	3482	9.1	6.1	56 645	1.09 (1.06 to 1.13)	1.04 (1.00 to 1.08)			
Unknown/missing	4208	11.1	5.2	81 363	0.95 (0.92 to 0.98)	0.90 (0.87 to 0.94)			
Asthma	6943	100.0	1.0	679 588					
Never smoked	3436	49.5	0.9	382 327	1.00 (reference)	1.00 (reference)	7.1		
Current smoker	2164	31.2	1.4	156 123	1.51 (1.43 to 1.59)	1.29 (1.22 to 1.37)			
Former smoker	608	8.8	1.0	58 025	1.16 (1.07 to 1.27)	1.10 (1.00 to 1.20)			
Unknown/missing	735	10.6	0.9	83 113	1.04 (0.96 to 1.12)	1.01 (0.92 to 1.10)			
Bronchiolitis	3689	100.0	0.6	660 119					
Never smoked	1855	50.3	0.5	373 219	1.00 (reference)	1.00 (reference)	4.0		
Current smoker	1078	29.2	0.7	149 609	1.45 (1.34 to 1.56)	1.16 (1.08 to 1.25)			
Former smoker	287	7.8	0.5	56 561	1.02 (0.90 to 1.15)	0.98 (0.87 to 1.11)			
Unknown/missing	469	12.7	0.6	80 730	1.18 (1.06 to 1.30)	1.11 (1.01 to 1.24)			

\*Adjusted for: maternal age; infant gender, SIMD; mothers socioeconomic status; fathers socioeconomic status; parity; mode of delivery, infant feeding at 6–8 weeks and country of birth.

†The number of infants at risk from their first birthday to fifth birthday is the total number of infants aged 0–1 year (697 003) minus the specific outcome cases and any deaths or migration from 0–1 year.

PAF, population attributable fraction; SIMD, Scottish Index of Multiple Deprivation.

**Table 4** Likelihood of specific perinatal and mortality outcomes and smoking status at first booking

Smoking status at first booking	N	Column %	Row %	All infants (N)	Crude HR	Adjusted* HR	PAF (%)
					HR (95% CI)	HR (95% CI)	
<b>Bacterial meningitis</b>	<b>1251</b>	<b>100.0</b>	<b>0.2</b>	<b>697 003</b>			
Never smoked	525	42	0.1	391 861	1.00 (reference)	1.00 (reference)	11.8
Current smoker	450	36	0.3	159 819	2.09 (1.84 to 2.37)	1.49 (1.30 to 1.71)	
Former smoker	125	10	0.2	59 326	1.57 (1.29 to 1.91)	1.35 (1.11 to 1.64)	
Unknown/missing	151	12	0.2	85 997	1.34 (1.11 to 1.60)	1.29 (1.06 to 1.57)	
<b>Acute and chronic otitis media and mastoiditis</b>	<b>8879</b>	<b>100.0</b>	<b>1.3</b>	<b>697 003</b>			
Never smoked	4866	55	1.2	391 861	1.00 (reference)	1.00 (reference)	1.7
Current smoker	2148	24	1.3	159 819	1.05 (1.00 to 1.11)	1.07 (1.02 to 1.14)	
Former smoker	755	9	1.3	59 326	1.02 (0.94 to 1.10)	1.01 (0.94 to 1.10)	
Unknown/missing	1110	13	1.3	85 997	1.10 (1.03 to 1.18)	1.13 (1.06 to 1.22)	
<b>Bronchopulmonary dysplasia</b>	<b>358</b>	<b>100.0</b>	<b>0.1</b>	<b>697 003</b>			
Never smoked	141	39	0.0	391 861	1.00 (reference)	1.00 (reference)	10.1
Current smoker	109	30	0.1	159 819	1.90 (1.48 to 2.44)	1.50 (1.15 to 1.96)	
Former smoker	22	6	0.0	59 326	1.03 (0.66 to 1.61)	0.90 (0.57 to 1.42)	
Unknown/missing	86	24	0.1	85 997	2.79 (2.13 to 3.65)	3.21 (2.45 to 4.22)	
<b>Long bone fracture</b>	<b>5757</b>	<b>100.0</b>	<b>0.8</b>	<b>697 003</b>			
Never smoked	3156	55	0.8	391 861	1.00 (reference)	1.00 (reference)	-1.5
Current smoker	1466	25	0.9	159 819	1.11 (1.04 to 1.18)	0.94 (0.88 to 1.01)	
Former smoker	532	9	0.9	59 326	1.11 (1.01 to 1.21)	1.06 (0.96 to 1.16)	
Unknown/missing	603	10	0.7	85 997	0.93 (0.85 to 1.02)	0.92 (0.83 to 1.01)	

\*Adjusted for: maternal age; infant gender, SIMD; mothers socioeconomic status; fathers socioeconomic status; parity; mode of delivery, infant feeding at 6–8 weeks and country of birth.

PAF, population attributable fraction; SIMD, Scottish Index of Multiple Deprivation.

Meningitis, bronchopulmonary dysplasia, asthma, bronchiolitis and acute upper and lower respiratory infections (in the first year of life) make large contributions to the increased risk of hospital admission; ear infections are less prominent, probably because nearly all are treated in primary care, though still a significant risk.

The strength of the association between maternal smoking and increased infant and child morbidity and mortality, consistent with other studies, provides convincing evidence of the benefits of maternal smoking abstinence on child health.

### Generalisability

This study presents results from a ‘whole population’ cohort of infants born in Scotland over a 13-year period from 1997 to 2009. Children have been followed through to March 2012. These results are contemporary and are likely to generalise to most other UK countries and other countries with high levels of smoking during pregnancy.

**Acknowledgements** Omotomilola Ajetunmbi, Information Services Division, NHS National Services Scotland, Edinburgh, Scotland, UK who created the dataset with BW for the breastfeeding and hospitalisation analysis.

**Contributors** All authors helped plan this study. RL analysed the data and commented on the manuscript drafts. CF supervised RL and commented on the manuscript drafts. DMT wrote the manuscript. BW commented on the manuscript drafts and helped create the database. RW commented on the manuscript drafts.

**Funding** The authors have not declared a specific grant for this research from any funding agency in the public, commercial or not-for-profit sectors.

**Competing interests** None declared.

**Patient consent for publication** Not required.

**Provenance and peer review** Not commissioned; externally peer reviewed.

**Data sharing statement** No additional unpublished data are available from the study.

**Author note** Technical appendix, statistical code and dataset available from: Information Services Division NHS National Services Scotland, Edinburgh, Scotland.

**Open access** This is an open access article distributed in accordance with the Creative Commons Attribution Non Commercial (CC BY-NC 4.0) license, which permits others to distribute, remix, adapt, build upon this work non-commercially, and license their derivative works on different terms, provided the original work is properly cited, appropriate credit is given, any changes made indicated, and the use is non-commercial. See: <http://creativecommons.org/licenses/by-nc/4.0/>.

### REFERENCES

1. Flenady V, Koopmans L, Middleton P, *et al*. Major risk factors for stillbirth in high-income countries: a systematic review and meta-analysis. *Lancet* 2011;377:1331–40.

2. ONS (Office for National Statistics). Infant and perinatal mortality in England and Wales by social and biological factors. *Stat Bull* 2011, Key findings. [www.ons.gov.uk/ons/dcp171778\\_300596.pdf](http://www.ons.gov.uk/ons/dcp171778_300596.pdf)
3. Dietz PM, England LJ, Shapiro-Mendoza CK, *et al*. Infant morbidity and mortality attributable to prenatal smoking in the U.S. *Am J Prev Med* 2010;39:45–52.
4. Health Inequalities Unit, Department of Health. Review of the Health Inequalities Infant. Mortality PSA Target 2007; [www.perinatal.nhs.uk/smoking/health%20inequalities%20report%202007.pdf](http://www.perinatal.nhs.uk/smoking/health%20inequalities%20report%202007.pdf)
5. Delpisheh A, Kelly Y, Rizwan S, *et al*. Population attributable risk for adverse pregnancy outcomes related to smoking in adolescents and adults. *Public Health* 2007;121:861–8.
6. Gilliland FD, Li YF, Peters JM. Effects of maternal smoking during pregnancy and environmental tobacco smoke on asthma and wheezing in children. *Am J Respir Crit Care Med* 2001;163:429–36.
7. Button TM, Maughan B, McGuffin P. The relationship of maternal smoking to psychological problems in the offspring. *Early Hum Dev* 2007;83:727–32.
8. Batstra L, Hadders-Algra M, Neeleman J. Effect of antenatal exposure to maternal smoking on behavioural problems and academic achievement in childhood: prospective evidence from a Dutch birth cohort. *Early Hum Dev* 2003;75(1-2):21–33.
9. Godfrey C, Pickett KE, Parrot S, *et al*. Estimating the costs to the NHS of smoking in pregnancy for pregnant women and infants, 2010. [http://phrc.lshtm.ac.uk/papers/PHRC\\_A3-06\\_Final\\_Report.pdf](http://phrc.lshtm.ac.uk/papers/PHRC_A3-06_Final_Report.pdf)
10. Leonardi-Bee J, Jere ML, Britton J. Exposure to parental and sibling smoking and the risk of smoking uptake in childhood and adolescence: a systematic review and meta-analysis. *Thorax* 2011;66:847–55.
11. ISD Scotland website. Smoking History and Booking. <http://www.isdscotland.org/Health-Topics/Maternity-and-Births/Publications/data-tables.asp?id=1791#1791>
12. NHS Executive. New NHS Smoking Cessation Services. Health Service Circular, 1999. [http://www.dh.gov.uk/prod\\_consum\\_dh/groups/dh\\_digitalassets/@dh/@en/documents/digitalasset/dh\\_4011979.pdf](http://www.dh.gov.uk/prod_consum_dh/groups/dh_digitalassets/@dh/@en/documents/digitalasset/dh_4011979.pdf) (accessed on 29 July 2011).
13. NICE (National Institute for Health and Care Excellence). Smoking: stopping in pregnancy and after childbirth, 2010. Public Health Guidance 26. <https://www.nice.org.uk/guidance/ph26>
14. Shipton D, Tappin DM, Vadiveloo T, *et al*. Reliability of self reported smoking status by pregnant women for estimating smoking prevalence: a retrospective, cross sectional study. *BMJ* 2009;339:b4347.
15. Tappin DM, MacAskill S, Bauld L, *et al*. Smoking prevalence and smoking cessation services for pregnant women in Scotland. *Subst Abuse Treat Prev Policy* 2010;5:1.
16. HSCIC (Health and Social Care Information Centre). Infant feeding survey – 2010. 2011. <http://www.hscic.gov.uk/catalogue/PUB00648>
17. Ajetunmobi OM, Whyte B, Chalmers J, *et al*. Breastfeeding is associated with reduced childhood hospitalization: Evidence from a scottish birth cohort (1997–2009). *J Pediatr* 2015;166:620–5.
18. National Privacy Advisory Committee. [http://www.nhsns.org/pages/corporate/privacy\\_advisory\\_committee.php](http://www.nhsns.org/pages/corporate/privacy_advisory_committee.php) (Accessed Aug 2013).
19. Fleming M, Kirby B, Penny KI. Record linkage in Scotland and its applications to health research. *J Clin Nurs* 2012;21:2711–21.
20. Ajetunmobi O, Whyte B, Chalmers J, *et al*. Informing the ‘early years’ agenda in Scotland: understanding infant feeding patterns using linked datasets. *J Epidemiol Community Health* 2014;68:83–92.
21. ISD Scotland. Stillbirths and infant deaths. <http://www.isdscotland.org/health-Topics/Maternity-and-Births/Stillbirth-and-Infant-Deaths/>
22. Saihu HM, Aliyu MH, Pierre-Louis BJ, *et al*. Levels of excess infant deaths attributable to maternal smoking during pregnancy in the United States. *Matern Child Health J* 2003;7:219–27.
23. Källén K. The impact of maternal smoking during pregnancy on delivery outcome. *Eur J Public Health* 2001;11:329–33.
24. Baba S, Wikström AK, Stephansson O, *et al*. Changes in snuff and smoking habits in Swedish pregnant women and risk for small for gestational age births. *BJOG* 2013;120:456–62.
25. Metzger MJ, Halperin AC, Manhart LE, *et al*. Association of maternal smoking during pregnancy with infant hospitalization and mortality due to infectious diseases. *Pediatr Infect Dis J* 2013;32:e1–e7.
26. Goldacre MJ, Wotton CJ, Maisonneuve JJ. Maternal and perinatal factors associated with subsequent meningococcal, Haemophilus or enteroviral meningitis in children: database study. *Epidemiol Infect* 2014;142:371–8.
27. Sørensen HT, Labouriau R, Jensen ES, *et al*. Fetal growth, maternal prenatal smoking, and risk of invasive meningococcal disease: a nationwide case-control study. *Int J Epidemiol* 2004;33:816–20.
28. Yusuf HR, Roach RW, Baughman WS, *et al*. Maternal cigarette smoking and invasive meningococcal disease: a cohort study among young children in metropolitan Atlanta, 1989–1996. *Am J Public Health* 1999;89:712–7.
29. Koehoorn M, Karr CJ, Demers PA, *et al*. Descriptive epidemiological features of bronchiolitis in a population-based cohort. *Pediatrics* 2008;122:1196–203.
30. Ferrante G, Antona R, Malizia V, *et al*. Smoke exposure as a risk factor for asthma in childhood: a review of current evidence. *Allergy Asthma Proc* 2014;35:454–61.
31. Mund M, Louwen F, Klingelhofer D, *et al*. Smoking and pregnancy—a review on the first major environmental risk factor of the unborn. *Int J Environ Res Public Health* 2013;10:6485–99.
32. Erickson AC, Arbour LT. Heavy smoking during pregnancy as a marker for other risk factors of adverse birth outcomes: a population-based study in British Columbia, Canada. *BMC Public Health* 2012;12:102–11.
33. Centers for Disease Control and Prevention. PRAMStat. 2011. [http://nccd.cdc.gov/PRAMStat/rdPage.aspx?rdReport=DRH\\_PRAMS.ExploreByTopic&isClassId=CL9&isTopicId=TOP27&go=GO](http://nccd.cdc.gov/PRAMStat/rdPage.aspx?rdReport=DRH_PRAMS.ExploreByTopic&isClassId=CL9&isTopicId=TOP27&go=GO)
34. ONS. Office for National Statistics). Infant and perinatal mortality in England and Wales by social and biological factors. *Stat Bull* 2011. Key findings [www.ons.gov.uk/ons/dcp171778\\_300596.pdf](http://www.ons.gov.uk/ons/dcp171778_300596.pdf)
35. Doll R, Peto R, Boreham J, *et al*. Mortality in relation to smoking: 50 years’ observations on male British doctors. *BMJ* 2004;328:1519.