BMJ Open Effect of a 2-day communication skills training on nursing and midwifery students' empathy: a randomised controlled trial

Mustapha Alhassan¹

To cite: Alhassan M. Effect of a 2-day communication skills training on nursing and midwifery students' empathy: a randomised controlled trial. *BMJ Open* 2019;**9**:e023666. doi:10.1136/bmjopen-2018-023666

► Prepublication history and additional material for this paper are available online. To view these files, please visit the journal online (<http://dx.doi.org/10.1136/bmjopen-2018-023666>).

Received 25 April 2018

Revised 22 November 2018

Accepted 30 January 2019

ABSTRACT

Introduction Empathy is crucial to the fundamental aim and achievement of nursing and midwifery goals. Researchers agree on the positive role empathy plays in interpersonal relationships when providing healthcare. Models of good communication have been developed to assist nurses, midwives and doctors to improve their ability to communicate with patients. This study investigated the effect of a 2-day communication skills training (CST) on nursing and midwifery students' empathy in a randomised controlled trial.

Methods The two groups had a baseline data collection at the same time. The intervention group had a CST, followed by post-test on day 3. The control group had post-test on day 4 just before their CST. The empathy outcome was measured with Jefferson Scales of Empathy-Health Professions Student version. Both groups had a follow-up test at the same time 6 months after the CST.

Results In this study, there was no statistically significant difference in the scores of empathy between the groups $F(1, 171)=0.18, p=0.675$. The intervention group had baseline T1 (M=109.8, SD=9.8, $d=0.160$), and post-test T2 (M=111.9, SD=9.0, $d=0.201$), whereas the control group had baseline T1 (M=107.9, SD=11.46, $d=0.160$), and post-test T2 (M=110.0, SD=11.0, $d=0.201$). Baseline data were collected on 15 June 2013.

Conclusions This study has shown that empathy may not be enhanced within a short period after CST.

Strengths and limitations of this study

- This study used nursing and midwifery students who were actively involved. The use of various methods like group discussions, roleplays, videos, short presentations and brainstorming sessions in the delivery of the communication skills training was also a positive development, since such methods take care of individual differences.
- There was allocation concealment to the researcher, research assistants and the participants.
- The empathy measure used was Jefferson Scales of Empathy-Health Professions Student version has good construct validity and criterion-related validity which has been reported.
- A limitation of this study is the generalisation of the results to other healthcare professionals. As a self-report outcome, results of this study cannot be generalised beyond the characteristics of this sample.
- Confounding factors can also limit the generalisability of this study; the study could not control for interaction between the groups during the period of the study. This could lead to the problem of contamination between the groups (ie, those in the intervention group talking to those in the control group after their training session days).

INTRODUCTION

Empathy is crucial to the fundamental aim and achievement of nursing and midwifery goals.¹ Within the nursing and midwifery field, such skills are considered indicative of best practice.² It has also been stressed that empathy is a necessary factor in the provision of quality nursing care.³ Researchers agree on the positive role empathy plays in interpersonal relationships when providing healthcare.^{4,5}

Models of good communication have been developed to assist nurses, midwives and doctors to improve their ability to communicate with patients.⁶⁻¹² A health maintenance organisation (Kaiser Permanente in the USA)

developed the Four Habits Model, which they have used for more than 20 years and is an effective programme for clinical communication.^{6,7} The model has been anchored into four habits: 'invest in the beginning (Habit I), elicit patients' perspective (Habit II), demonstrate empathy (Habit III), and invest in the end (Habit IV)'.^{6,7} The habits from this theory was the basis of the communication skills training (CST) that was developed and used for this study. The other theoretical model called the Person-Centred Nursing Framework (PCNF)¹³ was an essential component of the CST. Emphasis was made on PCNF necessary care processes of working with the patients beliefs and values,

© Author(s) (or their employer(s)) 2019. Re-use permitted under CC BY-NC. No commercial re-use. See rights and permissions. Published by BMJ.

School of Allied Health Sciences, University for Development Studies, Tamale, Ghana

Correspondence to

Dr Mustapha Alhassan; mustapha@uds.edu.gh

engagement, shared decision making, having sympathetic presence and providing holistic care.¹³

Objective

To investigate the effect of a 2-day CST on nursing and midwifery students' (NMS) empathy in a randomised controlled trial.

METHODOLOGY

Design and sample

This study was a 2 (intervention condition, between) x 3 (time, repeated) design in a randomised controlled trial (RCT) conducted at Tamale Nurses and Midwives College Ghana. The sample consisted of nursing students (n=181) and midwifery students (n=49).

Power analysis

The sample size of the participants was based on a power analysis. Relationship have been shown between training interventions and improved communication skills, measured with Roter Interaction Analysis System, with an effect size between medium and high.¹⁴ Fixing the effect size medium ($d=0.25$), using a two-tailed significance test ($p=0.05$), a sample size of 197 resulted in an acceptable power coefficient of 0.95.¹⁵ The sample size was computed using G*Power software.^{15 16}

Informed consent

Informed consent was not written and participants were told that taking part in the CST and answering the questionnaires meant their consent and an agreement to any publication from it. Participants were informed of the objectives of the study but not in detail such that they would know that the study wants to determine their empathy level, but were also given opportunity to ask questions to enable them to decide to take part in this study. Participants were informed that they could refuse to take part in the research at any time without having to face any consequence.

Patient and public involvement

Participants in this study were NMS and patients were not involved. The students as well as their tutors were involved in the design of the CST guide.

Criteria of inclusion and exclusion

The inclusion and exclusion criteria are presented in box 1.

Randomisation

There was allocation concealment to the researcher, research assistants and the participants. The researcher (MA) and research assistants conducted this by allowing participants to pick numbers written on papers, which had been randomly shuffled in a box. NMS were separated before random assignment to ensure that both professions were approximately equally represented

Box 1 Inclusion and exclusion criteria

Inclusion criteria

- ▶ Nursing and midwifery students (NMS) in their second year at Tamale Nursing and Midwifery College.
- ▶ NMS whose ages were above 18 years.
- ▶ NMS in Tamale Nursing and Midwifery College who were available for follow-up data collection after 6 months.

Exclusion criteria

- ▶ NMS who were not studying at Tamale Nursing and Midwifery College.
- ▶ NMS whose ages were below 18 years.
- ▶ NMS in Tamale Nursing and Midwifery College who were not available for follow-up data collection after 6 months.

in the groups. Therefore, participants were randomly assigned to either intervention group or a control group.

Procedure

The two groups had a baseline data collection (T1) at the same time. The intervention group had a CST, followed by post-test (T2) on day 3. The control group had post-test (T2) on day 4 just before their CST. The CST for both groups were the same. The tutors were aware of the training and data collection for the intervention and control group; however, the intervention and control groups were not aware. The outcome was measured with Jefferson Scales of Empathy-Health Professions Student version (JSE HPS version).¹⁷ Both groups had a follow-up test (T3) at the same time 6 months after the CST (figure 1).

Outcome measure

The outcome was empathy measured with JSE HPS version.⁴ JSE HPS version was in English. This questionnaire was administered in English. This is because the students are very fluent in English. They are taught in English from primary school because English is the official language in Ghana, and they practice in English. There are different versions of the JSE. The versions are comparable in content. Slight changes are made in the words such that the text will be suitable for the planned health professionals. JSE HPS version¹⁷ has 20 items in a Likert-type format from 1 (strongly disagree) to 7 (strongly agree). It has 10 negatively worded items. The negatively worded items were items 1, 3, 6, 7, 8, 11, 12, 14, 18 and 19.¹⁷

The scoring of the questionnaire was according to the scoring algorithm of JSE. According to JSE, 'a respondent must answer at least 16 (80%) of the 20 items; otherwise the form should be regarded as incomplete and excluded from the data analysis. If a respondent fails to answer 4 or fewer items, the missing values should be replaced with the mean score calculated from the items the respondent completed'.¹⁷ To score the questionnaire, the negatively worded items were reversed scored (from 1 [strongly agree] to 7 [strongly disagree]), while the other items are directly scored on their Likert weights from 1 (strongly

Figure 1 Flowchart showing enrolment, randomisation, communication skills training and data collection.

disagree) to 7 (strongly agree). The total score was the sum of all item scores. The higher the total empathy scores, the higher the empathic behavioural orientation. The maximum total score for each participant is 140 and the minimum score is 20. Higher total scores indicate higher empathy, whereas lower total scores indicate lower empathy.¹⁷ According to the owners of JSE, it takes

5–10 min to complete, although they do not endorse a time limit for completing it.¹⁷

Psychometric properties

Construct validity and criterion-related validity of the JSE HPS version have been reported.¹⁸ Hojat *et al*¹⁷ have reported that internal consistency reliability of this

Table 1 Demographic data

	Intervention group (n=93)	Control group (n=80)
Characteristics	n (%)	n (%)
Age		
>18 years	5 (6)	1 (1)
19–21 years	42 (45)	32 (40)
22–24 years	41 (44)	45 (57)
25–27 years	2 (2)	1 (1)
28–30 years	3 (3)	1 (1)
31 years and above	0 (0)	0 (0)
Gender		
Female	68 (73)	44 (55)
Male	25 (27)	36 (45)
Specialty		
Nursing student	62 (67)	69 (86)
Midwifery students	31 (33)	11 (14)
Marital status		
Married	2 (2)	9 (11)
Unmarried	90 (97)	70 (88)
Divorced	1 (1)	1 (1)
Religion		
Christianity	51 (55)	30 (38)
Islam	40 (43)	48 (60)
Other	2 (2)	2 (2)
Number of children		
No child	92 (99)	72 (90)
One child	1 (1)	2 (2)
Two children	0 (0)	4 (5)
Three children	0 (0)	2 (3)
Four children and above	0 (0)	0 (0)
Ethnicity		
Akan	11 (12)	5 (6)
Dagomba	28 (30)	34 (43)
Ewe	2 (2)	5 (6)
Fanti	6 (6)	3 (4)
Frafra (Grunsi)	10 (12)	2 (2)
Ga-Adangme	3 (3)	0 (0)
Gonja	8 (9)	3 (4)
Kotokoli	0 (0)	3 (4)
Basare/Bisa	0 (0)	2 (2)
Kasina/Bulsa	0 (0)	3 (4)
Dagati/Sisala	5 (5)	4 (5)
Other tribes	20 (21)	16 (20)
Academic writing and communication (AWC)		
None	10 (11)	13 (16)
One week	0 (0)	0 (0)

Continued

Table 1 Continued

	Intervention group (n=93)	Control group (n=80)
Characteristics	n (%)	n (%)
Two weeks	0 (0)	1 (1)
Three weeks	0 (0)	0 (0)
One month	1 (1)	0 (0)
Two months	0 (0)	1 (1)
Three months	3 (3)	2 (3)
Four months (one semester)	70 (75)	57 (71)
Two semesters	5 (6)	6 (8)
Three semesters	3 (3)	0 (0)
Four semesters	0 (0)	0 (0)
Above four semesters	1 (1)	0 (0)

n=sample size in a particular group.

version was 0.89 for medical students and 0.87 for house officers. Hojat *et al*⁴ has reported a test-retest reliability for the JSE HPS version as 0.65 ($p<0.01$). In their report, they said it was relatively low in magnitude, but acceptable for that kind of instrument considering the time interval between the test.⁴

Data analysis

Analysis of variance was used to test the hypothesis that there were statistically significant differences between the two groups at three time points. A Shapiro-Wilk's test ($p<0.05$)^{19 20} and a visual inspection of their histograms showed variable scores were approximately normally distributed. I have included the results as supplementary file in the form of principal component analysis, χ^2 test, independent t-test and Scree plot (see online supplementary tables 1–3 and figure 1).

A significance level of $p<0.05$ was planned. However, because several independent analyses (5) were performed on the data, the significance level of $p<0.05$ was adjusted to $p<0.01$ in interpreting the results using Bonferroni correction.²¹ 'The Bonferroni correction is an adjustment made to p values when several dependent or independent statistical tests are being performed simultaneously on a single data set'.²¹ In this study, Bonferroni correction was computed by taking the critical p value (α) and divided it by the several dependent analyses (0.05/5) resulting in $p<0.01$. All data were analysed using SPSS.

Communication skills training

The author (MA), who was the main trainer, designed and developed the training guide using 'Four Habits Model'²² and PCNF.¹³

Subsequently, the researcher trained a cotrainer (AAM) who assisted in the CST as well as in the data collection. The data was analysed by the author (MA) without blinding. The trainers used various methods to deliver

Table 2 Descriptive statistics of empathy

Time	Group	N=173				95% CI for Cohen's d	
		n	M	SD	Cohen's d	Lower	Upper
Baseline (T1)	Intervention	93	109.8	9.8	0.160	-0.139	0.458
	Control	80	107.9	11.5			
Post-test (T2)	Intervention	93	111.9	9.0	0.201	-0.098	0.499
	Control	80	110.0	11.0			
Follow-up (T3)	Intervention	93	109.4	10.4	-0.252	-0.551	0.048
	Control	80	111.9	8.3			

N, total sample size; n, group sample size; M, mean score.

the training. The methods were small group discussions, brainstorming, personal experience from participants, group reports, roleplaying, questions and answers, videos and summaries. Therefore, the training was on shared agenda. This approach made it possible for participants to share their previous training knowledge and ideas.

At the end of the training, participants were provided with photocopies of some relevant material as well as useful reference books and literature that will enable nurses and midwives to learn effective communication with patients.

RESULTS

Demographic data

Participants (n=173) were made of intervention group (n=93) and control group (n=80). The demographic data are presented in [table 1](#).

Descriptive statistics

The scores showed that there were slight increases in the intervention group from baseline T1 (M=109.8, SD=9.8, $d=0.160$) to post-test T2 (M=111.9, SD=9.0, $d=0.201$) as compared with the control group from baseline T1 (M=107.9, SD=11.5, $d=0.160$) to post-test T2 (M=110.0, SD=11.0, $d=0.201$) ([table 2](#)).

Inferential statistics

This study showed that there was no statistically significant difference in the scores of empathy between the groups $F(1, 171)=0.18$, $p=0.675$ ([table 3](#)).

In this study, there was no statistically significant effect between CST, the demographic variables of age, marital

status, specialisation, ethnicity and religion as well as academic writing and communication ([table 4](#)).

DISCUSSIONS

In this study, there was no statistically significant difference between the groups $F(1, 171)=0.18$, $p=0.675$ ([table 3](#)). The findings from this study are in contrast to the findings from a similar study that showed enhancement of empathy in nurses.^{23–26} For example, a study reported that nursing students empathy moderately increased in scores (M=88.63; SD=8.93).^{23–24} Further in contrast, another study found statistically significant effect in empathy scores following a training.²⁵

Research has shown that there are a number of studies that doubt the effectiveness of empathy training programmes in nursing education and rather reported stability in empathy.^{27–35} A study by La Monica *et al*³⁰ did not find improvement in empathy outcomes. In a related study, they found stability in empathy after a short-term education (M=20.7–22.6; SD=3.0–5.0).³¹ In another research, it was reported that empathy was stable.³²

Research has demonstrated that there is a relationship between empathy and demographic variables of gender, education and experience. In this study, there were no statistically significant differences in empathy and the demographic variables of gender, age, marital status, specialisation, religion, number of children and ethnicity between the both groups. The findings from this study are inconsistent with other studies where females empathy scores are reported to be higher than

Table 3 Inferential statistics empathy

Source	Type III SS	df	MS	F	Sig.
Intercept	6 259 179.09	1	6 259 179.09	55379.73	0.000
Group	19.91	1	19.91	.18	0.675
Error	19326.92	171	113.02		

Significance level $p<0.01$.

Measurement is by time point.

Transformed variable is by average.

Effect of the communication skills training, the empathy scores, and the demographic variables.

df, degrees of freedom; F, statistic; MS, mean square; Sig, significance level; SS, sum of squares.

Table 4 Effect of the communication skills training, the empathy scores and the demographic variables

Source	Type III SS	df	MS	F	Sig.
Intercept	35 145.96	1	35 145.96	321.61	0.000
Group*gender	738.51	2	369.25	3.38	0.037
Group*age	431.25	2	215.63	1.97	0.142
Group*marital status	72.00	2	36.00	0.33	0.720
Group*specialisation	241.87	2	120.94	1.11	0.333
Group*religion	219.32	2	109.66	1.00	0.369
Group*ethnicity	440.74	2	220.37	2.02	0.137
Group*AWC	69.25	2	34.63	0.32	0.729
Error	17 266.55	158	109.28		

Significance level $p < 0.01$.

Measurement is by time point.

Transformed variable is by average.

AWC, academic writing and communication; df, degrees of freedom; F, statistic; MS, mean square; Sig., significance level; SS, sum of squares.

male empathy scores.^{23–25 36–42} For example, a study has demonstrated statistical significance in female empathy than male empathy ($p < 0.001$).²⁴ In addition, women were reported to show increase in mean empathy score than male colleagues ($M=5.55$, $SD=0.46$ and $M=5.35$, $SD=0.55$,³⁹ respectively).

This has further been buttressed in another study where the mean female empathy score ($M=110.8$; $SD=11.7$) was reportedly higher than that of male empathy score ($M=105.3$; $SD=13.5$; $p=0.0001$; $d=0.44$).²⁵ In contrast, there have been reports of stability in empathy between women and men.⁴²

Despite the above evidence of empathy in some nursing research in the short term following empathy training, there have been some doubts on empathy follow-up research.^{37 43 44} In this study, empathy did not show any statistically significant difference between the groups in a follow-up after 6 months. This study is consistent with another study that found nursing empathy after training did not improve after five times measurement ($F(1, 29)=3.91$, $p < 0.06$).⁴³ This doubt in follow-up has also been reported in an earlier study by Daniels *et al.*⁴⁵

In contrast, another study found empathy increased 3 months after CST.⁴⁴ However, another study reported decreases in empathy as students advance through their nursing programme.⁴⁶

It has also been found by some researchers that there is a positive correlation between nursing students empathy and patient outcomes.^{47–50} Yu and Kirk,⁵¹ in a systematic review of measurement of empathy in nursing research, indicated that in eight appraisal researches, there was enhancement of empathy levels of students but that it was unclear if such enhancement was sustainable.

The results from this study confirm previous studies findings on nursing and midwifery training that empathy cannot be enhanced in a short period following CST.^{31–35 42 43} With this similar finding, there is a need for

further studies to determine the effectiveness of CST in enhancing NMS empathy.

Most of the studies have focused on empathy levels of nurses, differences in empathy, relationship between empathy and demographics variables.⁵¹ However, there are limited studies in the area of empathy in NMS. There are varied studies and the results from the previous studies show low,^{45 47} moderately enhanced^{23 52} and high levels^{23 34 52–56} of self-reported nurses' empathy. Other findings on nursing and midwifery training have contradicted this current study by indicating that empathy can be enhanced with training.^{24 25 41} However, some studies have found that NMS empathy actually decreases after training.^{46 57} Other variables like age, gender, education and religion have been considered in research.²³

Despite the fact that some studies have focused on empathy training among healthcare professionals including NMS in other countries, there are no known studies in Ghana. This study will therefore add to the literature on how best to enhance CST.

CONCLUSIONS

This study has shown that empathy may not be enhanced within a short period after CST. The participants were made aware of empathy being an outcome of this study and since JES is self-reported, it may have impacted their self-report. Selection bias may have impacted the lack of significance. It is possible that participants that volunteered were more empathetic compared with baseline and JES is self-reported. More so, the 2-day training time was not enough and that could have accounted for no enhancement of empathy.

This is the first RCT using CST in a nursing and midwifery school in Ghana. A study of this nature may better be evaluated by multicentre location in RCT across several regions. This may offer a much better comparison.

In this study, the women outnumbered the men both in the intervention and control. This is a limitation in the sense that women turn to be empathetic than men. Also, 99% of participants were either Christians or Muslims, as one is aware that religion teaches its members to be empathetic towards one another and this could have an effect on the outcome of this study.

Despite the limitations and strengths of this current study, the following recommendations are made for future studies. This CST had used a 2-day training period. A longer training period could have offered a better comparison. It does look like participants did not have the opportunity to read and reflect on the 2-day training before the post-test. This study used one location and a multicentre location in RCT across several nursing and midwifery schools probably could provide better outcomes.

This study explored the effect of CST post-test and 6 months post-training; however, a long-term examination could have been very useful. Further studies exploring the longer-term impact of the CST in other healthcare professionals and multi-location using cluster sampling may be beneficial. There is a need for additional studies to find out which aspects of CST for NMS will enhance empathy.

Acknowledgements The author (MA) wishes to thank the research assistant (AAM), management of Tamale Nursing and Midwifery college and the nursing and midwifery students for their participation in this study.

Contributors MA designed, undertook data collection, data integrity, accuracy of the analyses and writing of this article.

Funding The authors have not declared a specific grant for this research from any funding agency in the public, commercial or not-for-profit sectors.

Competing interests None declared.

Patient consent for publication Not required.

Ethics approval The Research and Monitoring Department of Tamale Teaching Hospital, Tamale – Ghana, gave Ethical approval for this study on 5th May 2013. The approval number is TTH/R6M/SR/13/12.

Provenance and peer review Not commissioned; externally peer reviewed.

Data sharing statement The data sets supporting the conclusions of this article are included within the article (extra data are available by emailing mustaph@uds.edu.gh).

Author note MA is a lecturer at the University for Development Studies, School of Allied Health Sciences, Tamale - Ghana.

Open access This is an open access article distributed in accordance with the Creative Commons Attribution Non Commercial (CC BY-NC 4.0) license, which permits others to distribute, remix, adapt, build upon this work non-commercially, and license their derivative works on different terms, provided the original work is properly cited, appropriate credit is given, any changes made indicated, and the use is non-commercial. See: <http://creativecommons.org/licenses/by-nc/4.0/>.

REFERENCES

- Reynolds WJ, Scott B. Do nurses and other professional helpers normally display much empathy? *J Adv Nurs* 2000;31:226–34.
- McCabe C, Timmins F. Communication skills for nursing practice. *Palgrave Macmillan* 2013.
- Reynolds WJ, Scott B, Jessiman WC. Empathy has not been measured in clients' terms or effectively taught: a review of the literature. *J Adv Nurs* 1999;30:1177–85.
- Hojat M, Gonnella JS, Nasca TJ, et al. Physician empathy: definition, components, measurement, and relationship to gender and specialty. *Am J Psychiatry* 2002;159:1563–9.
- Allgood MR. Chapter 17: Rethinking Empathy in Nursing Education: Shifting to a Developmental View. *Annu Rev Nurs Educ* 2015. <http://www.highbeam.com/doc/1P3-842569171.html> (accessed 2 Feb 2015).
- Frankel RM, Stein T. Getting the most out of the clinical encounter: the four habits model. *J Med Pract Manage* 2001;16:184–91.
- Stein T, Frankel RM, Krupat E. Enhancing clinician communication skills in a large healthcare organization: a longitudinal case study. *Patient Educ Couns* 2005;58:4–12.
- Roter DL. Physician/patient communication: transmission of information and patient effects. *Med State Med J* 1983;32:260–5.
- Roter DL. Patient question asking in physician-patient interaction. *Health Psychol* 1984;3:395–409.
- ACOG Committee Opinion No. 587: Effective patient-physician communication. *Obstet Gynecol* 2014;123:389–93.
- Rogers CR. Client-centred therapy: its current practice, implications and theory. *Houghton Mifflin* 1951.
- Egan G. The skilled helper: a problem-management and opportunity-development approach to helping. *Cengage Learning* 2013.
- McCormack B, McCance T. *Person-centred Nursing: Theory and Practice*: Wiley, 2010.
- Roter D, Hall J, Kern D, et al. Improving physicians' interviewing skills and reducing patients' em. 1975. <http://www.ncbi.nlm.nih.gov/pubmed/7677554> (accessed 1 Jan 2015).
- Faul F, Erdfelder E, Buchner A, et al. Statistical power analyses using G*Power 3.1: tests for correlation and regression analyses. *Behav Res Methods* 2009;41:1149–60.
- Universität Düsseldorf: G*Power. <http://www.gpower.hhu.de/> (accessed 3 Sep 2018).
- Hojat M, Mangione S, Nasca TJ, et al. The Jefferson scale of physician empathy: development and preliminary psychometric data. *Educ Psychol Meas* 2001;61:349–65.
- Krupat E, Frankel R, Stein T, et al. The Four Habits Coding Scheme: validation of an instrument to assess clinicians' communication behavior. *Patient Educ Couns* 2006;62:38–45.
- Shapiro SS, Wilk MB. An analysis of variance test for normality (complete samples). *Biometrika* 1965;52:591–611.
- Razali NM, Wah YB. Power comparisons of shapiro-wilk, kolmogorov-smirnov, lilliefors and anderson-darling tests. *J Stat Model Anal* 2011;2:21–33.
- Bland JM, Altman DG. Multiple significance tests: the Bonferroni method. *BMJ* 1995;310:170.
- Gulbrandsen P, Krupat E, Benth JS, et al. 'Four Habits' goes abroad: report from a pilot study in Norway. *Patient Educ Couns* 2008;72:388–93.
- Bailey S. Levels of empathy of critical care nurses. *Aust Crit Care* 1996;9:121–7.
- Ouzouni C, Nakakis K. An exploratory study of student nurses' empathy. *Health Sci J* 2012;6:534.
- Cutcliffe JR, Cassidy P. The development of empathy in students on a short, skills based counselling course: a pilot study. *Nurse Educ Today* 1999;19:250–7.
- Bas-Sarmiento P, Fernández-Gutiérrez M, Baena-Baños M, et al. Efficacy of empathy training in nursing students: a quasi-experimental study. *Nurse Educ Today* 2017;59:59–65.
- Moore PM, Rivera Mercado S, Grez Artigues M, et al. Communication skills training for healthcare professionals working with people who have cancer. *Cochrane Database Syst Rev* 2013;3:CD003751.
- Okaya K. Reliance between a patient and a nurse. *Nurs Today* 1995;10:6–11.
- Allgood MR. Empathy: the importance of recognizing two types. *J Psychosoc Nurs Ment Health Serv* 1992;30:14–17.
- La Monica EL, Wolf RM, Madea AR, et al. Empathy and nursing care outcomes. *Sch Inq Nurs Pract* 1987;1:197–213.
- Reynolds WJ, Presly AS. A study of empathy in student nurses. *Nurse Educ Today* 1988;8:123–30.
- Hodges SA. An experiment in the development of empathy in student nurses. *J Adv Nurs* 1991;16:1296–300.
- Williams B, Brown T, Boyle M, et al. Levels of empathy in undergraduate emergency health, nursing, and midwifery students: a longitudinal study. *Adv Med Educ Pract* 2014;5:299–306.
- Pålsson MB, Hallberg IR, Norberg A, et al. Burnout, empathy and sense of coherence among Swedish district nurses before and after systematic clinical supervision. *Scand J Caring Sci* 1996;10:19–26.
- Lauder W, Reynolds W, Smith A, et al. A comparison of therapeutic commitment, role support, role competency and empathy in three cohorts of nursing students. *J Psychiatr Ment Health Nurs* 2002;9:483–91.
- Chen D, Lew R, Hershman W, et al. A cross-sectional measurement of medical student empathy. *J Gen Intern Med* 2007;22:1434–8.

37. Hojat M, Gonnella JS, Mangione S, *et al.* Empathy in medical students as related to academic performance, clinical competence and gender. *Med Educ* 2002;36:522–7.
38. Kataoka HU, Koide N, Hojat M, *et al.* Measurement and correlates of empathy among female Japanese physicians. *BMC Med Educ* 2012;12:48.
39. Tavakol S, Dennick R, Tavakol M. Empathy in UK medical students: differences by gender, medical year and specialty interest. *Educ Prim Care* 2011;22:297–303.
40. DeVellis RF. Scale development: theory and applications. SAGE 2012.
41. Williams B, Brown T, McKenna L, *et al.* Empathy levels among health professional students: a cross-sectional study at two universities in Australia. *Adv Med Educ Pract* 2014;5:107–13.
42. Kliszcz J, Nowicka-Sauer K, Trzeciak B, *et al.* Empathy in health care providers—validation study of the Polish version of the Jefferson Scale of Empathy. *Adv Med Sci* 2006;51:219–25.
43. Evans GW, Wilt DL, Allgood MR, *et al.* Empathy: a study of two types. *Issues Ment Health Nurs* 1998;19:453–61.
44. Yates P, Hart G, Clinton M, *et al.* Exploring empathy as a variable in the evaluation of professional development programs for palliative care nurses. *Cancer Nurs* 1998;21:402–10.
45. Daniels TG, Denny A, Andrews D. Using microcounseling to teach RN nursing students skills of therapeutic communication. *J Nurs Educ* 1988;27:246–52.
46. Fallowfield L, Jenkins V, Farewell V, *et al.* Enduring impact of communication skills training: results of a 12-month follow-up. *Br J Cancer* 2003;89:1445–9.
47. Reid-Ponte P. Distress in cancer patients and primary nurses' empathy skills. *Cancer Nurs* 1992;15:283–92.
48. Murphy PA, Forrester DA, Price DM, *et al.* Empathy of intensive care nurses and critical care family needs assessment. *Heart Lung* 1992;21:25–30.
49. Olson JK. Relationships between nurse-expressed empathy, patient-perceived empathy and patient distress. *Image J Nurs Sch* 1995;27:317–22.
50. Olson J, Hanchett E. Nurse-expressed empathy, patient outcomes, and development of a middle-range theory. *Image J Nurs Sch* 1997;29:71–6.
51. Yu J, Kirk M. Measurement of empathy in nursing research: systematic review. *J Adv Nurs* 2008;64:440–54.
52. Watt-Watson J, Garfinkel P, Gallop R, *et al.* The impact of nurses' empathic responses on patients' pain management in acute care. *Nurs Res* 2000;49:191–200.
53. Aström S, Nilsson M, Norberg A, *et al.* Empathy, experience of burnout and attitudes towards demented patients among nursing staff in geriatric care. *J Adv Nurs* 1990;15:1236–44.
54. Aström S, Nilsson M, Norberg A, *et al.* Staff burnout in dementia care—relations to empathy and attitudes. *Int J Nurs Stud* 1991;28:65–75.
55. Warner RS. Nurses' empathy and patients' satisfaction with nursing care. *J N Y State Nurses Assoc* 1992;23:8–11.
56. Kuremyr D, Kihlgren M, Norberg A, *et al.* Emotional experiences, empathy and burnout among staff caring for demented patients at a collective living unit and a nursing home. *J Adv Nurs* 1994;19:670–9.
57. Kazanowski MK, Bennett LA. Engendering Empathy in Baccalaureate Nursing Students.