

BMJ Open Life-threatening alcohol-related traffic crashes in adverse weather: a double-matched case-control analysis from Canada

Donald A Redelmeier,¹ Fizza Manzoor

To cite: Redelmeier DA, Manzoor F. Life-threatening alcohol-related traffic crashes in adverse weather: a double-matched case-control analysis from Canada. *BMJ Open* 2019;**9**:e024415. doi:10.1136/bmjopen-2018-024415

► Prepublication history and additional material for this paper are available online. To view these files, please visit the journal online (<http://dx.doi.org/10.1136/bmjopen-2018-024415>).

Received 25 May 2018

Revised 15 November 2018

Accepted 21 January 2019



© Author(s) (or their employer(s)) 2019. Re-use permitted under CC BY-NC. No commercial re-use. See rights and permissions. Published by BMJ.

Medicine, Sunnybrook Health Sciences Centre, Toronto, Ontario, Canada

Correspondence to

Dr Donald A Redelmeier; dar@ices.on.ca

ABSTRACT

Importance Drunk driving is a major cause of death in North America, yet physicians rarely counsel patients on the risks of drinking and driving.

Objective To test whether the risks of a life-threatening alcohol-related traffic crash were further accentuated by adverse weather.

Design Double matched case-control analysis of hospitalised patients.

Setting Canada's largest trauma centre between 1 January 1995 and 1 January 2015.

Participants Patients hospitalised due to a life-threatening alcohol-related traffic crash.

Exposure Relative risk of a crash associated with adverse weather estimated by evaluating the weather at the place and time of the crash (cases) compared with the weather at the same place and time a week earlier and a week later (controls).

Results A total of 2088 patients were included, of whom the majority were drivers injured at night. Adverse weather prevailed among 312 alcohol-related crashes and was significantly more frequent compared with control circumstances. The relative risk of a life-threatening alcohol-related traffic crash was 19% higher during adverse weather compared with normal weather (95% CI: 5 to 35, $p=0.006$). The absolute increase in risk amounted to 43 additional crashes, extended to diverse groups of patients, applied during night-time and daytime, contributed to about 793 additional patient-days in hospital and was distinct from the risks for drivers who were negative for alcohol.

Conclusions Adverse weather was associated with an increased risk of a life-threatening alcohol-related traffic crash. An awareness of this risk might inform warnings to patients about traffic safety and counselling alternatives to drinking and driving.

INTRODUCTION

Alcohol-related traffic crashes cause substantial mortality and morbidity, accounting for ten thousand deaths annually in North America and contributing to one-third of total traffic fatalities.¹ Alcohol-related traffic fatality rates are higher in North America than many other countries that have greater

Strengths and limitations of this study

- Comprehensive analysis of patients hospitalised for life-threatening alcohol-related traffic crashes over two decades at Canada's largest trauma centre.
- Innovative case-only self-matched study design examining the crash location with the same time and place exactly 1 week earlier and 1 week later.
- Study limitations are inevitable because a randomised trial of drunk driving is not ethical, driving patterns vary in different regions and many important details were not available including traffic volumes, speeds, spacing and enforcement.
- Further limitations include uncertainties around the exact mechanisms of the increased risks and why the risks are distinct to drunk drivers.
- Additional limitations include a lack of data on other hazards and on the effectiveness of clinician counselling for mitigating traffic risks.

alcohol consumption per-capita.²⁻⁴ In addition, life-threatening alcohol-related traffic crashes in Canada and the USA result in over 300 000 patients hospitalised for brain trauma, spinal cord injuries, orthopaedic fractures or other non-fatal complications (leading to \$43 billion in societal costs annually).^{5,6} These patterns indicate that current public education, regulation and enforcement are insufficient for preventing drunk driving.^{7,8}

Motorists who drive drunk do so many times before attracting the attention of a healthcare provider.^{9,10} Epidemiological studies and statistical models estimate the average drunk driver needs to travel more than a million miles to cause one crash fatality.¹¹⁻¹³ This seemingly innocuous pattern tends to build a false sense of security from prior personal experiences; specifically, a mistaken belief that the individual can drive without incident if the road situation remains the same and free of other hazards.¹⁴ This faulty reasoning is particularly beguiling because alcohol is a

necessary but not a sufficient factor in triggering an alcohol-related traffic crash and because drunk drivers lack insights on how even minor hazards might precipitate a crash.¹⁵

One particularly common objective hazard is adverse weather that can create an extended disturbance for all who share the road.¹⁶ Adverse weather reduces visibility, decreases vehicle traction, creates visual glare, obscures reflective road markings and changes the patterns of vehicle cross traffic.^{17 18} Naturally, everyday driving entails an endless configuration of potential additional hazards that vary for each person and are easily forgotten after an uneventful trip.¹⁹ The unrecognised effect of these hazards, however, might create a fundamental mechanism explaining the complex link between drunk driving and traffic crashes. In this study, we test the association of adverse weather with the risk of a life-threatening alcohol-related traffic crash.

METHODS

Patient selection

We identified consecutive adults admitted to Canada's largest trauma centre, a tertiary care hospital that treats patients from crashes in the country's largest province.^{20–22} For a comprehensive analysis, we included all patients hospitalised for a crash (hereafter termed a life-threatening crash) including drivers, passengers or pedestrians since multiple individuals can be injured in traffic.^{23 24} We focused on those who tested positive for alcohol based on history, examination, assay or police report. Unclassified or atypical road incidents were excluded (eg, skateboard misadventures). Enrolment spanned from 1 January 1995 to 1 January 2015 yielding a complete sample for the two most recent available decades.²⁵

Clinical characteristics

We obtained clinical characteristics for patients based on hospital records using a standardised method validated in past research.²⁶ Information on the time, date and place of the crash was collected from paramedic reports if available and hospital records otherwise.²⁷ Information on patient age, sex, comorbidity, vital signs (after paramedic resuscitation), Injury Severity Score and Glasgow Coma Scale was based on chart review.^{28 29} Of note, alcohol testing was routine in hospital trauma protocols and did not require consent. Further clinical details included surgical procedures, intensive care unit admission, total length of stay and hospital mortality.³⁰ The available records lacked information on driver education, past infractions, addiction history, license suspensions, impact velocity, vehicle condition, distance travelled or intended destinations.

Crash setting

Data on crash locations spanned a wide diverse geographic area (1 million km²), were extracted in differing formats (street intersection, geographic coordinates, postal code)

and were subsequently transformed to exact geocodes for the crash site.^{31 32} Patients with missing or inexact crash locations were retained for analysis, denoted explicitly and subjected to sensitivity analysis. Geographic proximity to the trauma unit was estimated by Euclidean (straight-line) distance for those with known crash locations and by the median distance for those with missing or inexact crash locations. Crash time was recorded to the nearest hour to match the precision of standardised archived weather information.³³

Adverse weather

The official Canada Climate Data and Information Archive provided weather data indexed to date and hour, as validated in past research.³⁴ We focused on adverse conditions defined in the archive as rain, fog, drizzle, showers, snow, storms, freezing rain or freezing drizzle.^{35 36} All other conditions were defined as normal and included clear, mainly clear, mostly cloudy and cloudy. Daytime was crudely distinguished from night-time using simple thresholds of 07:00 and 19:00 hours.³⁷ We selected the weather station closest to the crash for patients with exact crash locations and the most central airport weather station for patients with inexact crash locations so no case was excluded (cases with inexact locations also subjected to sensitivity analysis).

Control comparisons

We identified two control days for each crash defined by the circumstances a week earlier and a week later (when presumably no other traffic crash was present).³⁸ A crash at midnight on 14 July 2011, for example, was compared with the same place at midnight on 7 July 2011 and 21 July 2011. This case-only design controlled for seasonal, daily and hourly trends; required no matching on individual patient characteristics; avoided ecological bias; and minimised multiple potential confounders including age, sex, genetics, personality, habits, education and road configuration.³⁹ The prevailing weather at the same time and place for crashes and control days was extracted in a blinded manner (no knowledge of outcome), with rare cases of missing weather data substituted by the immediately preceding hour so all comparisons were complete.

Statistical analysis

Our prespecified primary analysis involved a matched evaluation of individual cases comparing the prevalence of adverse weather on the crash day to the prevalence of adverse weather on the control days at the same time and place.⁴⁰ The relative risk of a crash associated with adverse weather was calculated using conditional logistic regression (accounting for 1:2 matching).^{41 42} Stratified analyses were conducted to further account for individual characteristics. Secondary analyses repeated the calculations for drivers who were negative for alcohol to check if the risks associated with adverse weather were distinct to drinking and driving. All estimates were calculated using exact 95% CIs and considered each patient a separate case.

Table 1 Patient characteristics

	Alcohol-positive	Alcohol-negative
	Patients (n=2088)	Patients (n=8111)
Age (years)		
<25	538 (26)	1698 (21)
25–44	968 (46)	2716 (33)
45–64	453 (22)	2244 (28)
≥65	129 (6)	1453 (18)
Male	1707 (82)	5191 (64)
Medical comorbidity*	732 (35)	2684 (33)
Protective device active†	772 (37)	4201 (52)
Abnormal vital signs‡	482 (23)	1465 (18)
Decreased Glasgow Coma Score§	549 (26)	1475 (18)
Position		
Driver	1362 (65)	5249 (65)
Passenger	294 (14)	1047 (13)
Pedestrian	432 (21)	1815 (22)
Night-time¶	1395 (67)	2599 (32)
Spring and summer	1184 (57)	4639 (57)
Weekend	944 (45)	2433 (30)
First decade**	1214 (58)	3642 (45)
Exact crash location	1341 (64)	4774 (59)
Injury Severity Score		
<15	644 (31)	2598 (32)
15–24	565 (27)	2318 (29)
25–34	441 (21)	1694 (21)
≥35	438 (21)	1501 (19)

Primary analysis based on alcohol-positive patients (alcohol-negative shown for context).

*Hypertension or diabetes most commonly.

†Denotes seatbelts or helmets.

‡Denotes hypotension (blood pressure <100), tachycardia (heart rate >120) or tachypnea (respiratory rate >25).

§Denotes decreased consciousness (value <15).

¶Night-time is 19:00 to 07:00 hours, daytime is 07:00 to 19:00 hours.

**First decade is 1995 to 2004, second decade is 2005 to 2014.

RESULTS

A total of 10 199 patients were injured because of a life-threatening traffic crash during the study, of whom 2088 (20%) tested positive for alcohol (exact concentrations unavailable for analysis). The majority of the alcohol-related crashes involved patients as drivers, most occurred at night and less than half used a seat-belt (table 1). Alcohol-related crashes were distributed throughout the year, although counts were marginally higher in the spring and summer months. As expected, alcohol-related crashes were more common on weekends than weekdays and slightly more numerous during the

Table 2 Acute medical care

	Alcohol-positive	Alcohol-negative
	Patients (n=2088)	Patients (n=8111)
Summary measure		
Ambulance transportation*	2030 (97)	7739 (95)
Blood transfusion†	693 (33)	2549 (31)
Surgery performed‡	1162 (56)	4406 (54)
Critical care admission§	1251 (60)	4105 (51)
Length of stay >7 days¶	1194 (57)	4630 (57)
Patient death**	174 (8)	815 (10)

*Manner of arrival to hospital.

†Denotes one or more units of red blood cell transfusions.

‡Defined as operating room procedure.

§Includes medical or surgical intensive care unit.

¶Interval in hospital from admission to discharge.

**Case fatality during index hospitalisation.

first decade than the second decade of the study. An exact crash location was identified for the majority of patients regardless of whether the crash was alcohol-related.

The average patient in an alcohol-related crash was a middle-aged adult with no medical comorbidity. Men were disproportionately involved, as were those younger than age 65 years and those who had not been wearing a seat-belt. Patients in an alcohol-related crash were no less seriously injured as measured by the distribution of abnormal vital signs, decreased Glasgow Coma Scale scores or Injury Severity Scale scores compared with patients who were negative for alcohol. Almost all patients were transported to hospital by ambulance (table 2). One-third of the patients required blood transfusions, over half required surgery and over half required a critical care admission. Ultimately, 174 patients died following an alcohol-related crash.

Overall, 312 of the 2088 (15%) alcohol-related crashes were characterised by adverse weather conditions. In contrast, 537 of the 4176 (13%) control days were characterised by adverse weather conditions at the same time and place (figure 1). The difference in prevailing weather equalled a 19% increased risk of a life-threatening alcohol-related traffic crash associated with adverse weather compared with normal weather (95% CI: 5 to 35, $p=0.006$). The absolute difference amounted to 43 additional life-threatening alcohol-related traffic crashes associated with adverse weather (one-in-seven of those observed). In contrast, drivers who were negative for alcohol showed no increased risk of a life-threatening traffic crash associated with adverse weather (estimate = $-1%$, 95% CI: $-11%$ to $+8%$, $p=0.704$).

The increased risk of a life-threatening alcohol-related traffic crash extended to diverse patient groups (figure 2). Drivers tended to predominate, yet the relative increase in risk associated with adverse weather also applied to




Figure 1 Venn diagram of summary data. Summary data based on 2088 alcohol-related life-threatening traffic crashes. Circles show counts of days with adverse weather at the time of the crash, at the control day 1 week before the crash and at the control day 1 week after the crash. For example, top circle indicates 312 of 2088 total crashes had adverse weather at the time and place of the crash. Main findings show disproportionate number of crash days with adverse weather compared with control days with adverse weather. OR indicates the relative frequency of adverse weather associated with a crash and is mathematically equal to the relative frequency of a crash associated with adverse weather (by the standard logic of case-control designs). OR calculated using exact methods that also account for matching in all triplets.

pedestrians and passengers (low counts). Similarly, night-time crashes were more numerous than daytime crashes, yet the increased relative risk associated with adverse weather applied regardless of the time of day. Analyses stratified by season, weekday, decade, crash location, age, sex, Injury Severity Scale scores and mortality all showed increased relative risks of an alcohol-related traffic crash associated with adverse weather and wide 95% CIs. No analysis showed the opposite pattern, no pairwise interaction term was statistically significant and all point-estimates overlapped the primary analysis.

The increased risk of a life-threatening alcohol-related traffic crash associated with adverse weather resulted in substantial inpatient hospital care. The mean length of hospital stay was similar for patients injured in adverse weather conditions compared with patients injured in normal weather conditions (figure 3). In total, the absolute increase in risk associated with adverse weather accounted for about 793 additional patient-days in hospital (online supplementary appendix). Similarly, the absolute increase in risk associated with adverse weather accounted for 52 additional surgical operations and 255




Figure 2 Analyses in patient subgroups. Forest plot showing relative risk of a life-threatening alcohol-related traffic crash associated with adverse weather. X-axis denotes relative risk with the null association indicated by a vertical line. Y-axis shows different analyses with full cohort analysis positioned at the top. Numbers enclosed by square brackets provide count of crashes with adverse weather and total sample size of cases in each subgroup. Solid circles indicate relative risk estimates and horizontal lines indicate 95% CIs. Values to the right of 1.00 denote increased risk and CIs that exclude 1.00 are statistically significant ($p < 0.05$). Findings show increased risk across diverse subgroups with all CIs overlapping the primary analysis.

additional patient-days in critical care (online supplementary appendix). The net economic consequences were equivalent to approximately \$1 million in additional economic costs (online supplementary appendix).

DISCUSSION

We studied about 2000 patients injured in a life-threatening alcohol-related traffic crash over twenty years in Canada. We found that adverse weather was prevalent in many cases, accounted for a further 19% increased relative risk and might explain one-in-seven life-threatening alcohol-related crashes. The increased risk associated with adverse weather extended to diverse patient groups, applied during night-time and daytime, and accounted for hundreds of additional patient-days in hospital. An increased relative risk of this magnitude is twice as large as driving without an air bag and an absolute risk of this magnitude is particularly important due to the high baseline risk of a traffic crash for all drunk drivers.⁴³⁻⁴⁶




Figure 3 Length of hospital stay. Kaplan-Meier graph of time spent in hospital by patients following 2088 alcohol-related life-threatening traffic crashes. X-axis shows time from admission in days. Y-axis shows proportion not yet discharged from hospital. Superimposed crossing lines denote crashes occurring in adverse weather and crashes occurring in normal weather conditions. Main findings show similar mean, median and distribution of length of hospital stay for both groups.

Several limitations of our findings merit emphasis. The study is not a randomised trial because the weather is impossible to control and drunk driving cannot be assigned in an ethical manner. Trauma centre medical assessments often underestimate the presence of alcohol, yet this imprecision tends to slant primary and stratified analyses towards the null.⁴⁷ Alcohol studies cannot be blinded easily so unconscious biases in clinicians might distort the assessment of outcome data.⁴⁸ Our analysis also lacks data on traffic volumes, distances, speeds, spacing and finer weather details at exact crash sites.⁴⁹ Finally, the background degree of traffic enforcement is uncertain, fluctuates on an hourly basis and stays confidential to avoid subterfuge.^{50 51}

Additional limitations relate to our study setting and justify replication in future research on other patients with life-threatening alcohol-related traffic crashes. The data reflect one large region that may not match drinking and driving patterns elsewhere.⁵² Canada is also notorious for long cold dark nights that are conducive to alcohol consumption and an impediment to roadside traffic police.^{53 54} Different regions have different traffic patterns and also vary in the age for legal drinking (19 years in most of Canada, 21 years in the USA).⁵⁵ Finally, we examined only one hazard that was objective and widespread yet daily traffic provides a large array of additional hazards that could precipitate a life-threatening alcohol-related crash.^{56–58}

Alcohol causes traffic crashes because of impaired judgement, decreased attention, reduced alertness and many other factors that limit the ability to compensate for hazards.⁵⁹ A further subtle mechanism is how alcohol lowers visual acuity when moving. Laboratory experiments indicate, for example, that three drinks of

alcohol cause a one-line loss on a Snellen eye chart test due to faulty visual tracking and reduced dynamic visual acuity.⁶⁰ Acute alcohol ingestion also causes decreased contrast sensitivity, sluggish glare adaptation and impaired risk perception that is unnoticed when stationary.^{61 62 63} The net effect is that eyesight deficits may be irrelevant when seated indoors (static vision), critical when driving in adverse weather (high-speed optical flow) and part of the false sense of security associated with drinking and driving.^{64 65}

Our study suggests adverse weather is directly relevant for alcohol-related crashes yet does not identify all the other hazards accentuating the traffic risks. Worsened surface glare, wheel traction and light backscatter may compromise how nearby traffic compensates for an impaired driver.⁶⁶ External sensors or other assistive vehicle technologies can malfunction when wet.⁶⁷ Traffic police may also dislike adverse weather and reduce enforcement in the rain.^{68 69} Together, these factors can help explain why adverse weather could be distinctly dangerous to drunk drivers; specifically, a crude OR of 25 associated with drunk driving might increase to a theoretical OR of 30 when driving in the rain (25×1.19). The net result could contribute to hundreds of patients requiring acute hospitalisation in North America each year (online supplementary appendix).⁷⁰

An increased risk of a life-threatening alcohol-related crash associated with adverse weather also means that police enforcement alone is not an easy solution against drunk driving.^{71 72} Traditional enforcement commonly includes sobriety checkpoints, mass media campaigns, encouraging seatbelts and random roadside alcohol breath testing.^{73 74} These interventions, of course, are effective and merit continuation.⁷⁵ In daily practice, however, the inconvenience of police enforcement (especially in adverse weather) helps explain the high ongoing rates of drunk driving in many countries.^{76 77} Clinicians wishing to save their patients from becoming more traffic injury statistics, therefore, might consider additional interventions beyond police enforcement.⁷⁸

Drunk driving is a life-threatening behaviour, yet little data are available to guide clinicians for prevention.⁷⁹ Current clinical efforts mostly include treating alcohol misuse or reporting unfit drivers to licensing authorities.^{80 81} Drunk driving, however, also stems from a patient's dismissal of health hazards that seem innocuous due to misleading past experiences. Clinicians intending to save patients from traffic injuries, therefore, might also counsel how crash risks vary substantially from one trip to the next due to adverse weather or other external hazards.⁸² Such counselling could mention a pre-planned taxi or ride-sharing option since both inebriation and uneventful past experiences can impair judgement.⁸³ Although not prescribed by a physician, this study suggests that alcohol is a drug that endangers patients and justifies tactical medical attention.

Patient involvement statement

Patients were not involved directly in the design, conduct or reporting of this research. We are grateful for all patients, families and clinicians involved in trauma care. We remain committed to disseminating study results to patients and the broad community.

Possible tweet

Adverse weather is associated with an increased risk of an alcohol-related traffic crash.

Acknowledgements We thank the following individuals for helpful comments on selected points: Allan Detsky, Michael Fralick, Saba Manzoor, Taaha Muhammad, William Raskin, Sheharyar Raza, Sharon Reece, Raffi Rush and Baiju Shah.

Contributors The lead author (DAR) wrote the first draft. Both authors (DAR, FM) contributed to study design, manuscript preparation, data analysis, results interpretation, critical revisions and final decision to submit. The lead author (DAR) had full access to all the data in the study, takes responsibility for the integrity of the data and is accountable for the accuracy of the analysis.

Funding This article was supported by a Canada Research Chair in Medical Decision Sciences, the Canadian Institutes of Health Research, the Ontario Ministry of Transportation and the Comprehensive Research Experience for Medical Students program at the University of Toronto.

Competing interests None declared.

Patient consent for publication Not required.

Ethics approval The study protocol was approved by the Research Ethics Board of Sunnybrook Health Sciences Center including a waiver for direct patient consent.

Provenance and peer review Not commissioned; externally peer reviewed.

Data sharing statement No additional data are available.

Open access This is an open access article distributed in accordance with the Creative Commons Attribution Non Commercial (CC BY-NC 4.0) license, which permits others to distribute, remix, adapt, build upon this work non-commercially, and license their derivative works on different terms, provided the original work is properly cited, appropriate credit is given, any changes made indicated, and the use is non-commercial. See: <http://creativecommons.org/licenses/by-nc/4.0/>.

REFERENCES

1. U.S. Department of Transportation, National Highway Traffic Safety Administration. National Highway Traffic Safety Administration. Traffic Safety Facts: Alcohol impaired driving. 2014 <http://www-nrd.nhtsa.dot.gov/Pubs/812231.pdf> (Cited 26 Jul 2016).
2. Sauber-Schatz EK, Ederer DJ, Dellinger AM, et al. Vital signs: motor vehicle injury prevention - United States and 19 comparison countries. *MMWR Morb Mortal Wkly Rep* 2016;65:672-7.
3. World Health Organization. *Recorded alcohol per capita (15+ years) consumption in litres of pure alcohol, from 1990*. Geneva: World Health Organization, 2016.
4. Fenelon A, Chen LH, Baker SP. Major causes of injury death and the life expectancy gap between the United States and other high-income Countries. *JAMA* 2016;315:609-11.
5. Majdan M, Plancikova D, Maas A, et al. Years of life lost due to traumatic brain injury in Europe: a cross-sectional analysis of 16 countries. *PLoS Med* 2017;14:e1002331.
6. Blincoc L, Miller TR, Zaloshnja E, et al. *The economic and societal impact of motor vehicle crashes, 2010 (Revised)*: U.S. Department of Transportation, National Highway Traffic Safety Administration, 2015.
7. Mello MJ, Nirenberg TD, Lindquist D, et al. Physicians' attitudes regarding reporting alcohol-impaired drivers. *Subst Abuse* 2003;24:233-42.
8. Spithoff S, Turner S. A systemic failure to address at-risk drinking and alcohol use disorders: the Canadian story. *CMAJ* 2015;187:479-80.
9. Zador PL, Krawchuk SA, Moore B. U.S. Department of Transportation, National Highway Traffic Safety Administration. Drinking and driving trips, stops by the police, and arrests: analyses of the 1995 National Survey of Drinking and Driving Attitudes and Behavior [Internet]. 2000 https://ntl.bts.gov/lib/26000/26600/26662/809_184.pdf (Cited 19 Jul 2017).
10. Redelmeier DA, Vinkatesh V, Stanbrook MB. Mandatory reporting by physicians of patients potentially unfit to drive. *Open Med* 2008;2:e8-e17.
11. Cavaioia AA, Strohmetsch DB, Abreo SD. Characteristics of DUI recidivists: a 12-year follow-up study of first time DUI offenders. *Addict Behav* 2007;32:855-61.
12. Solomon R, Chamberlain E, Abdoulaeva M, et al. Random breath testing: a Canadian perspective. *Traffic Inj Prev* 2011;12:111-9.
13. U.S. Department of Transportation: National Highway Traffic Safety Administration. National Highway Traffic Safety Administration. Traffic Safety Facts: 2014 data [Internet]. 2015 <https://crashstats.nhtsa.dot.gov/Api/Public/ViewPublication/812231> (Cited 18 Mar 2017).
14. Paxion J, Galy E, Berthelon C. Mental workload and driving. *Front Psychol* 2014;5:1344.
15. PLoS Medicine Editors. Preventing road deaths--time for data. *PLoS Med* 2010;7:e1000257.
16. Elvik R. Does the influence of risk factors on accident occurrence change over time? *Accid Anal Prev* 2016;91:91-102.
17. Hranac R, Sterzin E, Krechmer D, et al. *Empirical studies on traffic flow in inclement weather*. Washington (DC): U.S. Department of Transportation, National Highway Traffic Safety Administration, 2006.
18. Higgins L, Miles JD, Carlson P, et al. Nighttime visibility of prototype work zone markings under dry, wet-recovery, and rain conditions. *Transp Res Rec* 2009;2107:69-75.
19. Klauer SG, Guo F, Simons-Morton BG, et al. Distracted driving and risk of road crashes among novice and experienced drivers. *N Engl J Med* 2014;370:54-9.
20. McMurtry RY, Nelson WR, de la Roche MR. Current concepts in trauma: 2. The Sunnybrook Medical Centre trauma program: the first 11 years. *CMAJ* 1989;141:555-9.
21. Brenneman FD, Boulanger BR, McLellan BA, et al. Acute and long-term outcomes of extremely injured blunt trauma victims. *J Trauma* 1995;39:320-4.
22. Sunnybrook Health Sciences Centre. Facts & Figures. 2016 <http://sunnybrook.ca/content/?page=statistics> (Cited 19 July 2017).
23. Sunnybrook Health Sciences Centre. Sunnybrook Health Sciences Centre. Strategic Plan for Sunnybrook's Trauma Strategic Priority: 2015-2019 (Executive Overview). 2014 <https://sunnybrook.ca/uploads/1/programs/trauma-emergency-care/strategic-plan/trauma-strategic-plan-executive-overview.pdf> (Cited 13 Nov 2018).
24. Canadian Institute for Health Information. *Ontario Trauma Registry comprehensive data set data dictionary - May 2014*: Canadian Institute for Health Information, 2014. https://www.cihi.ca/en/services_otr_cds_dict_en.pdf. (Cited 13 Nov 2018).
25. Lustig AJ, Kurdyak PA, Thiruchelvam D, et al. Physician warnings in psychiatry and the risk of road trauma: an exposure crossover study. *J Clin Psychiatry* 2016;77:e1256-e1261.
26. Boulanger BR, McLellan BA, Sharkey PW, et al. A comparison between a Canadian regional trauma unit and an American level I trauma center. *J Trauma* 1993;35:261-6.
27. Redelmeier DA, Raza S. Life-threatening motor vehicle crashes in bright sunlight. *Medicine* 2017;96:e5710.
28. Kondziolka D, Schwartz ML, Walters BC, et al. The Sunnybrook neurotrauma assessment record: improving trauma data collection. *J Trauma* 1989;29:730-5.
29. Walters BC, McNeill I. Improving the record of patient assessment in the trauma room. *J Trauma* 1990;30:398-409.
30. Tien HC, Spencer F, Tremblay LN, et al. Preventable deaths from hemorrhage at a level I Canadian trauma center. *J Trauma* 2007;62:142-6.
31. Kelusky R. *2001 Geocode Street Guide*. Toronto: Toronto Works and Emergency Services, 2001.
32. Statistics Canada. *Land and freshwater area, by province and territory*: Statistics Canada, 2005.
33. Ahmed MM, Abdel-Aty M, Lee J, et al. Real-time assessment of fog-related crashes using airport weather data: a feasibility analysis. *Accid Anal Prev* 2014;72:309-17.
34. Fralick M, Denny CJ, Redelmeier DA. Drowning and the influence of hot weather. *PLoS One* 2013;8:e71689.
35. Meteorological Service of Canada. Glossary. Environment Canada; 1994-2014. http://climate.weather.gc.ca/glossary_e.html (Cited 26 Nov 2016).
36. Environment Canada. Manual of Surface Weather Observation. 2015 (Cited 26 Nov 2016).
37. Meteorological Service of Canada. Environment Canada; 1994-2014. http://climate.weather.gc.ca/historical_data/search_historic_data_e.html. (Cited 26 Nov 2016).
38. Redelmeier DA, Stewart CL. Driving fatalities on Super Bowl Sunday. *N Engl J Med* 2003;348:368-9.

39. Redelmeier DA, Tibshirani RJ. A simple method for analyzing matched designs with double controls: McNemar's test can be extended. *J Clin Epidemiol* 2017;81:51–5.
40. Redelmeier DA, Yarnell CJ. Can tax deadlines cause fatal mistakes? *Chance* 2013;26:8–14.
41. Hosmer D, Lemeshow S. *Applied logistic regression*. 2nd edn. New York (NY): Wiley and Sons, 2000:243–59.
42. Redelmeier DA, Tibshirani RJ. Methods for analyzing matched designs with double controls: excess risk is easily estimated and misinterpreted when evaluating traffic deaths. *J Clin Epidemiol* 2018;98:117–22.
43. Evans L. Traffic Safety. Bloomfield (MI): science serving society. 2004.
44. Zador PL, Krawchuk SA, Voas RB. Alcohol-related relative risk of driver fatalities and driver involvement in fatal crashes in relation to driver age and gender: an update using 1996 data. *J Stud Alcohol* 2000;61:387–95.
45. Cummings P, McKnight B, Rivara FP, et al. Association of driver air bags with driver fatality: a matched cohort study. *BMJ* 2002;324:1119–22.
46. Compton RP, Berning A. Drug and alcohol crash risk. *Journal of Drug Addiction, Education, and Eradication* 2015;11:29–46.
47. Gentilello LM, Villaveces A, Ries RR, et al. Detection of acute alcohol intoxication and chronic alcohol dependence by trauma center staff. *J Trauma* 1999;47:1131–5.
48. Colby SM, Barnett NP, Eaton CA, et al. Potential biases in case detection of alcohol involvement among adolescents in an emergency department. *Pediatr Emerg Care* 2002;18:350–4.
49. National Highway Traffic Safety Administration. *National motor vehicle crash causation survey: report to congress*. U.S. Department of Transportation: National Highway Traffic Safety Administration, 2008.
50. Fleming J. Learning to work together: police and academics. *Policing* 2010;4:139–45.
51. Ludwig A, Marshall M. Using crime data in academic research: issues of comparability and integrity. *Records Management Journal* 2015;25:228–47.
52. Sauber-Schatz EK, Ederer DJ, Dellinger AM, et al. Vital signs: motor vehicle injury prevention - united states and 19 comparison countries. *MMWR Morb Mortal Wkly Rep* 2016;65:672–7.
53. Li R, El-Basyouny K, Kim A. *A city-wide safety analysis of mobile speed enforcement: proceedings of the 25th CARSP conference: speeding and aggressive driving*. Ottawa (ON): Canadian Association of Road Safety Professionals, 2015.
54. Perreault S. *Impaired driving in Canada, 2015*. Ottawa (ON): Statistics Canada, 2016.
55. World Health Organization. *Global status report: alcohol policy*: World Health Organization, 2004.
56. Moskowitz H, Fiorentino D. *A review of the scientific literature regarding the effects of alcohol on driving-related behavior at blood alcohol concentrations of 0.08 grams per deciliter and lower*. U.S. Department of Transportation, National Highway Traffic Safety Administration, 2000.
57. Voas RB, Lacey JC. *Alcohol and highway safety 2006: a review of the state of knowledge*: U.S. Department of Transportation, National Highway Traffic Safety Administration, 2011.
58. Irwin C, Iudakhina E, Desbrow B, et al. Effects of acute alcohol consumption on measures of simulated driving: a systematic review and meta-analysis. *Accid Anal Prev* 2017;102:248–66.
59. Martin TL, Solbeck PA, Mayers DJ, et al. A review of alcohol-impaired driving: the role of blood alcohol concentration and complexity of the driving task. *J Forensic Sci* 2013;58:1238–50.
60. Schmäl F, Kunz R, Ortmann C, et al. Effect of ethanol on dynamic visual acuity during vertical body oscillation in healthy volunteers. *Eur Arch Otorhinolaryngol* 2000;257:485–9.
61. Nicholson ME, Andre JT, Tyrrell RA, et al. Effects of moderate dose alcohol on visual contrast sensitivity for stationary and moving targets. *J Stud Alcohol* 1995;56:261–6.
62. Andre JT. Visual functioning in challenging conditions: effects of alcohol consumption, luminance, stimulus motion, and glare on contrast sensitivity. *Journal of Experimental Psychology: Applied* 1996;2:250–69.
63. Deery HA, Love AW. The effect of a moderate dose of alcohol on the traffic hazard perception profile of young drink-drivers. *Addiction* 1996;91:815–27.
64. Lee DN. The optic flow field: the foundation of vision. *Philos Trans R Soc Lond B Biol Sci* 1980;290:169–79.
65. Bernardin F, Bremond R, Ledoux V, et al. Measuring the effect of the rainfall on the windshield in terms of visual performance. *Accid Anal Prev* 2014;63:83–8.
66. Qiu L, Nixon W. Effects of adverse weather on traffic crashes: systematic review and meta-analysis. *Transportation Research Record. Journal of the Transportation Research Board* 2008;2055:139–46.
67. Rasshofer RH, Spies M, Spies H. Influences of weather phenomena on automotive laser radar systems. *Advances in Radio Science* 2011;9:49–60.
68. Schafer JA, Mastrofski SD. Police leniency in traffic enforcement encounters: exploratory findings from observations and interviews. *J Crim Justice* 2005;33:225–38.
69. Chu HC. Risk factors for the severity of injury incurred in crashes involving on-duty police cars. *Traffic Inj Prev* 2016;17:495–501.
70. Voas RB, Torres P, Romano E, et al. Alcohol-related risk of driver fatalities: an update using 2007 data. *J Stud Alcohol Drugs* 2012;73:341–50.
71. Lerner BH. Drunk driving, distracted driving, moralism, and public health. *N Engl J Med* 2011;365:879–81.
72. Green RS, Kureshi N, Erdogan M. Legal consequences for alcohol-impaired drivers injured in motor vehicle collisions: A systematic review. *Accid Anal Prev* 2015;80:106–16.
73. Elder RW, Shults RA, Sleet DA, et al. Effectiveness of mass media campaigns for reducing drinking and driving and alcohol-involved crashes: a systematic review. *Am J Prev Med* 2004;27:57–65.
74. Bergen G, Pitan A, Qu S, et al. Publicized sobriety checkpoint programs: a community guide systematic review. *Am J Prev Med* 2014;46:529–39.
75. Naimi TS, Xuan Z, Sarda V, et al. Association of state alcohol policies with alcohol-related motor vehicle crash fatalities among US adults. *JAMA Intern Med* 2018;178:894.
76. Holder HD, Gruenewald PJ, Ponicki WR, et al. Effect of community-based interventions on high-risk drinking and alcohol-related injuries. *JAMA* 2000;284:2341–7.
77. Sloan FA, Eldred LM, Xu Y. The behavioral economics of drunk driving. *J Health Econ* 2014;35:64–81.
78. Rivara FP, Dunn C, Simpson E. *Addressing alcohol-impaired driving: training physicians to detect and counsel their patients who drink heavily*: U.S. Dept. of Transportation, National Highway Traffic Safety Administration, 2000.
79. Redelmeier DA, McLellan BA. Modern medicine is neglecting road traffic crashes. *PLoS Med* 2013;10:e1001463.
80. Redelmeier DA, Tien HC. Medical interventions to reduce motor vehicle collisions. *CMAJ* 2014;186:118–24.
81. Moyer A, Finney JW. Brief interventions for alcohol misuse. *CMAJ* 2015;187:502–6.
82. Redelmeier DA, Detsky AS. Clinical action against drunk driving. *PLoS Med* 2017;14:e1002231.
83. Decina LE, Foss R, Tucker ME, et al. *Alternative transportation programs: a countermeasure for reducing impaired driving*: U.S. Department of Transportation, National Highway Traffic Safety Administration, 2009.