BMJ Open Local prevalence of extended-spectrum beta-lactamase (ESBL) producing *Enterobacteriaceae* intestinal carriers at admission and co-expression of ESBL and OXA-48 carbapenemase in *Klebsiella pneumoniae*: a prevalence survey in a Spanish University Hospital

Cristina Díaz-Agero Pérez,^{1,2} Nieves López-Fresneña,^{1,2} Angela L Rincon Carlavilla,^{1,2} Marta Hernandez Garcia,^{2,3,4} Patricia Ruiz-Garbajosa,^{2,3,4} Jesús María Aranaz-Andrés,^{1,2} Friederike Maechler,⁵ Petra Gastmeier,⁶ Marc J M Bonten,⁷ Rafael Canton^{2,3,4}

To cite: Díaz-Agero Pérez C, López-Fresneña N, Rincon Carlavilla AL, *et al.* Local prevalence of extended-spectrum beta-lactamase (ESBL) producing *Enterobacteriaceae* intestinal carriers at admission and co-expression of ESBL and OXA-48 carbapenemase in *Klebsiella pneumoniae*: a prevalence survey in a Spanish University Hospital. *BMJ Open* 2019;**9**:e024879. doi:10.1136/bmjopen-2018-024879

► Prepublication history for this paper is available online. To view these files, please visit the journal online (<http://dx.doi.org/10.1136/bmjopen-2018-024879>).

Preliminary results from this study were presented at ECCMID 2016 on April 11, 2016, Amsterdam, The Netherlands.

Received 19 June 2018
Revised 28 November 2018
Accepted 9 January 2019

© Author(s) (or their employer(s)) 2019. Re-use permitted under CC BY-NC. No commercial re-use. See rights and permissions. Published by BMJ.

For numbered affiliations see end of article.

Correspondence to

Dr Cristina Díaz-Agero Pérez; cristina.diazagero@salud.madrid.org

ABSTRACT

Objective To assess the prevalence of extended-spectrum beta-lactamase (ESBL)-producing *Enterobacteriaceae* (ESBL-E) faecal carriers at admission in a University Hospital in Spain.

Design Prevalence survey.

Setting Pneumology, gastroenterology, urology and neurosurgery units at a university tertiary hospital in Madrid (Spain).

Participants A total of 10643 patients aged 18 and older admitted from March 2014 to April 2016 with a rectal swab taken at admission or as soon as possible within the first 48 hours.

Primary and secondary outcome measures Prevalence of ESBL-E faecal carriers and prevalence of ESBL-E infections at admission.

Results The prevalence of ESBL-E carriers at admission was 7.69% (CI 95% 7.18 to 8.19). Most of the isolates were *Escherichia coli* (77.51%), followed by *Klebsiella pneumoniae* (20.71%). Eighty-eight (10.41%) of ESBL-E were simultaneous ESBL and carbapenemase (CP) producers, 1.83% in the case of *E. coli* and 42.86% among *K. pneumoniae* isolates. Of the ESBL typed, 52.15% belonged to the cefotaximases (CTX-M-15) type and 91.38% of the CP were oxacillinase (OXA-48) type. Only 0.43% patients presented an active infection by ESBL-E at admission.

Conclusions The prevalence found in our study is very similar to that found in literature. However, we found a high percentage of simultaneous ESBL and CP producers, particularly in *K. pneumoniae*. Despite the high prevalence of colonised patients, the ESBL-infection rate at admission was very low.

BACKGROUND

The emergence of antimicrobial resistance represents a global challenge for healthcare

Strengths and limitations of this study

- This study is one of the most prolonged in time, with the largest number of patients, assessing colonisation by multidrug resistant micro-organisms. It includes adult participants of variable age groups and gender from a university hospital providing specialised assistance to 8.51% of the population of Madrid (Spain).
- The large number of patients included (10 643) gives strength to the results.
- Genes codifying extended-spectrum beta-lactamase (ESBL) and carbapenemase (CP) were characterised by polymerase chain reaction (PCR) and sequencing. Unfortunately total characterisation was not feasible in all isolates but only in 24.67% of total ESBL producing isolates and 73.86% of total CP producing isolates.

due to the limited treatment options. Extended-spectrum beta-lactamases (ESBL) are the main mechanisms of acquired resistance in Gram-negative bacteria. Until the late 90s, most ESBLs were isolated in nosocomial outbreaks, their prevalence was higher in *Klebsiella pneumoniae* than in *Escherichia coli*, and there was significant variation among countries, hospitals and wards.^{1 2} They were isolated at higher frequency in the intensive care units (ICU); recent surgery, catheterisation, urinary catheterisation, prolonged hospitalisation, ICU admission and previous use of cephalosporins and aminoglycosides were leading risk factors.^{3 4}

The situation today is very different since their prevalence has increased dramatically in the community, especially in urinary tract infections, where these enzymes are more frequently isolated in *E. coli*.^{5–8} The main clinical relevance of ESBL seems to be the inadequate empirical treatment delaying efficient antimicrobial treatment, for example, up to six times in the case of *E. coli* and *K. pneumoniae* ESBL (ie, 72 hours instead of 11 hours for susceptible strains).^{9 10} It is necessary to know the prevalence of microbial resistance in our geographical area and their epidemiological characteristics in order to establish the scope of the problem and analyse its evolution.

The aim of this study was to assess the prevalence of ESBL-producing *Enterobacteriaceae* (ESBL-E) faecal carriers at admission in hospital wards during an active surveillance screening programme (R-GNOSIS project).

METHODS

Study design and settings

The project falls within the R-GNOSIS study (Resistance of Gram-Negative Organisms: Studying Intervention Strategies) within the Work Package 5 "Patient isolation strategies for ESBL carriers in medical and surgical hospital wards", funded by the EU (FP7-HEALTH-2011-SINGLE STAGE-N°282512).

The University Hospital Ramón y Cajal is a public referral centre located in the North of Madrid (Spain). It provides specialised assistance to 558 373 citizens who represent 8.51% of the population of Madrid. With 1118 beds, it accounted for 31 179 admissions in the year 2014, 31 253 in 2015 and 31 847 in 2016. The pneumology (41 beds), gastroenterology (40 beds), urology (41 beds) and neurosurgery (20 beds) wards took part in the study.

Patients

Between 3 March 2014 and 3 April 2016, screening rectal swabs were obtained after verbal consent from all patients aged 18 and older, at admission or as soon as possible within the first 48 hours.

Patient involvement

Patients were not directly involved in the design and conception of the study. All patients were informed of the aim of the study and the consequences of a positive result (contact isolation and need for rectal screening at any hospital admission in the future to check the status) and gave their verbal consent to participate; if the patient refused, the swab was not taken. As soon as the microbiological result was known by the investigators, patients and their families were informed.

Laboratory analysis

The samples were seeded on ChromoID-ESBL and Chromo-ID CARBA/OXA-48 (BioMérieux, France) selective chromogenic agar plates. Bacterial identification was performed using the MALDI-TOF-MS (Bruker-Daltonics, Germany) mass spectrometry. ESBL and carbapenemase

(CP) production were phenotypically confirmed by the double-disk diffusion test, Hodge test and KPC/MBL/OXA-48 Confirm and ESBL AmpC screen kits (Rosco Diagnostica, Germany). Antimicrobial susceptibility was studied with microdilution (MicroScan, Beckman, CA) and gradient strips (Etests, BioMérieux, France). Genes codifying ESBL (*bla*_{SHV}, *bla*_{TEM}, *bla*_{CTX-M}) and CP (*bla*_{VIM}, *bla*_{KPC}, *bla*_{NDM}, *bla*_{OXA-48}) were characterised by PCR and sequencing.

Ethics

The study was carried out in accordance with the Declaration of Helsinki and Good Clinical Practice Guidelines (ICH-GCP-Guidelines, CPM/ICH/135/95) of the European Medicines Agency.

The study included all standard safeguards for ensuring the confidentiality of patient information and specifications stipulated in the Personal Data Protection Act 15/1999, of 13 December were followed.

Statistical analyses

A descriptive analysis of the variables collected was conducted; qualitative variables were expressed as percentages and quantitative variables as measures of central tendency (mean and median) and dispersion (SD and IQR). Pearson's χ^2 test was used to compare proportions and the Student's T-test was used to compare means. All statistical analyses were performed using SPSS Statistics V.19 (IBM) software.

RESULTS

During the research period 12 590 admissions of 9706 patients took place in the participating wards. In 84.5% of admissions, a rectal swab could be obtained within the first 48 hours of admission. [table 1](#).

Gender and mean age of included patients are shown in [table 2](#).

The prevalence of ESBL-E faecal carriers at admission was 7.69% ([table 3](#)). [Table 3](#) shows the distribution of carriers by ward, gender and age (mean and median).

Majority of patients colonised with ESBL-E were men, just like the majority of hospital patients, the difference not being statistically significant. The mean age of colonised patients was higher than the mean age of the

Table 1 Patients admitted to gastroenterology, pneumology, urology and neurosurgery wards and swabs taken

Ward	Admissions (n)	Swab at admission (n)	%
Gastroenterology	3380	2916	86.27
Pneumology	3240	2752	84.94
Urology	4685	3963	84.59
Neurosurgery	1285	1012	78.75
Total	12 590	10 643	84.55

Table 2 Gender and age of the included patients

Ward	Gender		Age (years)	
	Men (%)	Women (%)	Mean (SD)	Median (IQR)
Gastroenterology	1732 (59.39)	1184 (40.61)	66.53 (16.59)	69 (26.75)
Pneumology	1625 (59.05)	1127 (40.95)	70.72 (15.28)	74 (19)
Urology	3009 (75.93)	954 (24.07)	66.89 (14.56)	69 (20)
Neurosurgery	533 (52.67)	479 (47.33)	60.23 (16.52)	61 (25)
Total	6899 (64.82)	3744 (35.18)	64.91 (16.79)	67 (25)

total number of hospitalised patients (69.27SD 15.68 vs 64.91,SD 16.79), the difference being statistically significant ($p=0.0087$).

The difference in prevalence of colonisation at admission among the surveyed wards was statistically significant ($p=0.001$). The highest prevalence was found in the gastroenterology ward with 9.02%, the difference being significant when compared with the rest of wards ($p=0.01$). When comparing the prevalence between medical wards (pneumology and gastroenterology) and surgical wards (urology and neurosurgery), the difference was not statistically significant.

A total of 843 multiresistant *Enterobacteriaceae* were isolated in 818 patients, as 25 patients were colonised by more than one micro-organism at the time of admission (0.23%). Eighty-eight (10.44%) of the isolated *Enterobacteriaceae* were simultaneous ESBL and CP producers and 33.99% of these patients were known carriers, that is, their clinical records included a previous positive culture for ESBL-E.

The most frequently isolated ESBL-producing micro-organisms at admission were *E. coli* (77.70%; $n=655$), followed by *K. pneumoniae* (20.64%, $n=174$), with other species (*Enterobacter cloacae* 0.59%; *Citrobacter freundii* 0.36%; *Enterobacter aerogenes* 0.24%; *Citrobacter amalonaticus* 0.12%; *Citrobacter koseri* 0.12%; *Enterobacter asburiae* 0.12%; *Klebsiella oxytoca* 0.12%) being only 1.66 %. Among ESBL-*E. coli* isolates, 1.83% were simultaneous ESBL and CP producers ($n=12$). Among ESBL *K. pneumoniae* isolates, 43.10% were simultaneous ESBL and CP producers ($n=75$). Only one patient was colonised by a different ESBL and CP producer, *K. oxytoca*.

The typing of 208 beta-lactamases (24.67% of total ESBL) and 65 CP was possible (73.86% of total CP). Most of ESBL (83.17%) belonged to the CTX-M group, CTX-M-15 being the most numerous, followed by CTX-M-14. The remaining 16.83% belonged to the SHV group, SHV-12 being the most frequent (table 4). For the typed CP, 90.77% were OXA-48 type (table 5). In the case of four patients colonised simultaneously by two different ESBL-E (in two patients ESBL *E. coli* and ESBL *K. pneumoniae* and in the other ESBL+CP *E. coli* and ESBL+CP *K. pneumoniae* respectively), both micro-organisms carried the same enzyme type, CTX-M-15 in 3 of them and CTX-M-14 in 1 and OXA-48 in the case of CP.

Fifty-four patients presented an active infection by ESBL-E at admission, that is, 0.43% of patients admitted during the research period and 6.6% of ESBL-E intestinal carriers. Of those 54 patients, all except one also showed a positive rectal swab, 90.74% of those (49 patients) with the same specie causing the infection and 9.26% (five patients) with a different ESBL-E. Out of the diagnosed infections, 69.09% (38 urine cultures) were urinary tract infections, 14.55% bacteraemia ($n=8$; 1 of them secondary to a urinary tract infection), two community acquired pneumonias (3.64%), two surgical site infections (3.64%), two abscesses (3.64%), one lower respiratory infection (1.82%), one gastrostomy insertion site infection (1.82%) and one Fournier's gangrene (1.82%).

A total of 56 micro-organisms were isolated in the 55 positive clinical cultures as one of them was positive for two ESBL-E. The most frequently isolated micro-organism was, once again, *E. coli* (67.86%), followed by ESBL and

Table 3 ESBL-producing *Enterobacteriaceae* carriers at admission

Hospital admission wards	Gender		Age (years)		Prevalence (%) CI 95%
	Men (%)	Women (%)	Mean (SD)	Median (IQR)	
Gastroenterology	159 (60.23)	104 (39.77)	66.78 (16.62)	67.2 (26.64)	9.02 (7.96–10.08)
Pneumology	122 (61.31)	77 (38.69)	74.78 (14.36)	79 (15)	7.23 (6.25–8.22)
Urology	234 (80.69)	56 (19.31)	69.82 (14.04)	72 (21)	7.32 (6.49–8.14)
Neurosurgery	44 (66.67)	22 (33.33)	62.45 (17.26)	66.67 (25.84)	6.52 (4.95–8.09)
Total	559 (68.34)	259 (31.66)	69.27 (15.68)	72 (25)	7.69 (7.18–8.19)

ESBL, extended-spectrum beta-lactamases.

Table 4 Distribution of ESBL strains isolated and typed in rectal swabs at hospital admission

Enzyme	Micro-organism						Total (%)
	ESBL <i>Escherichia coli</i>	ESBL <i>Klebsiella pneumoniae</i>	ESBL <i>Enterobacter cloacae</i>	ESBL <i>Citrobacter freundii</i>	ESBL+CP <i>E. coli</i>	ESBL+CP <i>K. pneumoniae</i>	
CTX-M	1	–	–	–	–	–	1 (0.48)
CTX-M-1	10	4	–	–	–	–	14 (6.73)
CTX-M-9	10	3	–	–	–	–	13 (6.25)
CTX-M-14	23	1	–	–	–	2	26 (12.50)
CTX-M-15	35	31	1	–	3	40	110 (52.88)
CTX-M-27	6	–	–	–	–	–	6 (2.88)
CTX-M-32	2	–	–	–	–	–	2 (0.96)
CTX-M-55	1	–	–	–	–	–	1 (0.48)
SHV	1	1	–	–	–	–	2 (0.96)
SHV-2	1	1	–	–	–	–	2 (0.96)
SHV-12	10	8	–	1	–	5	24 (11.54)
SHV-28	–	5	–	–	–	1	6 (2.88)
SHV-31	–	1	–	–	–	–	1 (0.48)
Total	100	55	1	1	3	48	208 (100)

CP, carbapenemase; ESBL, extended-spectrum beta-lactamases.

CP *K. pneumoniae* (23.21%), ESBL *K. pneumoniae* (7.14%); *K. oxytoca* was isolated in one culture (1.79%).

DISCUSSION

In our study, the prevalence of ESBL-E carriers at admission was 7.69%, ranging between 6.52% and 9.02% depending on the ward. The prevalence of ESBL-E carriers in healthy individuals as well as in ambulatory and hospitalised patients has been researched in a number of studies. In all of them, *E. coli* is always the most frequently isolated micro-organism, as in our study (77.70%).^{11–19} In a meta-analysis published in 2016 which analysed prevalence studies in healthy persons and included 28 909 individuals from 66 studies, the mean global prevalence of colonisation was 14%, with great variability among regions.¹⁹ It was higher in Asia (with 46%) and Africa

(with 22%); in Europe the mean prevalence was 4%, with 3% in central Europe, 4% in northern Europe and 6% in southern Europe. Finally, in America, the mean prevalence was 2%, although it was admitted that there were very few studies for this region.²⁰

Our prevalence of intestinal carriers at admission is virtually the same as that found by a Dutch study published recently, which was 7.9% in patients coming from their homes and 8.6% in patients coming from long-term care facilities, a distinction not made in our research.²¹ Studies in three different areas in Spain (Madrid, Barcelona and Zaragoza) show that the prevalence of carriers has increased in the last few years, reaching rates ranging from 5.5% and 8.1% in 2002 and 2004 respectively, as in our study findings.^{11 13 16} In another study performed in Seville, the prevalence of carriers among patients admitted to emergency units was 7.4%, also very similar to our figure.²²

In our facility, 10.44% of ESBL micro-organisms were simultaneous CP producers, with 85.22% being *K. pneumoniae*, 13.64% *E. coli* and 1.14% *K. oxytoca*. Of the 65 CP typed (73.86% of total CP), the vast majority of them (90.77%) belonged to the OXA-48 type. This fact is especially important in the case of *K. pneumoniae*, with 43.10% being ESBL and CP producers (90.57% OXA-48). ESBL and CP *K. pneumoniae* were responsible for 23.21% of the infections diagnosed at hospital admission (69.27% of them urinary tract infections). We did not find a similar study to compare our data with but we think this finding must be deeply analysed. We found one case of KPC-3 (ESBL+CP *E. coli*) and one case of NDM-1 (ESBL+CP *K. pneumoniae*).

Table 5 Distribution of carbapenemase strains isolated and typed in rectal swabs at hospital admission

Enzyme	Micro-organism		
	ESBL+CP <i>Escherichia coli</i>	ESBL+CP <i>Klebsiella pneumoniae</i>	Total (%)
KPC-3	1	–	1 (1.54)
NDM-1	–	1	1 (1.54)
OXA-48	11	48	59 (90.77)
VIM-1	–	4	4 (6.15)
Total	12	53	65 (100)

CP, carbapenemase; ESBL, extended-spectrum beta-lactamases.

Male gender has been identified as a risk factor for the intestinal colonisation by ESBL-E.^{7 20 21 23 24} In our study, as in Valverde *et al*, the majority of colonised patients were men, but they were also the majority of the total number of hospitalised patients, the difference not being statistically significant.¹¹ Age is another risk factor identified in the bibliography; in our study, the mean age of colonised patients was higher than the mean age of hospitalised patients (69.27 years vs 64.91 years), the difference being statistically significant in this case ($p=0.0087$).^{23 24}

The prevalence of carriers at admission was higher in the gastroenterology ward, despite the patients being younger than the mean, with a difference being statistically significant when compared with the rest of included wards. In other published studies, liver disease has been identified as a risk factor for intestinal colonisation by ESBL-E, one of the possible explanations being the prophylactic use of fluoroquinolones to prevent spontaneous bacterial peritonitis in patients with chronic liver disease.^{25 26} Another risk factor for ESBL-E carriage recently described in the literature is the use of proton pump inhibitors (PPI) and these type of patients often receive PPIs and other medication for gastro-oesophageal reflux disease.^{27 28} In our case, we cannot provide an explanation as risk factors for every patient were not recorded.

Unfortunately total characterisation was not feasible in all isolates due to budget issues; so we decided to analyse a random selection. We were able to determine 24.67% of total ESBL producing isolates; that low percentage is a limitation of our study and the results could differ if all the ESBLs had been analysed but they are compatible with the epidemiology described in literature. The main enzyme group was CTX-M, the most common according to literature, followed by SHV, with the CTX-M-15 group prevailing with 52.88%.^{8 12-14 19 21 22 24}

In the last years, ESBL-E infections have become an increasing concern; in the USA for example, 140 000 hospital-acquired ESBL-E infections are estimated to occur per year.²⁹ Infections by these bacteria are associated with higher mortality rates and higher hospital costs compared with antibiotic-sensitive micro-organisms.³⁰ However, few studies have associated the fact of being an intestinal carrier of ESBL-E with the development of infections caused by these bacteria. A recent cohort study performed in patients with haematological malignancies found a 3.5-fold greater risk of developing bacteraemia by ESBL-E among colonised patients when compared with non-colonised patients; despite the fact that mortality was similar in both groups, colonisation was associated with longer hospital stays, shorter survival period and higher costs.³¹ On the contrary, another similar study did not find correlation between ESBL-E colonisation and infection in neutropenic patients.³² In our study 55 ESBL-E infections were diagnosed at admission and almost 70% were urinary tract infections. This means that 0.43% of patients were admitted with an ESBL-E infection, which represents 6.59% of the colonised patients. Only in one patient with ESBL-E infection at admission, no ESBL-E

was isolated in the rectal swabs. Even though the vast majority of infections were found in colonised patients, the total prevalence of infection was very low, and only in eight cases it consisted of bacteraemia (one of those secondary to a urinary tract infection). In two cases patients died during hospital admission, although their infection had been fully resolved and death was caused by an underlying oncological disease.

This study, one of the most prolonged in time and with the largest number of patients, confirms once again the extension of ESBL-E intestinal colonisation in the community, showing however, a low prevalence of infection. It is necessary to continue with the epidemiological surveillance of these micro-organisms, in order to acquire a better knowledge of the implications of being an intestinal carrier of ESBL-E. The high percentage of ESBL and CP *K. pneumoniae* producers must also be more deeply studied.

Author affiliations

¹Servicio de Medicina Preventiva y Salud Pública, Hospital Universitario Ramón y Cajal, Madrid, Spain

²Instituto Ramón y Cajal de Investigación Sanitaria (IRYCIS), Madrid, Spain

³Servicio de Microbiología, Hospital Universitario Ramón y Cajal, Madrid, Spain

⁴Red Española de Investigación en Patología Infecciosa (REIPI), Madrid, Spain

⁵Charité Universitätsmedizin Berlin, Berlin, Germany

⁶Infection Control, Charité, Berlin, Germany

⁷Department of Medical Microbiology, University Medical Center Utrecht, Utrecht, The Netherlands

Contributors Conception and design of study: MJMB, RC, PG, FM. Data collection: CD-AP, NL-F, ALR-C, MH-G, PR-G. Analysis and/or interpretation of data: CD-AP, NL-F, ALR-C, MH-G, PR-G, JMA-A, RC. Drafting the manuscript: CD-AP, NL-F, ALR-C, MH-G. Revising the manuscript critically for important intellectual content: RC, PR-G, JMA-A, MB, PG, FM. Approval of the version of the manuscript to be published: CD-AP, NL-F, ALR-C, MH-G, PR-G, JMA-A, FM, PG, MB, RC.

Funding The project falls within the R-GNOSIS study (Resistance of Gram-Negative Organisms: Studying Intervention Strategies), within the Work Package 5 Patient isolation strategies for ESBL carriers in medical and surgical hospital wards, funded by the European Union (FP7-HEALTH-2011-SINGLE STAGE-N°282512). MH-G is supported with a contract from Instituto de Salud Carlos III of Spain (iP-FIS program, ref. IF14/00022).

Competing interests None declared.

Patient consent for publication Not required.

Ethics approval The Ethics Committee of Clinical Research (Comité Ético de Investigación Clínica del Hospital Universitario Ramón y Cajal, Madrid, Spain) formally reviewed and approved the study protocol on October 2013 (Ref. 251 – 13). A waiver of written informed consent of individual patients in the participating wards was requested and granted by the Committee as well as by the Medical Direction since the study did not expose patients to any novel risk and no investigational drugs, devices, or procedures were involved and verbal consent was considered sufficient.

Provenance and peer review Not commissioned; externally peer reviewed.

Data sharing statement Extra data are available by emailing: cristina.diazagero@salud.madrid.org.

Open access This is an open access article distributed in accordance with the Creative Commons Attribution Non Commercial (CC BY-NC 4.0) license, which permits others to distribute, remix, adapt, build upon this work non-commercially, and license their derivative works on different terms, provided the original work is properly cited, appropriate credit is given, any changes made indicated, and the use is non-commercial. See: <http://creativecommons.org/licenses/by-nc/4.0/>.

REFERENCES

- Winokur PL, Canton R, Casellas JM, *et al.* Variations in the prevalence of strains expressing an extended-spectrum beta-lactamase phenotype and characterization of isolates from Europe, the Americas, and the Western Pacific region. *Clin Infect Dis* 2001;32:S94–103.
- Babini GS, Livermore DM. Antimicrobial resistance amongst *Klebsiella* spp. collected from intensive care units in Southern and Western Europe in 1997–1998. *J Antimicrob Chemother* 2000;45:183–9.
- Paterson DL, Bonomo RA. Extended-spectrum beta-lactamases: a clinical update. *Clin Microbiol Rev* 2005;18:657–86.
- Sirof D. Extended-spectrum plasmid-mediated beta-lactamases. *J Antimicrob Chemother* 1995;36:19–34.
- Pitout JD, Hanson ND, Church DL, *et al.* Population-based laboratory surveillance for *Escherichia coli*-producing extended-spectrum beta-lactamases: importance of community isolates with blaCTX-M genes. *Clin Infect Dis* 2004;38:1736–41.
- Rodríguez-Baño J, Navarro MD, Romero L, *et al.* Bacteremia due to extended-spectrum beta-lactamase-producing *Escherichia coli* in the CTX-M era: a new clinical challenge. *Clin Infect Dis* 2006;43:1407–14.
- Ben-Ami R, Schwaber MJ, Navon-Venezia S, *et al.* Influx of extended-spectrum beta-lactamase-producing enterobacteriaceae into the hospital. *Clin Infect Dis* 2006;42:925–34.
- Woerther PL, Burdet C, Chachaty E, *et al.* Trends in human fecal carriage of extended-spectrum β -lactamases in the community: toward the globalization of CTX-M. *Clin Microbiol Rev* 2013;26:744–58.
- García E. Bacteriemias por *Escherichia coli* productor de betalactamasas de espectro extendido (BLEE): Significación clínica y perspectivas actuales. España. *Revista Española de Quimioterapia* 2011;24:57–66.
- Lautenbach E, Patel JB, Bilker WB, *et al.* Extended-spectrum beta-lactamase-producing *Escherichia coli* and *Klebsiella pneumoniae*: risk factors for infection and impact of resistance on outcomes. *Clin Infect Dis* 2001;32:1162–71.
- Valverde A, Coque TM, Sánchez-Moreno MP, *et al.* Dramatic increase in prevalence of fecal carriage of extended-spectrum beta-lactamase-producing *Enterobacteriaceae* during nonoutbreak situations in Spain. *J Clin Microbiol* 2004;42:4769–75.
- Munday CJ, Whitehead GM, Todd NJ, *et al.* Predominance and genetic diversity of community- and hospital-acquired CTX-M extended-spectrum beta-lactamases in York, UK. *J Antimicrob Chemother* 2004;54:628–33.
- Miró E, Mirelis B, Navarro F, *et al.* Surveillance of extended-spectrum beta-lactamases from clinical samples and faecal carriers in Barcelona, Spain. *J Antimicrob Chemother* 2005;56:1152–5.
- Moubareck C, Daoud Z, Hakimé NI, *et al.* Countrywide spread of community- and hospital-acquired extended-spectrum beta-lactamase (CTX-M-15)-producing *Enterobacteriaceae* in Lebanon. *J Clin Microbiol* 2005;43:3309–13.
- Rodrigues C, Shukla U, Jog S, *et al.* Extended-spectrum β -Lactamase-producing Flora in Healthy Persons. *Emerg Infect Dis* 2005;11:981–2.
- Castillo García FJ, Seral García C, Pardos De la Gandara M, *et al.* Prevalence of fecal carriage of ESBL-producing *Enterobacteriaceae* in hospitalized and ambulatory patients during two non-outbreak periods. *Eur J Clin Microbiol Infect Dis* 2007;26:77–8.
- Kader AA, Kumar A, Kamath KA. Fecal carriage of extended-spectrum beta-lactamase-producing *Escherichia coli* and *Klebsiella pneumoniae* in patients and asymptomatic healthy individuals. *Infect Control Hosp Epidemiol* 2007;28:1114–6.
- Pallecchi L, Bartoloni A, Fiorelli C, *et al.* Rapid dissemination and diversity of CTX-M extended-spectrum beta-lactamase genes in commensal *Escherichia coli* isolates from healthy children from low-resource settings in Latin America. *Antimicrob Agents Chemother* 2007;51:2720–5.
- Karanika S, Karantanos T, Arvanitis M, *et al.* Fecal colonization with extended-spectrum beta-lactamase-producing *enterobacteriaceae* and risk factors among healthy individuals: a systematic review and metaanalysis. *Clin Infect Dis* 2016;63:310–8.
- Shitrit P, Reisfeld S, Paitan Y, *et al.* Extended-spectrum beta-lactamase-producing *Enterobacteriaceae* carriage upon hospital admission: prevalence and risk factors. *J Hosp Infect* 2013;85:230–2.
- Platteel TN, Leverstein-van Hall MA, Cohen Stuart JW, *et al.* Predicting carriage with extended-spectrum beta-lactamase-producing bacteria at hospital admission: a cross-sectional study. *Clin Microbiol Infect* 2015;21:141–6.
- Rodríguez-Baño J, López-Cerero L, Navarro MD, *et al.* Faecal carriage of extended-spectrum beta-lactamase-producing *Escherichia coli*: prevalence, risk factors and molecular epidemiology. *J Antimicrob Chemother* 2008;62:1142–9.
- Ruppé E, Pitsch A, Tubach F, *et al.* Clinical predictive values of extended-spectrum beta-lactamase carriage in patients admitted to medical wards. *Eur J Clin Microbiol Infect Dis* 2012;31:319–25.
- Ben-Ami R, Rodríguez-Baño J, Arslan H, *et al.* A multinational survey of risk factors for infection with extended-spectrum beta-lactamase-producing enterobacteriaceae in nonhospitalized patients. *Clin Infect Dis* 2009;49:682–90.
- Pasricha J, Koessler T, Harbarth S, *et al.* Carriage of extended-spectrum beta-lactamase-producing enterobacteriaceae among internal medicine patients in Switzerland. *Antimicrob Resist Infect Control* 2013;2:20.
- Han JH, Nachamkin I, Zaoutis TE, *et al.* Risk factors for gastrointestinal tract colonization with extended-spectrum β -lactamase (ESBL)-producing *Escherichia coli* and *Klebsiella* species in hospitalized patients. *Infect Control Hosp Epidemiol* 2012;33:1242–5.
- Hamprecht A, Rohde AM, Behnke M, *et al.* Colonization with third-generation cephalosporin-resistant *Enterobacteriaceae* on hospital admission: prevalence and risk factors. *J Antimicrob Chemother* 2016;71:2957–63.
- Huizinga P, van den Bergh MK, van Rijen M, *et al.* Proton pump inhibitor use is associated with extended-spectrum β -Lactamase-producing *enterobacteriaceae* rectal carriage at hospital admission: a cross-sectional study. *Clin Infect Dis* 2017;64:361–3.
- Centers for Disease Control and Prevention (2013). *Antibiotic resistance threats in the United States*: U.S. Department of Health and Human Services; Centers for Disease Control and Prevention, 2017.
- Stewardson AJ, Allignol A, Beyersmann J, *et al.* The health and economic burden of bloodstream infections caused by antimicrobial-susceptible and non-susceptible *Enterobacteriaceae* and *Staphylococcus aureus* in European hospitals, 2010 and 2011: a multicentre retrospective cohort study. *Euro Surveill* 2016;21.
- Cornejo-Juárez P, Suárez-Cuenca JA, Volkow-Fernández P, *et al.* Fecal ESBL *Escherichia coli* carriage as a risk factor for bacteremia in patients with hematological malignancies. *Support Care Cancer* 2016;24:253–9.
- Arnan M, Gudíol C, Calatayud L, *et al.* Risk factors for, and clinical relevance of, faecal extended-spectrum β -lactamase producing *Escherichia coli* (ESBL-EC) carriage in neutropenic patients with haematological malignancies. *Eur J Clin Microbiol Infect Dis* 2011;30:355–60.