

## Supplementary information

### Association between caesarean section delivery and obesity in childhood: a longitudinal cohort study in Ireland

Gwinyai Masukume,<sup>1,2</sup> Fergus P McCarthy,<sup>1,2,3</sup> Philip N Baker,<sup>4</sup> Louise C Kenny,<sup>5</sup> Susan MB Morton,<sup>6</sup> Deirdre M Murray,<sup>1,7</sup> Jonathan O'B Hourihane,<sup>1,7</sup> Ali S Khashan<sup>1,8</sup>

<sup>1</sup>The Irish Centre for Fetal and Neonatal Translational Research (INFANT), University College Cork, Cork, Ireland

<sup>2</sup>Department of Obstetrics and Gynaecology, University College Cork, Cork, Ireland

<sup>3</sup>Department of Women and Children's Health, School of Life Course Sciences, King's College London, London, United Kingdom

<sup>4</sup>College of Life Sciences, University of Leicester, Leicester, United Kingdom

<sup>5</sup>Department of Women's and Children's Health, Faculty of Health and Life Sciences, University of Liverpool, Liverpool, United Kingdom


<sup>6</sup>Centre for Longitudinal Research - He Ara ki Mua, University of Auckland, Auckland, New Zealand

<sup>7</sup>Department of Paediatrics and Child Health, University College Cork, Cork, Ireland

<sup>8</sup>School of Public Health, University College Cork, Cork, Ireland

#### Corresponding author:

Fergus P McCarthy: [Fergus.mccarthy@ucc.ie](mailto:Fergus.mccarthy@ucc.ie)



**Supplementary Figure 1** Mean body mass index (BMI) from birth to 5 years of age by delivery mode and sex. Please note that the time axis has been expanded below age 1 year to allow clearer visualisation.

**Supplementary Table 1** Missing data for body fat % at age two months.

Characteristic	Body fat % data available at age two months (n %) n=1033	Body fat % data missing at two age months (n %) n=272	P value <sup>a</sup>
Maternal age (years), median IQR	31 (28-33)	30 (28-33)	0.6021
Ethnicity <sup>b</sup>			0.558
Caucasian	1018 (98.5)	269 (98.9)	
Other	15 (1.5)	3 (1.1)	
Schooling (years primary and secondary), median IQR	13 (13-14)	13 (13-14)	0.5227
Marital status <sup>b</sup>			0.879
Single	100 (9.7)	23 (8.5)	
Married	725 (70.2)	195 (71.7)	
Stable relationship not married	207 (20.0)	54 (19.9)	
Sex of baby <sup>b</sup>			0.081
Male	540 (52.3)	126 (46.3)	
Female	493 (47.7)	146 (53.7)	
Pre-eclampsia <sup>b</sup>	40 (3.9)	9 (3.3)	0.664
Maternal BMI at 15 weeks (kg/m <sup>2</sup> ), median IQR	24.1 (22.1-26.9)	23.7 (22.0-26.7)	0.2455
Gestational age (weeks), median IQR	40 (39-41)	40 (39-41)	0.4624
Number of cigarettes per day at 15 weeks SCOPE visit, mean ( $\pm$ SD) <sup>c</sup>	0.5 ( $\pm$ 2.2)	0.4 ( $\pm$ 2.0)	0.2517
Birth weight (g), median IQR	3460 (3150-3770)	3475 (3160-3750)	0.9099

IQR – Interquartile range, BMI – Body mass index, SD – standard deviation, SCOPE – Screening for pregnancy endpoints.

<sup>a</sup> Mann-Whitney test

<sup>b</sup> Pearson's  $\chi^2$  test or Fisher's exact

<sup>c</sup> Two-sample t test

**Supplementary Table 2** Mode of delivery and body mass index at age two years. Non-marosomic.

BMI category (normal BMI – base outcome)	RRR (95% CI)	P value	AdjRRR (95% CI)**	P value
<b>Thin</b>				
Unassisted vaginal	Reference		Reference	
Operative vaginal	1.41 (0.84-2.36)	0.188	1.51 (0.89-2.58)	0.130
Prelabour LSCS	0.26 (0.24-1.62)	0.357	0.67 (0.27-1.68)	0.398
LSCS in labour	0.73 (0.32-1.64)	0.443	0.83 (0.37-1.90)	0.664
<b>Overweight or obese</b>				
Unassisted vaginal	Reference		Reference	
Operative vaginal	0.98 (0.58-1.64)	0.929	0.93 (0.54-1.59)	0.789
Prelabour LSCS	0.93 (0.44-1.95)	0.842	0.95 (0.44-2.05)	0.891
LSCS in labour	1.01 (0.51-1.98)	0.982	0.89 (0.44-1.82)	0.747

N for adjusted model = 921. Multinomial logistic regression. BMI – Body mass index, RRR (Relative Risk Ratio), CI (Confidence intervals), Adj (Adjusted).

\*\*Adjusted for maternal age, education, ethnicity, marital status, infant sex, maternal smoking during pregnancy, maternal BMI at the first antenatal visit, gestational age (at delivery), birth weight and pre-eclampsia

**Supplementary Table 3** Mode of delivery and body mass index at age five years. Non-macrosomic.

BMI category (normal BMI – base outcome)	RRR (95% CI)	P value	AdjRRR (95% CI)**	P value
<b>Thin</b>				
Unassisted vaginal	Reference		Reference	
Operative vaginal	1.59 (0.76-3.32)	0.221	1.85 (0.85-4.04)	0.120
Prelabour LSCS	0.73 (0.20-2.63)	0.629	0.46 (0.14-1.55)	0.209
LSCS in labour	1.09 (0.38-3.14)	0.880	1.14 (0.39-3.34)	0.815
<b>Overweight or obese</b>				
Unassisted vaginal	Reference		Reference	
Operative vaginal	1.43 (0.87-2.36)	0.161	1.77 (1.03-3.04)	0.038
Prelabour LSCS	0.89 (0.41-1.95)	0.768	0.86 (0.36-2.08)	0.750
LSCS in labour	1.59 (0.85-2.98)	0.150	2.37 (1.19-4.68)	0.014

N for adjusted model = 741. Multinomial logistic regression. BMI – Body mass index, RRR (Relative Risk Ratio), CI (Confidence intervals), Adj (Adjusted).

\*\*Adjusted for maternal age, education, ethnicity, marital status, infant sex, maternal smoking during pregnancy, maternal BMI at the first antenatal visit, gestational age (at delivery), birth weight and pre-eclampsia