

BMJ Open What factors are associated with resilient outcomes in children exposed to social adversity? A systematic review

Deirdre Gartland,^{1,2} Elisha Riggs,^{1,3} Sumaiya Muyeen,¹ Rebecca Giallo,^{1,3} Tracie O Afifi,⁴ Harriet MacMillan,⁵ Helen Herrman,⁶ Eleanor Bulford,¹ Stephanie J Brown^{1,2,3}

To cite: Gartland D, Riggs E, Muyeen S, *et al.* What factors are associated with resilient outcomes in children exposed to social adversity? A systematic review. *BMJ Open* 2019;**9**:e024870. doi:10.1136/bmjopen-2018-024870

► Prepublication history and additional material for this paper are available online. To view these files, please visit the journal online (<http://dx.doi.org/10.1136/bmjopen-2018-024870>).

Received 4 July 2018

Revised 24 January 2019

Accepted 30 January 2019



© Author(s) (or their employer(s)) 2019. Re-use permitted under CC BY-NC. No commercial re-use. See rights and permissions. Published by BMJ.

For numbered affiliations see end of article.

Correspondence to

Dr Deirdre Gartland;
deirdre.gartland@mcri.edu.au

ABSTRACT

Objectives Children exposed to social adversity—hardship as a result of social circumstances such as poverty or intergenerational trauma—are at increased risk of poor outcomes across the life course. Understanding what promotes resilient outcomes is essential for the development of evidence informed intervention strategies. We conducted a systematic review to identify how child resilience is measured and what factors are associated with resilient outcomes.

Design Systematic search conducted in CINAHL, MEDLINE and PsychInfo from January 2004 to October 2018 using the keywords ‘resilien* and child*’ in the title or abstract. Eligible studies: (1) described children aged 5–12 years; (2) identified exposure to social adversity; (3) identified resilience; and (4) investigated factors associated with resilience.

Outcome measures (1) approaches to identifying resilience and (2) factors associated with resilient outcomes.

Results From 1979 studies retrieved, 30 studies met the inclusion criteria. Most studies were moderate to high quality, with low cultural competency. Social adversity exposures included poverty, parent loss, maltreatment and war. Only two studies used a measure of child resilience; neither was psychometrically validated. Remaining studies classified children as resilient if they showed positive outcomes (eg, mental health or academic achievement) despite adversity. A range of child, family, school and community factors were associated with resilient outcomes, with individual factors most commonly investigated. The best available evidence was for cognitive skills, emotion regulation, relationships with caregivers and academic engagement.

Conclusions While there is huge variation in the type and severity of adversity that children experience, there is some evidence that specific individual, relational and school factors are associated with resilient outcomes across a range of contexts. Such factors provide an important starting point for effective public health interventions to promote resilience and to prevent or ameliorate the immediate and long-term impacts of social adversity on children.

BACKGROUND

Why is it important to understand resilience in children?

Social adversity can be defined as exposure to hardship as a result of social circumstances, for example, poverty, racial discrimination, maltreatment, intergenerational trauma or

Strengths and limitations of this study

- This is the first systematic review summarising research about the individual, family, school, social and community factors associated with resilient outcomes in children across a broad range of social adversities including child maltreatment, war and poverty.
- The synthesis was limited to a descriptive review of the studies due to the significant heterogeneity in the measurement approaches, categorisation of resilience and the resilience factors investigated.
- The studies reviewed were limited to those published in English.

community violence. Significant numbers of children are exposed to social adversity in childhood. One in eight adults report childhood sexual abuse, one in four report physical abuse^{1,2} and more than one in four children are exposed to intimate partner violence.^{3,4} Globally, 28 million children are uprooted by war and conflict with attendant trauma, hardship and disadvantage.⁵ One in 10 Australians live below the internationally accepted poverty line, with almost one quarter of these being dependent children,⁶ and Indigenous families experience a far greater cumulative load of early life stress and social adversity, including intergenerational trauma.^{7,8}

Experiences of social adversity have multiple direct and indirect effects on the social, mental and physical health of individuals and families.^{9,10} Children exposed to social adversity are more likely to experience emotional and behavioural problems, mental health disorders, speech and language problems, learning difficulties and physical disorders.^{4,11–15} Furthermore, childhood experiences of adversity are associated with lifelong negative impacts on health and well-being.^{16–20} There is also clear evidence of a clustering of social adversity, including

trauma, in families and communities.^{16 10 21–23} In summary, child exposure to social adversity is common, and there is clear evidence of negative impacts for many children. At the same time, there is also evidence that some children show positive outcomes.

What is resilience?

Resilience has been defined as positive developmental outcomes in the face of adversity or stress,²⁴ being relatively resistant to psychosocial risk experiences²⁵ and the development of competence despite chronic stress.²⁶ While differing in terminology, these descriptions encompass the two common factors necessary for defining resilience: first, the experience of adversity or stress, and second, the achievement of positive outcomes during or following the exposure to adversity. Current research predominantly views resilience as the *process* by which individuals draw on personal characteristics and resources in their environment to withstand and negotiate adversity—a dynamic process across contexts and over the life course.^{27 28}

Resilience can be considered within an ecological framework, with risk and protective factors present within an individual; in their proximal environments of family, school or work; and at the more distal community and societal levels. A range of factors has been identified as supporting resilient outcomes in adolescence and adulthood. Individual factors such as coping style, cognition, optimism and self-esteem,^{29–31} positive family relationships or social connectedness,^{30–32} and community factors such as school engagement,³³ church or support groups.^{32 34} Most of the resilience research to date has focused on adult or adolescent resilience to trauma or other types of adversity (although the trauma may occur in childhood). There is limited research available on factors that support resilience in middle childhood (5–12 years).

Why focus on childhood?

While there have been recent systematic reviews of adult resilience factors,^{35 36} there is only one systematic review on child resilience and this focuses on interventions to build resilience.³⁷ As the key developmental tasks in adolescence or adulthood differ greatly from those in childhood, there may be key differences in the factors that support resilience in children. For example, while adolescents are developing independence from family and focusing on relationships with peers, a key developmental task for children is building a close supportive bond with a caregiver. Resilience factors identified as important for adults or adolescents, such as optimism, positive family relationships and school engagement^{30 31 33} have obvious relevance for children, while others such as employment, church and community support may be less important. There may also be factors that are key in childhood that have not been identified in the adult literature. Exploring the factors that are associated with resilient outcomes in children is essential to improve outcomes for the millions of children experiencing poverty, violence and other social adversities.

While the data are complex, economic evaluations suggest that investment in evidence-based interventions in childhood can have significantly greater and longer lasting impacts per dollar spent, compared with intervening in adulthood.^{38 39} Additionally, early intervention has the potential to interrupt the intergenerational transmission of trauma and disadvantage.⁴⁰ While there is huge variation in the challenges children may experience, ranging from societal factors, such as poverty, to individual exposures, such as sexual abuse, there are likely to be resilience factors that are important across different contexts and adversities. Better evidence as to what promotes resilient outcomes in children exposed to trauma or adversity is essential for effective interventions to prevent or ameliorate the immediate and lifelong impacts.^{41 42}

Aims

The purpose of this systematic review was to identify studies that have examined resilience in children exposed to social adversity. Specifically, the objectives were to identify: (1) how resilience was assessed in research studies; and (2) the individual, family, social, school, community/neighbourhood and societal factors associated with positive outcomes or resilience in children who have experienced social adversity.

METHODS

Titles and abstracts of papers identified by the search terms described below were screened by three reviewers (DG, EB and SM) against the inclusion and exclusion criteria. Papers remaining after this initial screening were imported into EndNote, and full-text articles were obtained. A more in-depth examination of the Methods section of each paper was conducted to ensure studies matched the inclusion criteria. Disagreements as to inclusion/exclusion were discussed with a fourth reviewer (ER) where necessary, with final decisions agreed by consensus.

Patient and public involvement

Patients were not involved in this study.

Inclusion criteria

Table 1 shows the study inclusion and exclusion criteria. Eligible studies were limited to research studies describing original research and written in English. Reviews, case reports, commentaries, government reports and non-peer-reviewed papers were excluded. The population of primary interest was children aged 5–12 years who had been exposed to ‘social adversity’. Social adversity was defined as exposure to trauma or hardship as a result of social circumstances, for example, poverty, racial discrimination, violence, war or maltreatment. Studies in which participants were exposed to hardship as a result of physical rather than social factors, such as natural disasters, cancer, genetic conditions or physical disability were not included. Studies that included some children

Table 1 Review inclusion and exclusion criteria

	Inclusion criteria	Exclusion criteria
Study design	Cohort studies, cross-sectional studies, randomised trials and qualitative studies reporting original research.	Reviews, case reports and commentaries.
Type of paper	Peer-reviewed published papers.	Non-peer-reviewed papers, grey literature.
Participants	Children aged 5–12 years exposed to ‘social adversity’ defined as exposure to trauma or hardship as a result of social circumstances (eg, poverty, war or maltreatment).	Exposure to hardship as a result of physical rather than social factors, for example, natural disasters, cancer, genetic disease or physical disability.
Informants	Children, parents/carers, school personnel and health professionals.	
Outcomes of interest	(1) How child resilience is identified or assessed (eg, psychometric measure or defined criterion). (2) Factors associated with resilient outcomes.	

outside the age range were included if enough detail was provided to ascertain that the majority of participants were aged 5–12 years. While much research has been conducted over the previous 50 years on factors associated with positive child outcomes, for example, success in school or positive coping skills, given that our focus was on child resilience, studies were required to identify or name ‘resilience’ as a focus of the investigation.

The objectives of this review were twofold: (1) to identify how child resilience is identified/measured and (2) the factors associated with resilient outcomes in children. Factors were considered within the socioecological domains of individual, family, social, school and neighbourhood or community.

Search strategy

The search strategy was developed in consultation with a librarian. We searched published literature from January 2004 to October 2018. We limited the search to the previous 10 years (at the time of developing the protocol) to focus on the more recent approach to resilience research, where resilience is seen as a dynamic process rather than a static individual characteristic. Selection of electronic databases was based on the topic of our systematic review and included two general databases (MEDLINE and CINAHL) and a databases more specific to the topic area (PsychInfo). Databases were searched for articles in English with *resilien** and *child** in the title or abstract. Papers with *genetic* or only *adolesce** (without *child*) in the title or abstract were excluded as per criteria described above. This simple approach was adopted in preference to a more complex search strategy as the significant heterogeneity in adversity exposures, child outcomes and resilience factors described in the literature meant that more complex searches missed papers we were already aware of.

Data extraction

Data were extracted using a structured data abstraction form, with the first 10% of papers independently abstracted by two reviewers, and the results were

compared to ensure adherence to the protocol, check the quality of the data and check consistency in data abstraction. Data extracted included study design, country, sample size, participant characteristics (eg, age and gender), informant (eg, parent, social worker and/or child), social adversity exposure, resilience definition/criteria, resilience measure (if included), resilience factors measured and factors associated with resilience. The quality of the included studies was appraised by one author using the Health Evidence Bulletin Wales critical appraisal tool adapted from the Critical Appraisal Skills Programme.⁴³ This tool assesses key domains of study quality, including clarity of aims, appropriateness and rigour of design and analysis, risk of bias and relevance of results. To appraise the studies in terms of culturally safe and participatory methods, a cultural competency appraisal tool was also used.⁴⁴

RESULTS

The initial search identified 1979 papers, which was reduced to 348 after initial screening of titles and abstracts (see [figure 1](#)). A further 318 studies were excluded after obtaining the full papers and examining study methods in greater detail. Papers were excluded due to: not investigating child resilience and/or factors associated with resilience (n=786); participants outside the age range or age not specified (n=283); and not an original research report. Thirty papers describing 30 studies met the inclusion criteria.

[Table 1](#) provides details of the 30 included studies conducted from 2004 to 2017. The majority of studies was conducted in North America (n=20, 67%). The remaining studies were conducted in Africa (2), Asia (3), Australia (1) and Europe (4). Participants ranged in age from 2 years to 16 years (n=26 studies), with the mean age ranging from 4.5 years to 11.9 years (n=19 studies). Twelve studies (40%) were prospective longitudinal




Figure 1 Flow chart of paper selection.

cohort studies and 18 were cross-sectional studies (60%). Two studies (7%) included qualitative methods.

The most common social adversity described was poverty (n=14, 47%) (see [Table 2](#)). Belonging to a minority population group, such as an ethnic minority, is associated with a higher risk of social disadvantage and adversity across the life course.^{9 40 45 46} Seven studies reported that participants belonged to an ethnic minority (23%), with six of these seven studies also identifying poverty as an adversity exposure. The seventh study was an examination of cross-ethnic friendships in South Asian British children in the context of perceived ethnic discrimination.⁴⁷ None of the included studies separately identified Indigenous or refugee children, although Cortina *et al*⁴⁸ examined refugee status as a risk factor for children living in rural Southern Africa.

Seven studies described children who had been abused or neglected (23%) and four (13%) described children exposed to violence (including intimate partner violence, community or school violence). Eight studies described children with interruptions in the caregiving relationship: five (17%) were about parental loss or illness (HIV, mental illness, incarceration or death) and three reported on children in out-of-home care (10%). One study described children living in a war-affected zone (3%).

Critical appraisal of the included studies using the Health Evidence Bulletin Wales critical appraisal tool determined that the majority of the studies were of high quality (see online supplementary table 1a). In terms of culturally safe and participatory study methods,⁴⁴ most of the included studies were of low quality, with only one study—investigating resilience in South African children⁴⁸—classified as high quality (see online supplementary table 1b).

Measuring resilience in middle childhood

Two studies (7%) used a resilience measure. Bagci *et al*⁴⁷ used a brief resilience scale developed by Bartko

and Eccles⁴⁹ to assess resilience in 247 South Asian British children. The scale was originally developed for American adolescents and asks respondents 'How often are you very good at' on four items related to problem solving, planning, bouncing back from bad experiences and learning from mistakes. Bagci *et al*⁴⁷ report a relatively low scale reliability of 0.6. No other psychometric properties were reported.

Somchit and Sriyaporn⁵⁰ report using a resilience framework developed by Grotberg to develop a resilience measure to examine perceptions of adversity and academic achievement in 267 Thai children. The paper provides no information about the properties of the newly developed measure, and it is unclear how Grotberg's resilience framework was operationalised. There is no information included in the paper about the items, scales or psychometric properties of the measure.

The remaining 28 studies classified children as resilient by specifying positive outcomes on one or more measures of mental health (eg, internalising problems or life satisfaction) or functioning (eg, cognitive outcomes, academic achievement or social behaviours) in the context of exposure to adversity or trauma. Of the 12 longitudinal cohort studies, all studies used positive functioning in a single domain to define resilience. Six studies used a measure of positive mental health to classify children as resilient (see [table 2](#)), four identified an aspect of child behaviour (eg, low levels of antisocial behaviour) and two used an academic measure (eg, positive classroom behaviour and academic performance). Lagasse *et al*⁵¹ used positive outcomes across both social and mental health domains but investigated each separately (ie, resilience was not having a mental illness diagnosis or lower levels of delinquent behaviour rather than a composite of a positive outcome in both areas).

There was considerable variation in the approaches used to define resilience. For example, Halevi *et al*⁵² classified Israeli war-exposed children as resilient if they exhibited a lack of psychopathology on the Development and Well-Being Assessment. Brown *et al*⁵³ used latent growth curve analysis of poor black American students' internalising symptoms over time and classified students who exhibited stable low level internalising symptoms as resilient. While both studies used mental health to identify children as resilient, they employed very different measures and approaches. Across all 30 studies, the measures used and the methodological approach to identifying children as resilient or non-resilient/vulnerable varied greatly.

In the 18 cross-sectional studies, three studies identified resilience using a composite of positive child outcomes across more than one domain, for example, mental health and social competence.^{54–56} Lin *et al*⁵⁷ used a composite of three outcome measures but all were within the domain of mental health (below clinical cut-off for depression, anxiety and internalising/externalising problems). The remaining studies used a positive child outcome in a single domain of mental, behavioural or academic outcomes to identify resilience in children exposed to

Table 2 Summary of reviewed papers (n=30)

First author (year)	Country	Sample		Age Range	Mean (SD)	% Female		Informant	Exposure	Resilience measure or criteria	Resilience factors measured	Factors significantly associated with resilience
		size	Age			Mean (SD)	Female					
Longitudinal cohort studies												
Brown (2013) ⁵³	USA	1603	5–11	NR	NR	0	Child, Parent, Teacher, School representative.	Poverty, Ethnic minority.	↓ Internalising problems.	Cognitive skills, Family environment, School environment, Neighbourhood quality.	Cognitive skills.	
Coohney (2011) ⁵⁷	USA	702	6–10	7.8 (2.9)	53.8		Child, Caregiver, Child protective services investigators.	Maltreatment.	↑ Math score, ↑ Reading.	Intelligence, Daily living skills, School – engagement, Peer relationships, Emotional support – caregiver.	Intelligence, Daily living skills.	
Dubowitz(2016) ⁸³	USA	943	4–6	NR	NR	51.0	Child, Caregiver.	Maltreatment.	↓ Externalising/internalising problems, ↑ Developmental functioning.	Caregiver mental health, Social ecology of family.	Caregiver mental health, Social ecology (employment and less children).	
Flouri (2014) ⁵⁸	UK	16916	3–7	NR	NR	49.1	Child, Parent.	Poverty.	↓ Emotional/behavioural problems.	Self-regulation, Verbal cognitive ability.	Self-regulation, Verbal cognitive ability.	
Halevi (2016) ⁵²	Israel	232	1.5–11	NR	NR	NR	Child, Mother.	War/conflict.	↓ Psychopathology.	Maternal uncontained style, Child social engagement, Maternal distress, Gender.	Maternal uncontained style, Child social engagement, Maternal distress, Gender (girl).	
Honer (2012) ⁸⁵	USA	1733	Grades 4, 5, 6 and 7	NR	NR	53.7	Child, Parent.	Poverty, Ethnic minority.	Health promoting behaviour.	Social connectedness, Humour, Healthy behaviour, Being loved, Self-concept.	Self-concept, Being loved.	
Jaffee (2007) ⁵⁹	UK	2181	5–7	NR	NR	51.0	Mother, Teacher, Child, Research worker.	Maltreatment.	↓ Antisocial behaviour.	Cognitive skills, Temperament, Maternal warmth, Sibling warmth, Neighbourhood quality (informal social control and social cohesion), Norms for antisocial behaviour.	Cognitive skills, Neighbourhood quality, Temperament.	
Lagasse (20016) ⁵¹	USA	517	0–11	NR	NR	NR	Child, Caregiver, Teacher.	Poverty, Parent substance abuse, Community violence.	↓ Crimes against people, ↓ Delinquency, ↓ Drug use, ↓ Depression.	Relatedness to others, Effortful control.	Relatedness to others, Effortful control.	
Manly (2013) ⁸⁴	USA	170	4–6	NR	NR	52.4	Child, Parent, Teacher.	Maltreatment.	↑ Classroom behaviour, ↑ Academic performance.	Cognitive function, Ego resiliency, Flexibility, Positive affect, Self-esteem, Creativity.	Cognitive function, Ego resiliency.	
Ng (2014)	China	150	7–12	8.8 (1.0)	50.7		Child.	Poverty.	↑ Life satisfaction.	Hope, Community support, Community integration, Community participation, Community organisation.	Hope, Community support.	

Continued

Table 2 Continued

First author (year)	Country	Sample size	Age Range	%		Resilience measure or criteria	Exposure	Informant	Resilience factors measured	Factors significantly associated with resilience
				Mean (SD)	Female					
Ramos-Olazagasti (2010) ⁷²	USA/Puerto Rico	1271	5–13	11.6 (NR)	50.0	↓ Internalising symptoms.	Poverty. Ethnic minority. Community violence.	Child. Parent.	Parental monitoring. Religiosity.	Parental monitoring. ↓Religiosity.
Yoon (2018) ⁶⁴	USA	449	4–5	4.5	48.3	Three classes identified by modelling externalising problems time 1–time 4. Resilient=low/stable group.	Maltreatment.	Caregiver. Child protection services.	Child prosocial skills. Caregiver well-being	Child prosocial skills. Caregiver well-being.
Cross-sectional studies										
Anthony (2008) ⁶²	US	157	NR	11.9 (1.3)	51.6	↑ Academic grades. ↓ Drug use. ↓ Delinquency. ↓ Antisocial behaviour.	Poverty.	Child. Teacher.	Self-esteem. Coping. Academic self-efficacy. School commitment. Social support – family. Social support – peers. Social support – neighbourhood. Social support – adults not family. Neighbourhood cohesion. Positive parenting.	Self-esteem. Coping. School commitment. Social support – family. Social support – peers. Social support – neighbourhood. Neighbourhood cohesion. Positive parenting.
Bagci (2014) ⁴⁷	UK	247	NR	11 (0.3)	56.3	Resilience Scale (Bartko and Eccles).	Ethnic minority (discrimination).	Child.	Friendship quality. Friendship quantity.	Friendship quality. Friendship quantity.
Bell (2013) ⁷⁰	Canada	531	5–9	7.4 (1.4)	47.3	↓ Emotional/behavioural problems.	Out of home care.	Caregiver. Welfare workers. Child aid societies.	Social support – peers. Parenting skills. Academic performance. Positive values. Social skills. Positive identity.	Positive identity. Parenting skills. Positive values.
Borman (2014) ⁸⁰	USA	925	6–12	NR	NR	↑ Mathematics achievement.	Poverty. Ethnic minority.	Child. Parent. Teacher. School principal. School district personnel.	Student – engagement. Student – positive attitude. Internal locus of control. Self-efficacy – math. Supportive school environment: Safe. Orderly. Positive student–teacher relations. School resources (eg, teacher years and class size). Effective school (eg, monitoring of students and principal leadership).	Student – engagement. Student – positive attitude. Internal locus of control. Self-efficacy – math. Supportive school environment: Safe. Orderly. Positive student–teacher relations.

Continued

Table 2 Continued

First author (year)	Country	Sample		Age Range	Mean (SD)	% Female	Informant	Exposure	Resilience measure or criteria	Resilience factors measured	Factors significantly associated with resilience
		size	Age								
Cicchetti (2007) ⁵⁴	USA	677		6–12	9.5 (2.3)	45.2	Child. Peer. Camp counsellor.	Maltreatment.	Composite across seven indicators: ↓ Depressive symptoms. ↑ Social competence (prosocial, disruptive/aggressive and withdrawn). ↓ Internalising problems. ↓ Externalising problems. ↓ School risk. Resilient=high functioning in 5–7 areas.	Social skills. Ego-resilience. Ego-control.	Ego-resilience. Ego-control.
Cortina (2013) ⁴⁸	South Africa	1025		10–12	NR	49.2	Child. Teacher. Pre-existing records.	Poverty.	↓ Emotional/behavioural problems. ↓ Depression/anxiety. ↑ Peer relationships.	Maternal characteristics (eg, age, SES, education and refugee status). Household size. Household stability (mother present and months in household). Breast fed.	Maternal education. Non-refugee status. SES. Mother partnered. Older child age.
Dallaire (2013) ⁶⁹	USA	210		7–12	9.1 (1.1)	56.0	Child. Parent. Peer. Researcher.	Parent incarceration.	↓ Aggressive behaviour.	Empathy.	Empathy.
Fantuzzo (2012) ³³	USA	3535		8	NR	0.0	Teacher. Pre-existing records.	Cumulative risk experiences including poverty, ethnic minority, maltreatment and homelessness.	↑ Academic achievement.	Academic engagement.	Academic engagement.
Flores ⁵⁵	USA	133		NR	8.7 (1.8)	30.8	Child. Camp counsellor. Peer.	Maltreatment. Poverty. Ethnic minority.	Composite of high functioning across nine indicators of: ↑ Behavioural functioning. ↑ Social competence. ↓ Externalising/internalising problems. Resilient=high functioning in 6–9 areas.	Intelligence. Ego-resilience. Ego-control. Relationship with adult outside family. Social skills.	Ego-resilience. Ego-control. Relationship with adult outside family.
Graham (2009) ⁵⁶	USA	219		6–12	8.5 (2.2)	50.2	Child. Mother.	Intimate partner. Violence.	Composite of: ↓ Internalising/externalising problems. ↓ Depressive symptoms. ↑ Global self-worth/social competence. ↓ Interpersonal worries/anxiety. Four clusters: resilient=high competence and low adjustment problems.	Positive maternal mental health. Parenting warmth. Parenting effectiveness. Family functioning.	Maternal mental health. Parenting effectiveness.

Continued

Table 2 Continued

First author (year)	Country	Sample		Age Range	Mean (SD)	% Female	Informant	Exposure	Resilience measure or criteria	Resilience factors measured	Factors significantly associated with resilience
		size	Range								
Holiday (2014)	USA	230	3–7	5.8 (0.4)	47.4	Child, Caregiver, Teacher	Poverty.	↑ School performance.	Parent engagement, Friendships, Culturally compatible school, Childcare, Physical health, Parent education level, Language, Literacy.	Childcare, Parent education level.	
Jones (2007) ⁷¹	USA	71	9–11	NR	56.3	Child.	Poverty, Community violence.	↓ Post Traumatic Stress Disorder symptoms.	Formal kinship support, Spirituality, Afrocentric support.	Formal kinship support, Spirituality, Afrocentric support.	
Lin (2004) ⁵⁷	USA	179	8–16	11.6 (2.5)	45.8	Child, Caregiver, Teacher.	Parent death/absence.	Composite: ↓ Externalising/internalising problems, ↓ Depression, ↓ Anxiety, Resilient=less than clinical cut point on all three measures.	Caregiver mental health, Caregiver warmth, Caregiver discipline, Inhibition of emotional expression, Control beliefs, Self-esteem, Coping efficacy, Perception of negative events.		
Myers (2013) ⁶¹	USA	61	7–13	9.7 (1.6)	55.7	Child, Camp mentors.	Parent incarceration.	↓ Bullying.	Emotion regulation.	Emotion regulation.	
Rees (2013)	UK	193	7–15	NR	47.7	Child, Caregiver, Teacher.	Out-of-home care.	↓ Emotional/behavioural problems.	Mental health, Emotional literacy, Cognitive ability, Literacy attainment, Parental contact.	Parental contact.	
Somchit (2004) ⁵⁰	Thailand	267	9–16	11.9 (1.1)	54.3	Child.	Poverty.	Investigator developed measure from International Resilience Project.	Academic achievement, Gender.	Gender (girl).	
Qualitative studies											
Downie (2010) ⁶⁵	Australia	20	8–15	10.8 (2.1)	60.0	Child	Out-of-home care.	↑ Self-evaluation (self-concept and well-being).	Coping strategies, Emotional health (safety, love, family contact), Material needs met, Discipline, Contact with family members.	Emotional health (safety, care/love and family contact), Material needs met, Contact with family members.	
Ebersöhn (2015) ⁶⁸	South Africa	19	5–7	NR	57.9	Child, Mother.	HIV/AIDS	↓ Externalising/internalising problems.	Problem-focused coping, Emotional intelligence sense of belonging, Seek help/support/resources, Spirituality/religious identification, Positive future expectancy, Resolve/agency.	Problem-focused coping, Emotional intelligence sense of belonging, Seek help/support/resources, Spirituality/religious identification, Positive future expectancy, Resolve/agency.	

NR, not reported; SES, Socio Economic Status.

adversity (see table 2). Overall, no two papers employed the same method to identify children as resilient.

What factors are associated with resilient outcomes?

In addition to examining the way in which studies identified resilience in children, we also abstracted information on factors associated with resilient outcomes. Table 3 summarises the included studies according to a single primary social adversity to avoid artificially inflating results and the factors that were identified as significantly associated with child resilience. The factors investigated in each study have been grouped into the domains of individual, family, social, school and neighbourhood/community. Each domain will be described in turn.

Individual factors

Individual characteristics were identified as significantly associated with resilient outcomes in all social adversity contexts. Most studies included individual characteristics (n=24, 80%). A wide range of factors was investigated including gender, temperament, emotion regulation, cognitive skills, social skills, self-efficacy and self-esteem (see table 3). Five longitudinal studies identified a significant association between an aspect of cognition and resilient outcomes in the different adversity exposures. For example, in a large study of child emotional and behavioural resilience in the context of persistent poverty, verbal cognitive ability was found to be significantly associated with low levels of child internalising problems.⁵⁸ Brown *et al*⁵³ found moderate support for an association between cognitive abilities and stable low internalising problems over time in a large longitudinal cohort of 1603 boys living in poverty (and belonging to an ethnic minority). Jaffee *et al*⁵⁹ reported that maltreated boys with above average intelligence had a significantly higher likelihood of being in the resilient group; however this association was not evident for girls (resilience was defined as antisocial behaviour scores at or below the median of the non-maltreated group).

Eight studies identified associations between self-regulation and resilience: three longitudinal cohort studies, four cross-sectional and one qualitative study. Self-regulation refers to purposeful modulation of thoughts, emotions or behaviour or 'effortful regulation of the self by the self'.⁵⁸ Although Flouri *et al*⁵⁸ found that self-regulation benefited both poor and non-poor children, there was an increasing gap in outcomes over time between the poor children with high and low regulation, particularly in terms of internalising problems. The trajectories of internalising and externalising problems of poor and non-poor children with high self-regulation were similar, particularly as they got older.⁵⁸ Borman and Overman⁶⁰ in a cross-sectional study of 925 low SES third grade students purposively selected from a longitudinal cohort found a higher internal locus of control to be associated with resilient outcomes (as defined by performing better than expected in maths), irrespective of minority status. Emotion regulation also predicted group membership of

low rather than high bullying in 61 children attending a summer camp for children of incarcerated mothers.⁶¹

Two papers reported aspects of self-esteem to be significantly associated with resilient outcomes. Anthony⁶² identified four distinct clusters among 157 ethnically diverse children living in public housing in the USA. High levels of self-esteem and coping skills (among other factors) were found among children in the first cluster, who were found to be resilient in terms of low drug use and delinquency and better academic grades. A large longitudinal study of ethnically diverse children living in rural Texas similarly found that children with a self-perception of competence were less likely to engage in health risk behaviours (ie, smoking cigarettes, using marijuana, drinking alcohol or carrying a weapon).⁶³

As shown in table 2, a range of other child characteristics was associated with resilient outcomes in one or two studies including prosocial skills,⁶⁴ coping skills,^{57 65} hope,⁶⁶ daily living skills,⁶⁷ help seeking⁶⁸ and empathy.⁶⁹

Family and social support factors

Sixteen studies examined aspects of caregiver relationships, family environment and/or parenting skills across all the social adversity contexts, with significant factors identified in all contexts. For example, support from family was associated with significantly lower engagement in health risk behaviours in a sample of almost 2000 ethnically diverse children living in poverty in rural Texas.⁶³ In a mixed methods study, feeling loved, cared for and supported by their caregivers was significantly associated with emotional health for 20 children who were in the full time care of their grandparents.⁶⁵ For 219 children who had been exposed to intimate partner violence in the previous year, parenting warmth was one of the factors that significantly differentiated resilient children from those with severe problems.⁵⁶

Fewer studies (n=11) investigated social factors associated with resilient outcomes across the different adversity contexts. Broader social support from friends or adults outside the family was identified as significant in the face of poverty, child abuse, violence and parental absence/loss. For example, in a longitudinal cohort of 517 children aged 11 years who had been exposed to violence (child abuse, family violence, violent friends and/or community violence), the authors reported that children who had positive 'relatedness to others' scores (a composite of positive relationships and prosocial behaviour with teachers, parents, friends and peers) were more resilient in terms of experiencing less school-specific delinquency (truancy or suspension), depression, oppositional defiance disorder and conduct disorder.⁵¹ Furthermore, in a cross-sectional study, the ability to form a positive relationship with an adult figure outside of the immediate family predicted resilience in maltreated Latino children.⁵⁵ Finally, investigation of the benefits of cross-ethnic friendships in 247 South Asian British children identified that friendship quality, but not the number of friends, was associated with psychological well-being and resilient outcomes.⁴⁷

Table 3 Significant resilience factors (grouped into themes) in the individual, family, social, school and community domains by primary adversity exposure*

Adversity	Individual domain			Family domain			Social domain			School domain			Community domain		
	First author, year	n†	Factor (no. of papers)‡	n†	Factor (no. of papers)‡	n†	Factor (no. of papers)‡	n†	Factor (no. of papers)‡	n†	Factor (no. of papers)‡	n†	Factor (no. of papers)‡	n†	Factor (no. of papers)‡
Poverty ethnic discrimination (6/11 involved an ethnic minority)	Anthony ⁶² , 2008. Bagci ⁴⁷ , 2014. Borman ⁶⁰ , 2004. Brown ⁵³ , 2013. Cortina ⁴⁸ , 2013. Fantuzzo ⁶³ , 2012. Flour ⁵⁸ , 2014. Holliday, 2014. Horne ⁶³ , 2012. Ng, 2014. Somchit ⁵⁰ , 2004.	8	Emotion regulation (1). Cognitive abilities (2). Coping (1). Hope (1). Self-esteem (1). Self-efficacy (1). Self-identity (1). Gender (1).	6	Family relationships (1). Parenting skills (1). Family SES (2).	4	Family relationships (1). Friends (2).	5	School environment (1). Teachers (1). Student engagement (3).	5	Social support (1). Friends (2).	5	School environment (1). Teachers (1). Student engagement (3).	5	Neighbourhood (1). Community (1).
Parent absence (illness, death, incarceration and out-of-home care)	Bell ⁷⁰ , 2013. Dallaire ⁶⁹ , 2013. Downie ⁶⁵ , 2010. Ebersohn ⁶⁸ , 2010. Lin ⁵⁷ , 2004. Myers ⁶¹ , 2013. Rees, 2013.	7	Cognition (1). Coping (1). Emotion regulation (2). Empathy (1). Self-efficacy (1). Self-identity (1). Help seeking (1).	4	Family relationships (1). Caregiver relationship (1). Parenting skills (1). Family SES (1).	2	Family relationships (1). Caregiver relationship (1).	2	Social support (1).	1	Social support (1).	1	Social support (1).	1	Spirituality (1).
Maltreatment (one or more of physical abuse, emotional abuse, sexual abuse and neglect)	Cicchetti ⁵⁴ , 2007. Coohey ⁶⁷ , 2011. Dubowitz ⁶³ , 2016. Flores ⁶⁵ , 2009. Jaffee ⁵⁹ , 2007. Manly ⁶⁴ , 2013. Yoon ⁶⁴ , 2017.	7	Cognition (3). Emotion regulation (3). Temperament (1). Positive affect (1). Living skills (1). Prosocial skills (1).	4	Caregiver relationship (1). Family environment (1). Parent mental health (1). Caregiver well-being (1).	3	Caregiver relationship (1). Family environment (1). Parent mental health (1). Caregiver well-being (1).	3	Social support (1).	0	Social support (1).	0	Social support (1).	1	Neighbourhood (1).
Violence (intimate partner and community)	Graham-Bermann ⁵⁶ , 2009. Jones ⁷¹ , 2007. Lagasse ⁵¹ , 2006. Ramos-Olazagasti ⁷² , 2013.	2	Emotion regulation (1). Positive affect (1).	3	Family relationships (1). Parenting skills (2). Parent mental health (1).	2	Family relationships (1). Parenting skills (2). Parent mental health (1).	2	Social support (2). Social skills (1).	0	Social support (2). Social skills (1).	0	Social support (2). Social skills (1).	2	Spirituality (2). Culture (1).
War	Helevi, 2016	1	Gender (1).	1	Parent mental health (1).	1	Parent mental health (1).	1	Social skills (1).	0	Social skills (1).	0	Social skills (1).	0	

*Papers included multiple exposures but are presented in a single primary adversity category to avoid artificially inflating counts.

†Number of papers reporting on individual, family, social, school or community resilience factors.

‡Number of studies identifying significant associations with resilient outcomes.

School factors

School factors were investigated in two contexts only: five studies in the context of poverty and one study in the context of parental loss/illness. School factors included both student characteristics, such as academic engagement, and school environment factors. Student academic engagement was reported to mediate the association between risk experiences (a count that included child maltreatment, low maternal education at birth, homeless shelter stay, poor prenatal care, preterm birth/low birth weight and lead exposure) and both reading and mathematics outcomes in a cross-sectional study involving almost 4000 third grade African-American boys.³³ For ethnically diverse children living in public housing developments in the USA, school commitment was significantly associated with resilient outcomes in terms of lower delinquency and drug use and better school grades.⁶² A single study investigated parental loss and reported that academic performance was not significantly associated with resilient outcomes (lower levels of emotional and/or behavioural problems) in a sample of 531 children living in out-of-home care.⁷⁰

In relation to aspects of the school environment, a large cross-sectional study of 925 students living in poverty examined the school characteristics for resilient (above expected mathematics achievement) versus non-resilient students (below expected mathematics achievement).⁶⁰ The authors reported few differences for *school resources* (eg, class sizes) or *effective school environment* (eg, strong principal leadership) but did find significant differences with respect to *supportive school community*. Resilient students were significantly more likely to come from schools with positive student–teacher relationships, a safe and orderly environment and that were supportive of family involvement.

Community factors

Eight studies explored community factors associated with resilient outcomes, seven of which identified significant associations between diverse aspects of community or culture. For example, in a large longitudinal cohort study of children living in relatively high crime neighbourhoods, those children who lived in communities with high social cohesion and informal social control were more likely to be resilient to maltreatment.⁵⁹ Perceived community support was associated with life satisfaction in a sample of 75 children living in poverty in Hong Kong.⁶⁶ Finally, spirituality was associated with resilience in a qualitative study involving institutionalised African children and African-American children living in a high crime, high poverty area in Texas.^{68 71} However, in one longitudinal prospective cohort study, family religiosity was associated with higher levels of internalising symptoms for Puerto Rican children.⁷²

DISCUSSION

While there is a growing body of resilience research, to date, there has been a limited focus on child resilience in the context of social adversity and disadvantage, especially among children aged 5–12 years. Of the 30 studies

included in this systematic review, only two employed a structured measure of resilience and neither of these measures was validated. Most studies relied on categorisation to identify resilience (exposure to adversity+positive achievement in a particular outcome=resilience). Across the 28 studies using this approach, there was significant variation in outcome measurement and how children were classified as resilient/non-resilient. These findings are consistent with a recent systematic review of the operationalisation of resilience in longitudinal studies by Cosco *et al.*³⁵ The authors comment that the considerable ‘heterogeneity in adversity/adaptation dyads and operationalisation methods’ (p. 5) meant they were unable to meaningfully review the actual protective factors investigated in the studies they reviewed. This existing variation in the identification of resilience makes comparisons across studies, populations and contexts problematic. The adoption of a more consistent approach to defining and measuring child resilience would enhance our understanding of resilience across different contexts and improve our ability to examine the effectiveness of interventions.^{26 35 41 73 74}

Most of the included studies identified child resilience by measuring positive outcomes in a single domain, such as mental health. Resilience is a multidimensional construct that may vary across domains.^{26 73–75} Positive outcomes in a single domain may mask negative outcomes in other areas, and conversely, while adversity and trauma will impact the mental health of all exposed children to some degree, they may exhibit resilience in other domains such as academic or behavioural. Therefore, the identification of factors associated with resilience based on a single domain may not be generalisable to other domains or settings.

These limitations aside, there appears to be a range of factors that may contribute to resilience in children across different social ecological domains. At an individual level, factors such as emotion regulation, cognitive skills, empathy or a positive outlook have been associated with resilient outcomes. In the family and social domains, good relationships with caregivers and positive parenting approaches appear key, but this is also the case for social support from friends or other adults. There was less evidence available for school factors associated with resilient outcomes for children. A safe and orderly school environment, positive relationships with teachers and student academic engagement appear to be associated with resilient outcomes in the context of poverty. Few studies addressed community factors, but community cohesion and links with cultural identity, including spiritual beliefs, also appear to be associated with resilience in children.

As noted above, there was significant heterogeneity in the measurement approaches, categorisation of resilience and the resilience factors investigated in the reviewed literature. This meant we were not able to summarise the papers quantitatively to compare effect sizes for variables associated with resilient outcomes using meta-analyses.

Limitations of this review include the breadth of resilience research, which required the exclusion of a large number of papers after a brief review. In hindsight, some relevant papers may have been missed due to the databases chosen—a recent paper by Bramer *et al*⁷⁶ suggests that a combination of Embase, MEDLINE, Web of Science Core Collection and Google Scholar performs best for systematic reviews. We did not have sufficient funding to include papers published in languages other than English. The studies on the whole were moderate to high quality; however, many scored poorly in terms of employing culturally safe and participatory study methods. This may introduce bias in our understanding of the resilience factors that are important for children from migrant or minority backgrounds. None of the included studies reported data for Indigenous or refugee children. This may be because these children did not participate or were not identified separately in the included studies. Given the greater burden of childhood trauma and social adversity experienced by Indigenous^{77 78} and refugee children,⁷⁹ this is a major gap in the literature.

The review findings provide a starting point for focusing research efforts on a broader conceptualisation of resilience that includes individual factors and factors in the socioecological context surrounding the child. Developmentally, this broader understanding of resilience is particularly important for children, who have a limited capacity to shape their world compared with adolescents and adults. It also highlights aspects of resilience that will be important to consider when attempting to improve child resilience in clinical or educational settings. Two recent systematic reviews conducting meta analyses of resilience interventions, one focused on adults and one on school-based interventions, concluded that resilience interventions, particularly those including cognitive behavioural therapy, appeared to be beneficial, at least in the short term.^{37 80} However, considerable heterogeneity across studies, weak methodologies, lack of longitudinal data and small sample sizes were among the limitations noted. Given the breadth of factors identified as contributing to child resilience in the current review, it appears likely that resilience interventions will be most effective where both individual and broader relevant socioecological aspects are addressed and multilevel approaches are taken.^{37 41 81 82}

It is imperative to investigate resilience and develop interventions to build child resilience in a way that embraces more vulnerable, harder-to-reach populations, often not captured using standard research methods. It is also vital that all the relevant domains of resilience including self, family, school, social and neighbourhood/community strengths are incorporated. The development of a measure of child resilience factors that is relevant and appropriate for children who are more vulnerable to disadvantage and social adversity will have several advantages. It will support consistency across studies and contexts and the development and evaluation of intervention programmes, and it will improve assessment of child

outcomes in clinical, research and educational settings for families.

While there is huge variation in the type and severity of adversity that children may experience, there are a number of resilience factors associated with positive child outcomes across different contexts and exposures. Such factors provide an important starting point for effective public health interventions to promote resilience and to prevent or ameliorate the immediate and long-term impacts of social adversity on children.

Author affiliations

¹Intergenerational Health, Murdoch Children's Research Institute, Melbourne, Victoria, Australia

²Department of Pediatrics, University of Melbourne, Melbourne, Victoria, Australia

³Department of General Practice and Primary Health Care, University of Melbourne, Carlton, Victoria, Australia

⁴Department of Community Health Sciences and Department of Psychiatry, University of Manitoba, Winnipeg, Manitoba, Canada

⁵Departments of Psychiatry and Behavioural Neurosciences, and of Pediatrics, McMaster University, Hamilton, Ontario, Canada

⁶Orygen, The National Centre of Excellence in Youth Mental Health, Melbourne, Victoria, Australia

Contributors DG, ER, SM, RG and SJB devised the search protocol; EB, DG, ER and SM executed the protocol; SM and ER completed the critical appraisal; DG conducted the analysis and wrote the paper with input from RG, TOA, HM, HH and SJB. All authors have contributed to and approved the final manuscript.

Funding Funding for this study was provided by the Australian National Health and Medical Research Council Project Grant (#1064061) and Myer Family Foundation philanthropic funding. Murdoch Children's Research Institute is supported by the Victorian Government's Operational Infrastructure programme. HM is supported by the Chedoke Health Chair in Psychiatry. TOA is supported by a Canadian Institutes of Health Research (CIHR) New Investigator Award and a CIHR Foundations Grant. SJB is supported by a NHMRC Research Fellowship. RG is supported by a NHMRC Career Development Fellowship.

Disclaimer The funding bodies had no role in the study design, collection, analysis or interpretation of the data, writing the manuscript or the decision to submit the paper for publication.

Competing interests None declared.

Patient consent for publication Not required.

Provenance and peer review Not commissioned; externally peer reviewed.

Data sharing statement The full results of the quality appraisal are available upon request.

Open access This is an open access article distributed in accordance with the Creative Commons Attribution Non Commercial (CC BY-NC 4.0) license, which permits others to distribute, remix, adapt, build upon this work non-commercially, and license their derivative works on different terms, provided the original work is properly cited, appropriate credit is given, any changes made indicated, and the use is non-commercial. See: <http://creativecommons.org/licenses/by-nc/4.0/>.

REFERENCES

- Gartland D, Woolhouse H, Giallo R, *et al*. Vulnerability to intimate partner violence and poor mental health in the first 4-year postpartum among mothers reporting childhood abuse: an Australian pregnancy cohort study. *Arch Womens Ment Health* 2016;19:1091–100.
- Afifi TO, MacMillan HL, Boyle M, *et al*. Child abuse and mental disorders in Canada. *Can Med Assoc J* 2014;186:E324–32.
- Garcia-Moreno C, Jansen HA, Ellsberg M, *et al*. Prevalence of intimate partner violence: findings from the WHO multi-country study on women's health and domestic violence. *Lancet* 2006;368:1260–9.
- Gartland D, Woolhouse H, Mensah FK, *et al*. The case for early intervention to reduce the impact of intimate partner abuse on child

- outcomes: results of an Australian cohort of first-time mothers. *Birth* 2014;41:374–83.
5. United Nations Children's Fund. A child is a child: Protecting children on the move from violence, abuse and exploitation. 2017.
 6. Phillips B, Miranti R, Vidyattama Y, et al. *Social exclusion and disadvantage in Australia*. University of Canberra, ACT: The National Centre for Social and Economic Modelling, 2013.
 7. Weetra D, Glover K, Buckskin M, et al. Stressful events, social health issues and psychological distress in Aboriginal women having a baby in South Australia: implications for antenatal care. *BMC Pregnancy Childbirth* 2016;16:88.
 8. Askew DA, Schluter PJ, Spurling GK, et al. Urban Aboriginal and Torres Strait Islander children's exposure to stressful events: a cross-sectional study. *Med J Aust* 2013;199:42–5.
 9. Marmot M. Social determinants of health inequalities. *Lancet* 2005;365:1099–104.
 10. Murali V, Oyebo F. Poverty, social inequality and mental health. *Advances in Psychiatric Treatment* 2004;10:216–24.
 11. Mensah FK, Kiernan KE. Maternal general health and children's cognitive development and behaviour in the early years: findings from the millennium cohort study. *Child Care Health Dev* 2011;37:44–54.
 12. Elgar FJ, McGrath PJ, Waschbusch DA, et al. Mutual influences on maternal depression and child adjustment problems. *Clin Psychol Rev* 2004;24:441–59.
 13. Kitzmann KM, Gaylord NK, Holt AR, et al. Child witnesses to domestic violence: a meta-analytic review. *J Consult Clin Psychol* 2003;71:339–52.
 14. Deave T, Heron J, Evans J, et al. The impact of maternal depression in pregnancy on early child development. *BJOG* 2008;115:1043–51.
 15. O'Connor TG, Heron J, Golding J, et al. Maternal antenatal anxiety and children's behavioural/emotional problems at 4 years. *British Journal of Psychiatry* 2002;180:502–8.
 16. Young JC, Widom CS. Long-term effects of child abuse and neglect on emotion processing in adulthood. *Child Abuse Negl* 2014;38:1369–81.
 17. Afifi TO, Boman J, Fleisher W, et al. The relationship between child abuse, parental divorce, and lifetime mental disorders and suicidality in a nationally representative adult sample. *Child Abuse Negl* 2009;33:139–47.
 18. Webb RT, Antonsen S, Carr MJ, et al. Self-harm and violent criminality among young people who experienced trauma-related hospital admission during childhood: a Danish national cohort study. *Lancet Public Health* 2017;2:e314–22.
 19. Herrenkohl TI, Hong S, Klika JB, et al. Developmental impacts of child abuse and neglect related to adult mental health, substance use, and physical health. *J Fam Violence* 2013;28:191–9.
 20. Hughes K, Bellis MA, Hardcastle KA, et al. The effect of multiple adverse childhood experiences on health: a systematic review and meta-analysis. *Lancet Public Health* 2017;2:e356–66.
 21. Dubow EF, Ippolito MF. Effects of poverty and quality of the home environment on changes in the academic and behavioral adjustment of elementary school-age children. *J Clin Child Psychol* 1994;23:401–12.
 22. Rutter M. Poverty and child mental health: natural experiments and social causation. *JAMA* 2003;290:2063–4.
 23. Hawkins JD, Van Horn ML, Arthur MW. Community variation in risk and protective factors and substance use outcomes. *Prev Sci* 2004;5:213–20.
 24. Masten AS. Resilience in children threatened by extreme adversity: frameworks for research, practice, and translational synergy. *Dev Psychopathol* 2011;23:493–506.
 25. Cicchetti D. Annual Research Review: Resilient functioning in maltreated children—past, present, and future perspectives. *J Child Psychol Psychiatry* 2013;54:402–22.
 26. Luthar SS, Cicchetti D, Becker B. The construct of resilience: a critical evaluation and guidelines for future work. *Child Dev* 2000;71:543–62.
 27. Herrman H, Stewart DE, Diaz-Granados N, et al. What is resilience? *Can J Psychiatry* 2011;56:258–65.
 28. Masten AS, Tellegen A. Resilience in developmental psychopathology: contributions of the Project Competence Longitudinal Study. *Dev Psychopathol* 2012;24:345–61.
 29. Hooberman J, Rosenfeld B, Rasmussen A, et al. Resilience in trauma-exposed refugees: the moderating effect of coping style on resilience variables. *Am J Orthopsychiatry* 2010;80:557–63.
 30. Bowes L, Maughan B, Caspi A, et al. Families promote emotional and behavioural resilience to bullying: evidence of an environmental effect. *J Child Psychol Psychiatry* 2010;51:809–17.
 31. Ben-David V, Jonson-Reid M. Resilience among adult survivors of childhood neglect: a missing piece in the resilience literature. *Child Youth Serv Rev* 2017;78:93–103.
 32. Marriott C, Hamilton-Giachritsis C, Harrop C. Factors promoting resilience following childhood sexual abuse: a structured, narrative review of the literature. *Child Abuse Review* 2014;23:17–34.
 33. Fantuzzo J, LeBoeuf W, Rouse H, et al. Academic achievement of African American boys: a city-wide, community-based investigation of risk and resilience. *J Sch Psychol* 2012;50:559–79.
 34. Newbold B, Chung K, Hong E. *Resilience among single adult female refugees in Hamilton*. Ontario, 2013:65–75.
 35. Cosco TD, Kaushal A, Hardy R, et al. Operationalising resilience in longitudinal studies: a systematic review of methodological approaches. *J Epidemiol Community Health* 2017;71:98–104.
 36. Siriwardhana C, Ali SS, Roberts B, et al. A systematic review of resilience and mental health outcomes of conflict-driven adult forced migrants. *Confl Health* 2014;8:13.
 37. Dray J, Bowman J, Campbell E, et al. Systematic review of universal resilience-focused interventions targeting child and adolescent mental health in the school setting. *J Am Acad Child Adolesc Psychiatry* 2017;56:813–24.
 38. Heckmen JJ. *The case for investing in disadvantaged young children in big ideas for children: investing in our nation's future*. Washington, DC: First Focus, 2008.
 39. Wise S, da Silva L, Webster E, et al. *The efficacy of early childhood interventions*. Melbourne: Australian Institute of Family Studies and Melbourne Institute of Applied Economic and Social Research, 2006.
 40. Braveman P, Barclay C. Health disparities beginning in childhood: a life-course perspective. *Pediatrics* 2009;124:S163–75.
 41. Yoshikawa H, Whipps MD, Rojas NM. Commentary: new directions in developmentally informed intervention research for vulnerable populations. *Child Dev* 2017;88:459–65.
 42. Kinner SA, Borschmann R. Inequality and intergenerational transmission of complex adversity. *Lancet Public Health* 2017;2:e342–3.
 43. Health Evidence Bulletins Wales. Questions to assist with the critical appraisal of a systematic review. <http://heb.wcf.ac.uk/projectmethod/appendix5.htm2015>
 44. Riggs E, Gussy M, Gibbs L, et al. Assessing the cultural competence of oral health research conducted with migrant children. *Community Dent Oral Epidemiol* 2014;42:43–52.
 45. White K, Haas JS, Williams DR. Elucidating the role of place in health care disparities: the example of racial/ethnic residential segregation. *Health Serv Res* 2012;47:1278–99.
 46. In: Wrench J, Rea A, Ouali N, eds. *Migrants, ethnic minorities and the labour market: integration and exclusion in Europe*. 2nd edn: Springer, 2016.
 47. Bagci SC, Rutland A, Kumashiro M, et al. Are minority status children's cross-ethnic friendships beneficial in a multiethnic context? *Br J Dev Psychol* 2014;32:107–15.
 48. Cortina MA, Fazel M, Hlungwani TM, et al. Childhood psychological problems in school settings in rural Southern Africa. *PLoS One* 2013;8:e65041.
 49. Bartko WT, Eccles JS. Adolescent participation in structured and unstructured activities: a person-oriented analysis. *J Youth Adolesc* 2003;32:233–41.
 50. Somchit S, Sriyaporn P. The relationships among resilience factors, perception of adversities, negative behavior, and academic achievement of 4th- to 6th-grade children in Thad-Thong, Chonburi, Thailand. *J Pediatr Nurs* 2004;19:294–303.
 51. Lagasse LL, Hammond J, Liu J, et al. Violence and delinquency, early onset drug use, and psychopathology in drug-exposed youth at 11 years. *Ann N Y Acad Sci* 2006;1094:313–8.
 52. Halevi G, Djalovski A, Vengrober A, et al. Risk and resilience trajectories in war-exposed children across the first decade of life. *J Child Psychol Psychiatry* 2016;57:1183–93.
 53. Brown J, Barbarin O, Scott K. Socioemotional trajectories in black boys between kindergarten and the fifth grade: the role of cognitive skills and family in promoting resiliency. *Am J Orthopsychiatry* 2013;83:176–84.
 54. Cicchetti D, Rogosch FA. Personality, adrenal steroid hormones, and resilience in maltreated children: a multilevel perspective. *Dev Psychopathol* 2007;19:787–809.
 55. Flores E, Cicchetti D, Rogosch FA. Predictors of resilience in maltreated and nonmaltreated latino children. *Dev Psychol* 2005;41:338–51.
 56. Graham-Bermann SA, Gruber G, Howell KH, et al. Factors discriminating among profiles of resilience and psychopathology in children exposed to intimate partner violence (IPV). *Child Abuse Negl* 2009;33:648–60.

57. Lin KK, Sandler IN, Ayers TS, et al. Resilience in parentally bereaved children and adolescents seeking preventive services. *J Clin Child Adolesc Psychol* 2004;33:673–83.
58. Flouri E, Midouhas E, Joshi H. Family poverty and trajectories of children's emotional and behavioural problems: the moderating roles of self-regulation and verbal cognitive ability. *J Abnorm Child Psychol* 2014;42:1043–56.
59. Jaffee SR, Caspi A, Moffitt TE, et al. Individual, family, and neighborhood factors distinguish resilient from non-resilient maltreated children: a cumulative stressors model. *Child Abuse Negl* 2007;31:231–53.
60. Borman GD, Overman LT. Academic resilience in mathematics among poor and minority students. *Elem Sch J* 2004;104:177–95.
61. Myers BJ, Mackintosh VH, Kuznetsova MI, et al. Relationship processes and resilience in children with incarcerated parents: III. Teasing, bullying, and emotion regulation in children of incarcerated mothers. *Monogr Soc Res Child Dev* 2013;78:26–40.
62. Anthony EK. Cluster profiles of youths living in Urban poverty: factors affecting risk and resilience. *Soc Work Res* 2008;32:6–17.
63. Horner SD, Rew L, Brown A. Risk-taking behaviors engaged in by early adolescents while on school property. *Issues Compr Pediatr Nurs* 2012;35:90–110.
64. Yoon S. Fostering resilient development: protective factors underlying externalizing trajectories of maltreated children. *J Child Fam Stud* 2018;27:443–52.
65. Downie JM, Hay DA, Horner BJ, et al. Children living with their grandparents: resilience and wellbeing. *Int J Soc Welf* 2010;19:8–22.
66. Ew N, Chan CC, Lai MK. Hope and life satisfaction among underprivileged children in hong kong: the mediating role of perceived community support. *J Community Psychol* 2014;42:352–64.
67. Coohey C, Renner LM, Hua L, et al. Academic achievement despite child maltreatment: a longitudinal study. *Child Abuse Negl* 2011;35:688–99.
68. Ebersöhn L, Eloff I, Finestone M, et al. Telling stories and adding scores: measuring resilience in young children affected by maternal HIV and AIDS. *Afr J AIDS Res* 2015;14:219–27.
69. Poehlmann J, Eddy JM, Dallaire DH, et al. Relationship processes and resilience in children with incarcerated parents. *Monogr Soc Res Child Dev* 2013;78:7–25.
70. Bell T, Romano E, Flynn RJ. Multilevel correlates of behavioral resilience among children in child welfare. *Child Abuse Negl* 2013;37:1007–20.
71. Jones JM. Exposure to chronic community violence: resilience in African American children. *J Black Psychol* 2007;33:125–49.
72. Ramos-Olazagasti MA, Shrout PE, Yoshikawa H, et al. Contextual risk and promotive processes in puerto rican youths' internalizing trajectories in puerto rico and New York. *Dev Psychopathol* 2013;25:755–71.
73. Walsh WA, Dawson J, Mattingly MJ. How are we measuring resilience following childhood maltreatment? Is the research adequate and consistent? What is the impact on research, practice, and policy? *Trauma Violence Abuse* 2010;11:27–41.
74. Windle G, Bennett KM, Noyes J. A methodological review of resilience measurement scales. *Health Qual Life Outcomes* 2011;9:8–2554.
75. Ungar M. A constructionist discourse on resilience multiple contexts, multiple realities among at-risk children and youth. *Youth Soc* 2004;35:341–65.
76. Bramer WM, Rethlefsen ML, Kleijnen J, et al. Optimal database combinations for literature searches in systematic reviews: a prospective exploratory study. *Syst Rev* 2017;6:245.
77. Welfare AloHa. *The health and welfare of australia's Aboriginal and Torres Strait Islander peoples: Welfare AloHa*, 2015.
78. Mitrou F, Cooke M, Lawrence D, et al. Gaps in Indigenous disadvantage not closing: a census cohort study of social determinants of health in Australia, Canada, and New Zealand from 1981–2006. *BMC Public Health* 2014;14:1–17.
79. Hodes M, Anagnostopoulos D, Skokauskas N. Challenges and opportunities in refugee mental health: clinical, service, and research considerations. *Eur Child Adolesc Psychiatry* 2018;27:385–8.
80. Joyce S, Shand F, Tighe J, et al. Road to resilience: a systematic review and meta-analysis of resilience training programmes and interventions. *BMJ Open* 2018;8:e017858.
81. Cahill H, Beadle S, Farrelly R, et al. *Building resilience in children and young people: a literature review for the department of education and early childhood development youth research centre, melbourne graduate school of education*. Melbourne: University of Melbourne, 2017.
82. Zimmerman RS. Importance of resilience research and multi-level interventions. *Soc Sci Med* 2017;190:275–7.
83. Dubowitz H, Thompson R, Proctor L, et al. Adversity, maltreatment, and resilience in young children. *Acad Pediatr* 2016;16:233–9.
84. Manly JT, Lynch M, Oshri A, et al. The impact of neglect on initial adaptation to school. *Child Maltreat* 2013;18:155–70.