Correction: *RADAR study: protocol for an observational cohort study to identify early warning signals on the pathways to alcohol use disorder*

Slade T, Swift W, Mewton L, *et al.* RADAR study: protocol for an observational cohort study to identify early warning signals on the pathways to alcohol use disorder *BMJ Open* 2017;7:e018256. DOI:10.1136/bmjopen-2017-018256.

This article was previously published with an error.

In the methods and analysis section the text reads:

Participants will be eligible for RADAR if and when they report either: (1) frequent drinking (defined as consuming any quantity of alcohol at least 1–2 days/week over a 12-month period), or (2) semifrequent binge drinking (defined as consuming more than 4 standard drinks, ie, 40 g alcohol, in a single session at least 3–4 days/month over a 12-month period).

However, the correct text should read:

Participants will be eligible for RADAR if and when they report either: (1) frequent drinking (defined as consuming any quantity of alcohol at least 1–2 days/week over a 12-month period), or (2) semifrequent binge drinking (defined as consuming more than 4 standard drinks, ie, 40 g alcohol, in a single session at least 2–3 days/month over a 12-month period).

Open access This is an open access article distributed in accordance with the Creative Commons Attribution Non Commercial (CC BY-NC 4.0) license, which permits others to distribute, remix, adapt, build upon this work non-commercially, and license their derivative works on different terms, provided the original work is properly cited, appropriate credit is given, any changes made indicated, and the use is non-commercial. See: <http://creativecommons.org/licenses/by-nc/4.0/>.

© Author(s) (or their employer(s)) 2019. Re-use permitted under CC BY-NC. No commercial re-use. See rights and permissions. Published by BMJ.

BMJ Open 2019;9:e018256corr1. doi:10.1136/bmjopen-2017-018256corr1